

## Set Theory Definitions

---

### Set Membership, Equality, and Subsets

An element of a set is an object directly contained within that set. For example,  $1 \in \{1, 2, 3\}$  and  $\emptyset \in \{\emptyset\}$ , but  $1 \notin \emptyset$  and  $1 \notin \{2, 3\}$ . Note that  $1 \notin \{\{1\}\}$ , and  $\{1\} \notin \{1\}$ , but  $\{1\} \in \{\{1\}\}$ .

Two sets are equal if they contain the same elements. For example, we have that  $\{1, 2\} = \{2, 1\}$  and that  $\{\emptyset\} = \{\emptyset\}$ . However,  $\{\emptyset\} \neq \{\{\emptyset\}\}$ , because each set contains an element the other does not. A set and a non-set are never equal; in particular, this means  $x \neq \{x\}$  for any  $x$ .

A set  $A$  is a subset of a set  $B$  (denoted  $A \subseteq B$ ) if every element of  $A$  is also an element of  $B$ :

$$\mathbb{N} \subseteq \mathbb{Z} \quad \{1, 2, 3\} \subseteq \{1, 2, 3, 4\} \quad \{1\} \subseteq \{1, \{1\}, \{\{1\}\}\}$$

### Set Operations

The set  $\{x \mid \text{some property of } x\}$  is the set of all objects  $x$  that satisfy the given property. Formally, we have that  $w \in \{x \mid \text{some property of } x\}$  if and only if the specified property holds for  $w$ .

The set  $A \cup B$  is the set  $\{x \mid x \in A \text{ or } x \in B\}$ . Equivalently,  $x \in A \cup B$  if and only if  $x \in A$  or  $x \in B$ .

The set  $A \cap B$  is the set  $\{x \mid x \in A \text{ and } x \in B\}$ . Equivalently,  $x \in A \cap B$  precisely if  $x \in A$  and  $x \in B$ .

The set  $A - B$  is the set  $\{x \mid x \in A \text{ and } x \notin B\}$ . This set is also sometimes denoted  $A \setminus B$ .

The set  $A \Delta B$  is the set  $\{x \mid \text{exactly one of } x \in A \text{ and } x \in B \text{ is true}\}$ .

### Power Sets

The power set of a set  $S$ , denoted  $\wp(S)$ , is the set of all subsets of  $S$ . Using set-builder notation, this is the set  $\wp(S) = \{U \mid U \subseteq S\}$ . Cantor's Theorem states that  $|S| < |\wp(S)|$  for every set  $S$ .

### Special Sets

The set  $\emptyset = \{\}$  is the empty set containing no elements.

The set  $\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$  is the set of all natural numbers. We treat 0 as a natural number.

The set  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$  is the set of all integers.

The set  $\mathbb{R}$  consists of all the real numbers. The set  $\mathbb{Q}$  consists of all rational numbers.

### Cardinality

The cardinality of a finite set  $S$  (denoted  $|S|$ ) is the natural number equal to the number of elements in that set. The cardinality of  $\mathbb{N}$  (denoted  $|\mathbb{N}|$ ) is  $\aleph_0$  (pronounced "aleph-nought"). Two sets have the same cardinality if there is a way of pairing up each element of the two sets such that every element of each set is paired with exactly one element of the other set.