

Section 1: Analytic Probability

Problems by three through six by Chris Piech. Rest by past CS109 teaching teams

Overview of Section Materials

The warmup questions provided will help students practice concepts introduced in lectures. The section problems are meant to apply these concepts in more complex scenarios similar to what you will see in problem sets and exams.

Warmups

1. Fish Pond

Suppose there are 7 blue fish, 4 red fish, and 8 green fish in a large fishing tank. You drop a net into it and end up with 2 fish. What is the probability you get 2 blue fish?

2. Axioms of Probability

Decide whether each of the three statements below is true or false:

- $P(A) + P(A^C) = 1$. Recall that A^C means A "complement" or "not" A
- $P(A \cap B) + P(A \cap B^C) = 1$. Recall that \cap means "and"
- If $P(A) = 0.4$ and $P(B) = 0.6$ then it must be the case that $A = B^C$

3. Conditional Probability

What is the difference between these two terms $P(B|A)$ and $P(A \cap B)$? Imagine that B is the event that a student "correctly answer a multiple choice question" and A is the event that the same student "guesses randomly". Provide an explanation as well as a mathematical relationship between the two.

Problems

4. The Birthday Problem

When solving a counting problem, it can often be useful to come up with a generative process, a series of steps that "generates" examples. A correct generative process to count the elements of set A will (1) *generate every element of A* and (2) *not generate any element of A more than once*. If our process has the added property that (3) *any given step always has the same number of possible outcomes*, then we can use the product rule of counting.

Example: Say we want to count the number of ways to roll two (distinct) dice where one die is even and one die is odd. Our process could be: (1) roll the first die and note if the value is even or odd, then (2) count the number of ways the second die can be rolled for a value of the opposite parity. Since the first step has 6 options and the second step has 3 options regardless of the outcome of the first step, the number of possibilities is $6 * 3 = 18$.

Problem: Assume that birthdays happen on any of the 365 days of the year with equal likelihood (we'll ignore leap years).

- What is the probability that of the n people in class, at least two people share the same birthday?
- What is the probability that this class contains exactly one pair of people who share a birthday?

5. Self Driving Car

A self driving car has a 60% belief that there is a motorcycle to its left based on all the information it has received up until this point in time. Then, it receives a new, independent report from its left camera. The camera reports that there is **no** motorcycle. What is the updated belief that there is a motorcycle to the left of the car? The camera is an imperfect instrument. When there is truly no motorcycle, the camera will report “no motorcycle” 90% of the time. When there actually is a motorcycle, the camera will report “no motorcycle” 5% of the time.

Extra Practice

6. Flipping Coins

One thing that students often find tricky when learning combinatorics is how to figure out when a problem involves permutations and when it involves combinations. Naturally, we will look at a problem that can be solved with both approaches. Pay attention to what parts of your solution represent distinct objects and what parts represent indistinct objects.

Problem: We flip a fair coin n times, hoping (for some reason) to get k heads.


- How many ways are there to get exactly k heads? Characterize your answer as a *permutation* of H's and T's.
- For what x and y is your answer to part (a) equal to $\binom{x}{y}$? Why does this *combination* make sense as an answer?
- What is the probability that we get exactly k heads?

7. Counting

The Inclusion Exclusion Principle for three sets is:

$$|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$$

Explain why in terms of a venn-diagram.



8. Combinatorial Proofs

Prove why $\binom{n}{k} = \binom{n}{n-k}$.