

EE270

Large scale matrix computation, optimization and learning

Instructor : Mert Pilanci

Stanford University

Thursday, Jan 23 2020

Randomized Linear Algebra

Lecture 7: Least Squares Optimization and Random Projections

Recap: Johnson Lindenstrauss Lemma

- ▶ Let $\epsilon \in (0, \frac{1}{2})$. Given any set of points $\{x_1, \dots, x_n\}$ in \mathbb{R}^d , there exists a map $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $m = \frac{9 \log(n)}{\epsilon^2 - \epsilon^3}$ such that

$$1 - \epsilon \leq \frac{\|Sx_i - Sx_j\|_2^2}{\|x_i - x_j\|_2^2} \leq 1 + \epsilon$$

Recap: Johnson Lindenstrauss Lemma

- ▶ Let $\epsilon \in (0, \frac{1}{2})$. Given any set of points $\{x_1, \dots, x_n\}$ in \mathbb{R}^d , there exists a map $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $m = \frac{9 \log(n)}{\epsilon^2 - \epsilon^3}$ such that

$$1 - \epsilon \leq \frac{\|Sx_i - Sx_j\|_2^2}{\|x_i - x_j\|_2^2} \leq 1 + \epsilon$$

- ▶ Note that the target dimension m is **independent of the original dimension d** , and depends **only on the number of points n** and the accuracy parameter.

Recap: Johnson Lindenstrauss Lemma

- ▶ Let $\epsilon \in (0, \frac{1}{2})$. Given any set of points $\{x_1, \dots, x_n\}$ in \mathbb{R}^d , there exists a map $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $m = \frac{9 \log(n)}{\epsilon^2 - \epsilon^3}$ such that

$$1 - \epsilon \leq \frac{\|Sx_i - Sx_j\|_2^2}{\|x_i - x_j\|_2^2} \leq 1 + \epsilon$$

- ▶ Note that the target dimension m is **independent of the original dimension d** , and depends **only on the number of points n** and the accuracy parameter.
- ▶ more surprises: picking an $m \times d$ random matrix $S = \frac{1}{\sqrt{m}} G$ with $G_{ij} \sim N(0, 1)$ standard normal works with high probability!

True 'projections': random subspaces also work

- ▶ Pick $S_{(i)}$ uniformly random on the unit sphere
- ▶ Pick $S_{(i+1)}$ uniformly random on the unit sphere and $\perp S_{(i)}, \dots, S_{(1)}$
- ▶ S is a projection matrix, which projects onto a uniformly random subspace

$$\mathbb{P} \left\{ \left| \|Su\|_2 - \sqrt{\frac{m}{d}} \right| > t \right\} \leq 2e^{-\frac{t^2 d}{2}}$$

- ▶ Applying union bound for all points $i, j = 1, \dots, d$ gives a similar result
- ▶ Random i.i.d. S matrices are easier to generate and approximately orthogonal: $\mathbb{E}S^T S = I$

Computationally cheaper random matrices

- ▶ Gaussian $S_{ij} = \frac{1}{\sqrt{m}} N(0, 1)$
- ▶ Rademacher

$$S_{ij} = \begin{cases} +\frac{1}{\sqrt{m}} & \text{with probability } \frac{1}{2} \\ -\frac{1}{\sqrt{m}} & \text{with probability } \frac{1}{2} \end{cases} \quad (1)$$

- ▶ Bernoulli-Rademacher

$$S_{ij} = \begin{cases} +\frac{\sqrt{3}}{\sqrt{m}} & \text{with probability } \frac{1}{2} \\ 0 & \text{with probability } \frac{2}{3} \\ -\frac{\sqrt{3}}{\sqrt{m}} & \text{with probability } \frac{1}{2} \end{cases} \quad (2)$$

- ▶ other sparse matrices (e.g. one non-zero per column)
- ▶ Fourier transform based matrices

Random projection for Approximate Matrix Multiplication

- ▶ Let the approximate product of AB be $C = AS^T SB$

$$\mathbb{P}[\|AB - C\|_F > 3\epsilon\|A\|_F\|B\|_F] \leq \delta$$

- ▶ Follows from JL Moment property
- ▶ $S \in \mathbb{R}^{m \times n} \sim \frac{1}{\sqrt{m}} \times$ random i.i.d. sub-Gaussian, e.g., ± 1 , or $N(0, 1)$ with $m = \frac{c_1}{\epsilon^2} \log \frac{1}{\delta}$
- ▶ $S \in \mathbb{R}^{m \times n} \sim \frac{1}{\sqrt{m}} \times$ CountSketch matrix (one nonzero per column, which is ± 1 at a uniformly random location) with $m = \frac{c_2}{\epsilon^2 \delta}$
- ▶ $S \in \mathbb{R}^{m \times n} \sim \frac{1}{\sqrt{m}} \times$ Fast JL Transform with $m = \frac{c_3}{\epsilon} \log \frac{1}{\delta}$

Random projection for Approximate Matrix Multiplication

- ▶ Let the approximate product of AB be $C = AS^T SB$

$$\mathbb{P}[\|AB - C\|_F > 3\epsilon\|A\|_F\|B\|_F] \leq \delta$$

- ▶ Follows from JL Moment property
- ▶ $S \in \mathbb{R}^{m \times n} \sim \frac{1}{\sqrt{m}} \times$ random i.i.d. sub-Gaussian, e.g., ± 1 , or $N(0, 1)$ with $m = \frac{c_1}{\epsilon^2} \log \frac{1}{\delta}$
- ▶ $S \in \mathbb{R}^{m \times n} \sim \frac{1}{\sqrt{m}} \times$ CountSketch matrix (one nonzero per column, which is ± 1 at a uniformly random location) with $m = \frac{c_2}{\epsilon^2 \delta}$
- ▶ $S \in \mathbb{R}^{m \times n} \sim \frac{1}{\sqrt{m}} \times$ Fast JL Transform with $m = \frac{c_3}{\epsilon} \log \frac{1}{\delta}$
- ▶ Sparse JL and Fast JL are more efficient
- ▶ advantages: doesn't require any knowledge about matrices A and B (**oblivious**)
- ▶ optimal sampling probabilities depend on the column/row norms of A and B

Least Squares Regression

- ▶ Predict the value of a continuous target variable y
 $(a_1, b_1), \dots, (a_n, b_n)$
 $a_i \in \mathbb{R}^d$ and $b_i \in \mathbb{R}$
- ▶ Linear regression $f(a) = x^T a + x_0$

Least Squares Regression

- ▶ Predict the value of a continuous target variable y
 $(a_1, b_1), \dots, (a_n, b_n)$
 $a_i \in \mathbb{R}^d$ and $b_i \in \mathbb{R}$
- ▶ Linear regression $f(a) = x^T a + x_0$
- ▶ Performance measure: minimum sum of squares

$$\min_{x, x_0} \frac{1}{n} \sum_{i=1}^n (b_i - x^T a_i - x_0)^2$$

Least Squares Regression

- ▶ Predict the value of a continuous target variable y
 $(a_1, b_1), \dots, (a_n, b_n)$
 $a_i \in \mathbb{R}^d$ and $b_i \in \mathbb{R}$
- ▶ Linear regression $f(a) = x^T a + x_0$
- ▶ Performance measure: minimum sum of squares

$$\min_{x, x_0} \frac{1}{n} \sum_{i=1}^n (b_i - x^T a_i - x_0)^2$$

- ▶ we can add a regularization term $\lambda \|x\|_2^2$

$$\min_{x, x_0} \frac{1}{n} \sum_{i=1}^n (b_i - x^T a_i - x_0)^2 + \lambda \|x\|_2^2$$

Least Squares Regression

- ▶ Loss function:

$$L(x, x_0) = \frac{1}{n} \sum_{i=1}^n (b_i - x^T a_i - x_0)^2 + \lambda \|x\|_2^2$$

- ▶ $\frac{\partial}{\partial x_0} L(x, x_0) =$

$$\text{optimal } x_0^* = \frac{1}{n} \sum_{i=1}^n (y_i - x^T a_i) = \bar{b} - x^T \bar{a}$$

$$\text{where } \bar{a} = \sum_{i=1}^n a_i \text{ and } \bar{b} = \sum_{i=1}^n b_i$$

- ▶ plugging x_0^* in $L(x, x_0)$

$$L(x, x_0^*) = \frac{1}{n} \sum_{i=1}^n (b_i - \bar{b} - x^T (a_i - \bar{a}))^2 + \lambda \|x\|_2^2$$

Least Squares Regression

- ▶ Loss function:

$$L(x, x_0) = \frac{1}{n} \sum_{i=1}^n (b_i - x^T a_i - x_0)^2 + \lambda \|x\|_2^2$$

- ▶ $\frac{\partial}{\partial x_0} L(x, x_0) =$

$$\text{optimal } x_0^* = \frac{1}{n} \sum_{i=1}^n (y_i - x^T a_i) = \bar{b} - x^T \bar{a}$$

$$\text{where } \bar{a} = \sum_{i=1}^n a_i \text{ and } \bar{b} = \sum_{i=1}^n b_i$$

- ▶ plugging x_0^* in $L(x, x_0)$

$$L(x, x_0^*) = \frac{1}{n} \sum_{i=1}^n (b_i - \bar{b} - x^T (a_i - \bar{a}))^2 + \lambda \|x\|_2^2$$

define centered data $\tilde{a}_i = a_i - \bar{a}$ and $\tilde{b}_i = b_i - \bar{b}$

$$\min_x \|\tilde{A}x - \tilde{b}\|_2^2 + n\lambda \|x\|_2^2$$

Least Squares Regression

- ▶ Loss function:

$$L(x, x_0) = \frac{1}{n} \sum_{i=1}^n (b_i - x^T a_i - x_0)^2 + \lambda \|x\|_2^2$$

- ▶ $\frac{\partial}{\partial x_0} L(x, x_0) =$

$$\text{optimal } x_0^* = \frac{1}{n} \sum_{i=1}^n (y_i - x^T a_i) = \bar{b} - x^T \bar{a}$$

$$\text{where } \bar{a} = \sum_{i=1}^n a_i \text{ and } \bar{b} = \sum_{i=1}^n b_i$$

- ▶ plugging x_0^* in $L(x, x_0)$

$$L(x, x_0^*) = \frac{1}{n} \sum_{i=1}^n (b_i - \bar{b} - x^T (a_i - \bar{a}))^2 + \lambda \|x\|_2^2$$

define centered data $\tilde{a}_i = a_i - \bar{a}$ and $\tilde{b}_i = b_i - \bar{b}$

$$\min_x \|\tilde{A}x - \tilde{b}\|_2^2 + n\lambda \|x\|_2^2$$

$$\frac{\partial}{\partial x} L(x, x_0^*) = 2\tilde{A}^T (\tilde{A}x^* - \tilde{b}) + 2n\lambda x^* = 0$$

$$\text{optimal solution } x^* = (\tilde{A}^T \tilde{A} + n\lambda I)^{-1} \tilde{A}^T \tilde{b}$$

Autoregressive Models

$$b[n] = a[n + 1] \approx \sum_k x_k a[n - k]$$

- ▶ AR(2) model : two non-zero filter coefficients

$$a[n + 1] = -x_0 a[n] - x_1 a[n - 1]$$


and error term $e_n = 0$

- ▶ Example: Sine wave $a[n] = \sin(\alpha n)$ satisfies AR(2) model

Autoregressive models

- ▶ We can predict future values using

$$b[n] = \sum_k a[n - k]x_k$$



Least Squares Problems and Random Projection


- ▶ Given $A \in \mathbb{R}^{n \times d}$ and $b \in \mathbb{R}^d$
find the best linear fit $Ax \approx b$ according to

$$\min_{x \in \mathbb{R}^d} \|Ax - b\|_2^2$$

- ▶ no regularization, i.e., $\lambda = 0$
- ▶ If A is full column rank then
- ▶ $x_{LS} = (A^T A)^{-1} A^T b$

Geometry

$$\min_{x \in \mathbb{R}^d} \|Ax - b\|_2^2$$



Singular Value Decomposition

- ▶ Every $A \in \mathbb{R}^{n \times n}$ has a singular value decomposition

$$A = U \Sigma V^T$$

where $U \in \mathbb{R}^{n \times n}$ has orthonormal columns

Σ is diagonal with non-increasing non negative entries

V^T has orthonormal rows

- ▶ Pseudoinverse $A^\dagger = V \Sigma^{-1} U^T$

- ▶ Least Square solution

$$x_{LS} = (A^T A)^{-1} A^T b = A^\dagger b = V \Sigma^{-1} U^T b$$

Classical Methods for Least Squares

▶ **Direct methods**

- ▶ Cholesky decomposition: Form $A^T A$ and decompose $A^T A = R^T R$ where R is upper triangular. Solve normal equations $(A^T A)^{-1} = (R^T R)^{-1} A^T b$
 - ▶ QR decomposition: $A = QR$, solve $Rx = Q^T b$
 - ▶ Singular Value Decomposition: $x_{LS} = V \Sigma^{-1} U^T b$
- Direct methods have typically $O(nd^2)$ complexity

▶ **Indirect methods**

- ▶ Gradient descent with momentum (Chebyshev iteration)
- ▶ Conjugate Gradient
- ▶ Other iterative methods

Indirect methods have typically $O(\sqrt{\kappa}nd)$ complexity, where κ is the condition number

Faster Least Squares Optimization: Random Projection

- ▶ **Left-sketching**

Form SA and Sb where $S \in \mathbb{R}^{m \times n}$ is a random projection matrix

- ▶ Solve the smaller problem

$$\min_{x \in \mathbb{R}^d} \|SAx - Sb\|_2^2$$

- ▶ using any classical method.

Direct method complexity md^2

Faster Least Squares Optimization: Random Projection

- ▶ **Left-sketching**

Form SA and Sb where $S \in \mathbb{R}^{m \times n}$ is a random projection matrix

- ▶ Solve the smaller problem

$$\min_{x \in \mathbb{R}^d} \|SAx - Sb\|_2^2$$

- ▶ using any classical method.

Direct method complexity md^2

Approximation Result

- ▶ Let $S \in \mathbb{R}^{m \times d}$ be a Johnson-Lindenstrauss Embedding

$$x_{LS} = \arg \min_{x \in \mathbb{R}^d} \underbrace{\|Ax - b\|_2^2}_{f(x)}$$

$$\tilde{x} = \arg \min_{x \in \mathbb{R}^d} \|SAx - Sb\|_2^2$$

- ▶ If $m \geq \text{constant} \times \frac{\text{rank}(A)}{\epsilon^2}$ then,
- ▶ $f(x_{LS}) \leq f(\tilde{x}) \leq (1 + \epsilon^2)f(x_{LS})$
- ▶ $\|A(x_{LS} - \tilde{x})\|_2^2 \leq \epsilon^2$ with high probability

Gaussian Sketch

- ▶ Let S be $\frac{1}{m} \times$ i.i.d. Gaussian. $\mathbb{E}[S^T S] = I$

$$\tilde{x} = \arg \min_{x \in \mathbb{R}^d} \|SAx - Sb\|_2^2$$

- ▶ Is $\mathbb{E}[\tilde{x}]$ equal to x_{LS} ?

Questions?