

EE276: Homework #1

Due on Friday Jan 19, 5pm - Gradescope entry code: 2P885N

Note: Mutual information (denoted $I(X;Y)$) will be covered in the Tuesday, Jan 16th lecture. You only need this concept for three sub-parts of this homework.

1. **Example of joint entropy.** Let $p(x, y)$ be given by

		Y	
		0	1
X	0	$\frac{1}{3}$	$\frac{1}{3}$
	1	0	$\frac{1}{3}$


Find

- (a) $H(X), H(Y)$.
- (b) $H(X | Y), H(Y | X)$.
- (c) $H(X, Y)$.
- (d) $H(Y) - H(Y | X)$.
- (e) $I(X; Y)$.
- (f) Draw a Venn diagram for the quantities in (a) through (e).


2. **Entropy of Hamming Code.**

Hamming code is a simple error-correcting code that can correct up to one error in a sequence of bits. Now consider information bits $X_1, X_2, X_3, X_4 \in \{0, 1\}$ chosen uniformly at random, together with check bits X_5, X_6, X_7 chosen to make the parity of the circles even.


(eg: $X_1 + X_2 + X_4 + X_7 = 0 \pmod 2$)



Thus, for example,



becomes



That is, 1011 becomes 1011010.

(a) What is the entropy $H(X_1, X_2, \dots, X_7)$ of $\mathbf{X} := (X_1, \dots, X_7)$?

Now we make an error (or not) in one of the bits (or none). Let $\mathbf{Y} = \mathbf{X} \oplus \mathbf{e}$, where \mathbf{e} is equally likely to be $(1, 0, \dots, 0), (0, 1, 0, \dots, 0), \dots, (0, 0, \dots, 0, 1)$, or $(0, 0, \dots, 0)$, and \mathbf{e} is independent of \mathbf{X} .

(b) Show that one can recover the message \mathbf{X} perfectly from \mathbf{Y} . (Please provide a justification, detailed proof not required.)

(c) What is $H(\mathbf{X}|\mathbf{Y})$?

(d) What is $I(\mathbf{X}; \mathbf{Y})$?

(e) What is the entropy of \mathbf{Y} ?

3. **Entropy of functions of a random variable.** Let X be a discrete random variable. Show that the entropy of a function of X is less than or equal to the entropy of X by justifying the following steps:

$$H(X, g(X)) \stackrel{(a)}{=} H(X) + H(g(X) | X) \quad (1)$$

$$\stackrel{(b)}{=} H(X); \quad (2)$$

$$H(X, g(X)) \stackrel{(c)}{=} H(g(X)) + H(X | g(X)) \quad (3)$$

$$\stackrel{(d)}{\geq} H(g(X)). \quad (4)$$

Thus $H(g(X)) \leq H(X)$.

4. **Coin flips.** A fair coin is flipped until the first head occurs. Let X denote the number of flips required.

(a) Find the entropy $H(X)$ in bits. The following expressions may be useful:

$$\sum_{n=0}^{\infty} r^n = \frac{1}{1-r}, \quad \sum_{n=0}^{\infty} nr^n = \frac{r}{(1-r)^2}.$$

(b) A random variable X is drawn according to this distribution. Construct an “efficient” sequence of yes-no questions of the form, “Is X contained in the set S ?” that determine the value of X . Compare $H(X)$ to the expected number of questions required to determine X .

5. **Minimum entropy.** In the following, we use $H(p_1, \dots, p_n) \equiv H(\mathbf{p})$ to denote the entropy $H(X)$ of a random variable X with alphabet $\mathcal{X} := \{1, \dots, n\}$, i.e.,

$$H(X) = - \sum_{i=1}^n p_i \log(p_i).$$

What is the minimum value of $H(p_1, \dots, p_n) = H(\mathbf{p})$ as \mathbf{p} ranges over the set of n -dimensional probability vectors? Find all \mathbf{p} 's which achieve this minimum.

6. **Drawing with and without replacement.** An urn contains r red, w white, and b black balls. Suppose we draw $k \geq 2$ balls from the urn. Let $\mathbf{X} := (X_1, \dots, X_k)$, where X_i is the color of the i th ball drawn. Is the entropy of \mathbf{X} larger when the balls are drawn from the urn with replacement or without replacement? Set it up and show why. (There is both a hard way and a relatively simple way to do this.)

7. **Infinite entropy. [Bonus]**

This problem shows that the entropy of a discrete random variable can be infinite. (In this question you can take \log as the natural logarithm for simplicity.)

(a) Let $A = \sum_{n=2}^{\infty} (n \log^2 n)^{-1}$. Show that A is finite by bounding the infinite sum by the integral of $(x \log^2 x)^{-1}$.

(b) Show that the integer-valued random variable X distributed as:
 $P(X = n) = (An \log^2 n)^{-1}$ for $n = 2, 3, \dots$ has entropy $H(X)$ given by:

$$H(X) = \log A + \sum_{n=2}^{\infty} \frac{1}{An \log n} + \sum_{n=2}^{\infty} \frac{2 \log \log n}{An \log^2 n}$$

(c) Show that the entropy $H(X) = +\infty$ (by showing that the sum $\sum_{n=2}^{\infty} \frac{1}{n \log n}$ diverges).