

The MicroRNA Cluster miR-106b~25 Regulates Adult Neural Stem/Progenitor Cell Proliferation and Neuronal Differentiation

Jamie O. Brett¹, Valérie M. Renault¹, Victoria A. Rafalski^{1,2}, Ashley E. Webb¹, Anne Brunet^{1,2}

¹Department of Genetics; Stanford University School of Medicine; Stanford, CA 94305; USA

²Neurosciences Program; Stanford University School of Medicine; Stanford, CA 94305; USA

Key words: aging, neural stem cells, microRNAs, FoxO transcription factors, insulin signaling, neuronal differentiation

Received: 2/15/11; **Accepted:** 2/19/11; **Published:** 2/20/11

Corresponding author: Anne Brunet, PhD; **Email:** anne.brunet@stanford.edu

© Brett et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract: In adult mammals, neural stem cells (NSCs) generate new neurons that are important for specific types of learning and memory. Controlling adult NSC number and function is fundamental for preserving the stem cell pool and ensuring proper levels of neurogenesis throughout life. Here we study the importance of the microRNA gene cluster miR-106b~25 (miR-106b, miR-93, and miR-25) in primary cultures of neural stem/progenitor cells (NSPCs) isolated from adult mice. We find that knocking down miR-25 decreases NSPC proliferation, whereas ectopically expressing miR-25 promotes NSPC proliferation. Expressing the entire miR-106b~25 cluster in NSPCs also increases their ability to generate new neurons. Interestingly, miR-25 has a number of potential target mRNAs involved in insulin/insulin-like growth factor-1 (IGF) signaling, a pathway implicated in aging. Furthermore, the regulatory region of miR-106b~25 is bound by FoxO3, a member of the FoxO family of transcription factors that maintains adult stem cells and extends lifespan downstream of insulin/IGF signaling. These results suggest that miR-106b~25 regulates NSPC function and is part of a network involving the insulin/IGF-FoxO pathway, which may have important implications for the homeostasis of the NSC pool during aging.

INTRODUCTION

New neurons are generated in the mammalian brain throughout adult life. Slowly dividing and self-renewing neural stem cells (NSCs) are present in the subventricular zone (SVZ) of the lateral ventricles and in the subgranular zone (SGZ) of the hippocampal dentate gyrus. NSCs generate rapidly proliferating neural progenitor cells that ultimately differentiate to produce thousands of new neurons each day in adult rats [1]. The progeny of SVZ NSCs migrate to the olfactory bulb where they mature into inhibitory interneurons with roles in olfactory learning and memory [2]. SGZ NSCs produce excitatory neurons that integrate into the dentate gyrus and are critical for certain types of hippocampus-dependent learning and memory [1,3]. Neurogenesis declines with age [4-6] and is impaired by various types of stress [7] and brain inflammation [8]. Exercise and environmental enrichment increase neurogenesis, and can reverse the effects of aging [9,10]

and stress [11]. Excessive NSC proliferation, however, can promote functional exhaustion of these cells [12-14] and in some cases can lead to glioma, a form of brain cancer [15-17]. Thus, regulation of NSC proliferation and differentiation is pivotal for adult brain homeostasis and is disrupted during aging.

Intrinsic and extrinsic factors regulate NSC function largely by directing changes in gene expression. A number of transcription factors and chromatin modifiers control gene expression in adult NSCs, thereby affecting NSC number and ability to differentiate into multiple cell types. These regulators include the polycomb member Bmi1 [18-20], the transcriptional repressor Tlx [21,22], and the FoxO family of transcription factors [13,14]. MicroRNAs (miRNAs) represent an additional layer of gene expression control and have recently emerged as key regulators of embryonic and adult stem cells [23,24]. miRNAs are single-stranded ~23-nucleotide RNA molecules that are

usually derived from long primary host transcripts [25]. In the cytoplasm, miRNAs direct destabilization and translational repression of target mRNAs by binding sites usually in mRNA 3' untranslated regions. This miRNA-directed downregulation of gene expression generally requires a complementary match between the mRNA target site and the second to seventh nucleotides of the miRNA 5' end (the "seed sequence"). It also depends on other regions of complementary pairing between the mRNA site and the miRNA, the presence of other miRNA-targeted sites, and the mRNA structure at this region [26]. Several hundred miRNAs have been identified in humans and mice [27]. As each miRNA potentially targets hundreds of different mRNAs [28], miRNAs can coordinate cell behaviors by fine-tuning gene expression [26,29].

A number of miRNAs recently have been found to regulate adult NSCs in vivo and in culture [30]. For example, two miRNAs, let-7b and miR-9, inhibit NSC proliferation and promote neuronal differentiation by suppressing *Tlx* and the oncogenic chromatin regulator *Hmga2* [31-33]. In addition, miR-124 promotes differentiation of SVZ NSCs into neuroblasts by repressing the expression of the transcription factor *Sox9* [34]. Finally, miR-184 and miR-137 trigger NSC proliferation and inhibit differentiation by repressing the NSC fate-regulator *Numlike* [35] and the polycomb methyltransferase *Ezh2* [36], respectively. Thus, miR-124, miR-9, and let-7b elicit NSC differentiation, while miR-184 and miR-137 increase proliferation at the expense of differentiation potential. miRNAs that promote the expansion of NSCs while maintaining their ability to differentiate have not yet been identified.

The miRNAs in the miR-17 family are attractive candidates for this function. Specific miR-17 family members are overexpressed in a variety of cancers, including glioma and glioblastoma brain cancers [37-41], and promote cancer cell proliferation and survival [42-45]. Furthermore, in embryonic stem cells, miR-17 family members are repressed by the REST neuronal gene silencer [46], which negatively regulates neurogenesis [47]. miR-17 member expression in the brain declines between late embryonic and postnatal life [48], which correlates with the decline in neurogenesis that occurs during this period [49,50]. These results suggest that miR-17 members may be involved in promoting both proliferation and neurogenesis.

The miR-17 family consists of three paralogous polycistronic clusters on different chromosomes: miR-17~92 (miR-17, miR-18a, miR-19a, miR-20a, miR-19b-1, and miR-92a-1), miR-106b~25 (miR-106b, miR-93, and miR-25), and miR-106a~363 (miR-106a, miR-18b,

miR-20b, miR-19b-2, miR-92a-2, and miR-363). Members of each cluster belong to one of four groups with similar seed sequences and therefore similar mRNA targets [51]. Within the miR-17 family, members of the miR-106b~25 cluster (miR-106b, miR-93, and miR-25) appear to be the most strongly expressed in the adult brain [27,52]. Further suggesting a link between miR-106b~25 and neurogenesis, expression of the host gene for miR-106b~25, *Mcm7*, is reduced in a mouse model of Down syndrome with diminished numbers of neural progenitor cells and neurogenesis [53].

Interestingly, the miR-106b~25 genomic locus contains a consensus binding sequence for the FoxO transcription factors. FoxO factors are inhibited by the insulin/insulin-like growth factor-1 (IGF) signaling pathway [54-56] and have emerged as regulators of adult NSCs both in vitro and in vivo [13,14]. The FoxO family promotes longevity in a range of species [57-59] and is involved in nematode lifespan regulation by the miRNA *lin-4* [60]. FoxO3, one member of the FoxO family, has recently been associated with extreme longevity in humans [61-65]. The presence of a FoxO binding sequence in the miR-106b~25 genomic locus raises the possibility of an interaction between this miRNA cluster and the insulin/IGF-FoxO pathway in mammals.

Here we use primary cultures of neural stem/progenitor cells (NSPCs) from adult mice to show that miR-106b~25 promotes NSPC proliferation. Knocking down miR-25 decreases NSPC proliferation, and ectopically expressing miR-25 or the entire miR-106b~25 cluster increases proliferation. In NSPCs induced to differentiate, overexpressing miR-106b~25 enhances differentiation toward the neuronal lineage. We find that potential miR-25 target mRNAs are overrepresented in insulin/IGF signaling. Furthermore, we show that FoxO3 occupies a binding site near the promoter for miR-106b~25 in NSPCs, raising the possibility of a FoxO-miR-106b~25 feedback loop. Together, these results suggest that miR-106b~25 modulates adult NSPC proliferation and neuronal differentiation, which may have crucial implications for the maintenance of adult neurogenesis.

RESULTS

miR-106b, miR-93, and miR-25 are expressed in adult NSPC cultures

We examined the expression levels of the miR-106b~25 cluster members (miR-106b, miR-93, and miR-25; Figure 1A) in self-renewing or differentiating NSPCs isolated from young adult (3 month-old) mice. After the

first passage in culture, NSPCs were placed in self-renewal conditions or in differentiation conditions known to give rise to astrocytes, neurons, and oligodendrocytes [14,66]. We confirmed differentiation of NSPCs into these cell types by staining for markers of astrocytes (GFAP-positive), neurons (Tuj1-positive), and oligodendrocytes (O4-positive) [67] after seven days of differentiation (Figure 1B). We then tested the expression of miR-106b~25 by RT-qPCR in self-renewing and differentiating NSPCs (Figure 1C). We

found that miR-106b, miR-93, and miR-25 were all expressed in self-renewing NSPCs. Expression of these miRNAs was not significantly changed by multi-lineage differentiation, although these miRNAs tended to be slightly upregulated during differentiation. In contrast, miR-9, a miRNA known to be induced by NSPC differentiation [33], was significantly upregulated in differentiating NSPCs. Together, these results indicate that miR-106b~25 is expressed in both self-renewing and differentiating adult NSPCs.




Figure 1. The miR-106b~25 cluster is expressed in adult NSPCs in culture. (A) Genomic locus of the mouse miR-106b~25 cluster and its host gene, *Mcm7*. (B) NSPCs (age 12 weeks, passage 2) were grown in multi-lineage differentiation conditions (no EGF or bFGF, with 1% FBS) for 7 days and then stained for Tuj1 (a marker of neurons), GFAP (a marker of astrocytes), or O4 (a marker of oligodendrocytes). Scale bar: 100 μ m. (C) miRNA expression was determined by RT-qPCR in NSPCs in self-renewal conditions (with EGF and bFGF, no FBS) or differentiation conditions (no EGF or bFGF, with 1% FBS) for 4 days. Mean and SEM of gene expression relative to self-renewal conditions for 3 independent NSPC cultures (age 12 weeks, passage 2) are shown. One-sample two-tailed t-test, *: $p < 0.05$.

miR-25 is important for adult NSPC proliferation

We next tested whether miR-106b~25 is important for adult NSPC proliferation in self-renewal conditions. To inhibit miR-106b~25, we transfected NSPCs with locked nucleic acid (LNA)-modified oligonucleotides antisense to miR-106b, miR-93, or miR-25, or with a scrambled control LNA oligonucleotide. We assessed incorporation of the thymidine analog 5-ethynyl-deoxyuridine (EdU) in NSPCs transfected with LNA probes antisense to each of the miRNAs in the miR-106b~25 cluster or with control LNA probes. We found that miR-25 knockdown decreased EdU incorporation in NSPCs by 45% ($p=0.005$), whereas miR-106b or miR-93 knockdown did not significantly affect EdU incorporation in NSPCs (Figure 2). These results indicate that within the miR-106b~25 cluster, miR-25 is the most important for NSPC proliferation.




Figure 2. miR-25 is necessary for adult NSPC proliferation. NSPCs were transfected to knock down miR-106b, miR-93, or miR-25 or were transfected with a scrambled control oligonucleotide. Two days after transfection, NSPCs were incubated with EdU for 1 hour and then immediately fixed for analysis. (A) Representative photos for control knockdown and miR-25 knockdown. Scale bar: 100 μ m. (B) Mean and SEM of the proportion of EdU+ cells for each condition, for experiments on 5 independent NSPC cultures (age 8-14 weeks, passage 3-7). Paired two-tailed t-test, **: $p<0.01$.




Figure 3. Expression of miR-25 enhances adult NSPC proliferation. NSPCs were infected with an empty control retrovirus (expressing a GFP marker only) or a retrovirus expressing miR-25. NSPCs were grown to full neurospheres for about 1 week after infection before miRNA expression and proliferation were analyzed. (A) miR-25 expression was assessed with RT-qPCR in control versus miR-25-overexpressing NSPCs. Mean and SEM of 2 independent NSPC cultures (age 12 weeks, passage 2-5) are shown. (B) Representative photos for each condition. Scale bar: 100 μ m. (C) Control and miR-25-overexpressing NSPCs were dissociated and incubated with EdU for 1 hour. Mean and SEM of the proportion of EdU+ cells for each condition, for experiments on 4 independent NSPC cultures (age 12 weeks, passage 3-6), are shown. Paired two-tailed t-test, *: $p<0.05$.

Ectopic expression of miR-25 promotes proliferation in adult NSPCs

To test if miR-25 could promote proliferation in adult NSPCs, we ectopically expressed miR-25 in NSPCs using a retroviral vector containing the miR-25

precursor and green fluorescent protein (GFP). We verified by RT-qPCR that miR-25 was overexpressed, on average by 8-fold, in NSPCs after miR-25 retrovirus infection (Figure 3A). We found that ectopic miR-25 expression increased NSPC incorporation of EdU by 18% compared to the GFP-only control ($p=0.04$; Figure 3B,C).

Expression of the entire miR-106b~25 cluster also promotes adult NSPC proliferation

We next tested whether overexpressing the entire miR-106b~25 cluster in adult NSPCs could further enhance the proliferation increase caused by miR-25 overexpression. We generated a retroviral construct containing the 725-bp portion of the mouse gene encoding the miR-106b, miR-93, and miR-25 precursors. We verified by RT-qPCR that each member of miR-106b~25 was overexpressed in cells infected with miR-106b~25 retroviruses: miR-106b~25 expression was increased 10- to 30-fold in NSPCs infected with miR-106b~25 retroviruses compared to control retroviruses (Figure 4A). We assessed the proportion of cells that incorporated EdU or bromodeoxyuridine (BrdU), another thymidine analog, in miR-106b~25 expressing versus control NSPCs. Ectopic expression of miR-106b~25 increased thymidine analog incorporation by an average of 21% ($p=0.03$; Figure 4B,C), similar to miR-25 alone, supporting the idea that miR-25 is the main miR-106b~25 member influencing NSPC proliferation. Together, these results indicate that miR-106b~25 promotes adult NSPC proliferation, and this is likely due mainly to miR-25.

Expression of the miR-106b~25 cluster promotes neuronal differentiation of adult NSPCs

We examined how miR-106b~25 influences the generation of neurons from NSPCs during multi-lineage differentiation in culture. Because the short-term nature of LNA-mediated miRNA knockdown is not compatible with the duration of NSPC differentiation, we examined the effect of retrovirus overexpression of miR-106b~25 on neuronal differentiation. We infected NSPCs with retroviruses expressing miR-106b~25 or control retroviruses and then differentiated these cells for seven days. We stained cells for Tuj1, a marker of neurons, and determined the proportion of Tuj1-positive cells (Figure 5). Although infected NSPCs formed relatively few neurons – probably a consequence of the toxicity of the infection – we found that compared to control infection, miR-106b~25 expression consistently increased the proportion of Tuj1-positive cells, on average from 0.3% to 0.9% (2.6-fold; $p=0.005$). These results indicate that ectopic expression of miR-106b~25




Figure 4. Expression of the entire miR-106b~25 cluster also enhances adult NSPC proliferation. NSPCs were infected with an empty control retrovirus (expressing a GFP marker only) or a retrovirus expressing miR-106b, miR-93, and miR-25 simultaneously (miR-106b~25). NSPCs were grown to full neurospheres for about 1 week after infection before miRNA expression and proliferation were analyzed. (A) miR-106b, miR-93, and miR-25 expression was assessed with RT-qPCR in control versus miR-106b~25-overexpressing NSPCs. Mean and SEM of 4 independent NSPC cultures (age 12-14 weeks, passage 5-14) are shown. (B) Representative photos for each condition. Scale bar: 100 μ m. (C) Control and miR-106b~25-overexpressing NSPCs were dissociated and incubated with EdU or BrdU for 1 hour. Mean and SEM of the proportion of EdU+ or BrdU+ cells for each condition, for 6 experiments on independent NSPC cultures (age 12-14 weeks, passage 3-14), are shown. Paired two-tailed t-test, *: $p<0.05$.




Figure 5. miR-106b~25 enhances neurogenesis in culture. NSPCs were infected with an empty control virus or virus to overexpress miR-106b~25. Three days after infection, NSPCs were placed in differentiation conditions for 7 days, and then stained for Tuj1, a marker of neurons. **(A)** Representative photos for each condition. Scale bar: 50 μ m. **(B)** Mean and SEM of the proportion of Tuj1+ cells (total Tuj1+ cells/total DAPI-stained nuclei) normalized to control infection, for experiments on 4 independent NSPC cultures (age 12 weeks, passage 2), are shown. Paired two-tailed t-test, **: $p < 0.01$.

can enhance neurogenesis in culture.

miR-25 has a number of predicted targets in the TGF β and insulin/IGF-FoxO pathways

We next sought to identify the molecular networks involving miR-25, the main miR-106b~25 member controlling NSPC proliferation. Computational algorithms have been developed to predict miRNA binding sites on target mRNA transcripts, based on miRNA-target site complementarity, site context, and site conservation [26]. To examine miR-25 targets through multiple bioinformatics approaches, we first

used the TargetScan program [68] to predict the conserved mRNA targets of miR-25 (~600 targets) and then used the gene classification programs PANTHER [69,70] (Figure 6A) or GSEA [71] (Figure 6B) to associate biological processes and gene sets with these targets. In a parallel approach, we used the DIANA-miRPath program [72] to predict miR-25 targets (~150) with the DIANA-microT-3.0-Strict algorithm [73] followed by comparison with the Kyoto Encyclopedia of Genes and Genomes (KEGG) biological pathways [74] (Figure 6C). A number of interesting molecular networks were enriched for miR-25 targets, including p53 signaling, hypoxia signaling, and nitric oxide signaling, which are all important for NSC maintenance and activity [75-77]. Two signaling pathways in particular stood out from this target analysis: transforming growth factor β (TGF β)/bone morphogenic protein (BMP) signaling, which was enriched for miR-25 targets in all three bioinformatics approaches, and insulin/IGF signaling, which was enriched for miR-25 targets in the TargetScan-PANTHER analysis (Figure 6D). TGF β signaling has been shown to inhibit adult NSC proliferation and neurogenesis [78,79], suggesting that miR-25 might promote NSC proliferation and neuronal differentiation by repressing TGF β signaling. Activation of the insulin/IGF pathway, which inhibits FoxO factors [55], increases NSPC proliferation and self-renewal [80-83], and FoxO factors are necessary to maintain the relatively quiescent pool of adult NSCs [13,14]. The observation that the insulin/IGF-FoxO pathway is enriched for miR-25 targets is especially pertinent because the genomic locus of miR-106b~25 contains a conserved FoxO binding sequence (Figure 7A). Furthermore, there is crosstalk between TGF β signaling and the insulin/IGF-FoxO pathway in nematode longevity, mammalian stem cells, and cancer cells [84-86]. Taken together, these results suggest that modulation of the TGF β and insulin/IGF signaling pathways may mediate part of the effects of miR-25 in NSPCs.

The transcription factor FoxO3 binds to a site in the first intron of miR-106b~25/Mcm7

The precursors of miR-106b~25 members are all located in the thirteenth intron of the protein-coding gene Mcm7, a member of a DNA helicase family required for DNA replication [87]. The first intron of the Mcm7 gene contains a conserved core binding sequence (TTGTTTAC) for the FoxO proteins [88,89] (Figure 7A). As the FoxO factors, particularly FoxO3, are important for NSC self-renewal, proliferation, and differentiation [13,14], we tested whether FoxO3 could bind to this site in the first intron of miR-106b~25/Mcm7. We performed an electrophoretic




Figure 6. miR-25 targets genes involved in TGFβ and insulin/IGF signaling. (A) The PANTHER gene classification program was used to analyze TargetScan-predicted conserved targets for mouse miR-25 (~600 targets total). Shown are the top 5 biological pathways (ordered by Bonferroni-corrected binomial test p-values). (B) The GSEA program was used to analyze the same TargetScan-predicted target list as in (A), using the Canonical Pathways and GO Gene Sets categories. Shown are the top 5 categories (ordered by hypergeometric distribution-generated p-values). (C) The DIANA-microT program was used to generate a stringent list of mouse miR-25 targets. Shown are the top KEGG categories (ordered by Pearson's chi-square test p-values). (D) Pathway diagrams based on those in PANTHER Pathways for TGFβ and insulin/IGF-Akt signaling pathways, modified for simplicity and with select miR-25 predicted targets listed.

mobility shift assay (EMSA) in which recombinant FoxO3 was incubated with a 38-bp probe containing the FoxO binding sequence in the miR-106b~25 genomic locus. We found that FoxO3 caused a band shift of this probe, showing that FoxO3 directly binds this site in vitro (Figure 7B). To determine if FoxO3 is present at the binding site at the miR-106b~25 locus in NSPCs in the context of endogenous chromatin, we performed FoxO3 chromatin immunoprecipitation (ChIP) on NSPCs treated with brief growth factor removal and the PI3K inhibitor LY294002, to activate endogenous FoxO3 (Figure 7C). ChIP-qPCR showed that endogen-

ous FoxO3 occupies the binding site in the first intron of miR-106b~25/Mcm7 in cultured adult NSPCs. This enrichment was not present in FoxO3-null NSPCs, verifying the specificity of the FoxO3 ChIP. These results indicate that FoxO3 is bound at the genomic locus of the miR-106b~25 cluster.

To test if FoxO3 could upregulate the transcription of miR-106b~25/Mcm7, we generated a luciferase reporter construct containing a minimal SV40 promoter and the 500 bp surrounding the FoxO binding site in the first intron of miR-106b~25/Mcm7 (Figure 7A). We co-

transfected HEK 293T cells with this reporter construct and with plasmids to express wild-type FoxO3, a DNA-binding-defective inactive form of FoxO3, or constitutively active FoxO3. These luciferase assays revealed that constitutively active FoxO3 enhanced luciferase expression ($p=0.002$), and this was partly abrogated by mutating the FoxO binding site ($p=0.08$), indicating that FoxO3 acts as a transcriptional activator at this genomic locus in HEK 293T cells (Figure 7D).

We next investigated whether FoxO3 affects endogenous miR-106b~25 and Mcm7 expression in NSPCs by comparing the expression of miR-106b~25

and Mcm7 in cultured NSPCs from wild-type versus FoxO3-null adult mice (Figure 7E). FoxO3-null NSPCs had decreased abundance of Mcm7 mRNA ($p=0.01$), indicating that Mcm7 is a target gene of FoxO3. However, FoxO3-null NSPCs did not display decreased expression of the mature forms of miR-106b, miR-93, and miR-25, suggesting that FoxO3 does not directly upregulate miR-106b~25 and might even indirectly repress the expression of this cluster. Together, these results suggest a complex regulation in which FoxO3 activates the transcription of miR-106b~25/Mcm7, but may repress miR-106b~25 abundance, perhaps by a posttranscriptional mechanism or by acting at a separate promoter for miR-106b~25.




Figure 7. FoxO3 binds to a site in the first intron of miR-106b~25/Mcm7. (A) Location of the FoxO binding site (FHRE) within the first intron of the miR-106b~25/Mcm7 gene, and the sequence locations used for EMSA, ChIP, and luciferase experiments. (B) EMSA with recombinant FoxO3-GST and a radioactively-labeled (hot) probe corresponding to the FoxO binding site in miR-106b~25/Mcm7 (FHRE WT). + Ctrl: FoxO3-GST incubated with a probe for a known FoxO binding site. The specificity of the interaction was tested by increasing amounts of unlabeled (cold) probe or cold probe with mutations in the FoxO consensus binding sequence (FHRE Mut). (C) Wild-type and FoxO3-null NSPCs were dissociated and the next day treated with 4 hours growth factor removal followed by addition of LY294002 for 1 hour. Antibodies to FoxO3 or control IgG antibodies were used for ChIP. qPCR was used to assess the enrichment of FHRE and of a negative control site (- Ctrl). Shown is the relative enrichment for 1 experiment (age 12 weeks, passage 10). These results were confirmed in ChIP-Seq studies (Webb et al. submitted). (D) HEK 293T cells were co-transfected with a plasmid to express FoxO3 (empty control, wild-type FoxO3, FoxO3 lacking the DNA binding domain, or constitutively nuclear FoxO3), a firefly luciferase reporter containing FHRE with or without the FoxO consensus sequence mutated, and a Renilla luciferase reporter to normalize for transfection efficiency. As a positive control, a luciferase reporter containing a known FoxO3-activated site was used (+ Ctrl); as a negative control, a luciferase reporter without an enhancer site was used (- Ctrl). Luciferase activity was assessed two days after transfection. Mean and SEM for 4 independent experiments (- Ctrl, + Ctrl, and FHRE WT) or 2 independent experiments (FHRE Mut) are shown. Unpaired two-tailed t-test, **: $p<0.01$. (E) NSPCs from wild-type and FoxO3-null mice were isolated and cultured. Total RNA was collected, and the levels of mature miR-106b~25 members (relative to 5S RNA) and Mcm7 mRNA (relative to β -actin mRNA) were assessed by RT-qPCR. Mean and SEM of the FoxO3-null/wild-type fold change for 5-6 independent cultures (age 10-13 weeks, passage 2-5) are shown. One-sample two-tailed t-test, *: $p<0.05$.

DISCUSSION

miR-106b~25 members are known to promote cancer cell proliferation and survival [42,44], modulate embryonic stem cell differentiation [90], and promote reprogramming of mouse embryonic fibroblasts into induced pluripotent stem cells [91] – but the importance of miR-106b~25 has not been investigated in an adult stem cell population before. This study examined the role of miR-106b~25 in adult NSPCs. We found that miR-25 knockdown decreases NSPC proliferation, miR-25 or miR-106b~25 overexpression increases adult NSPC proliferation, and miR-106b~25 overexpression promotes neuronal differentiation. Furthermore, FoxO3 binds near the promoter for the host gene of miR-106b~25 and thus has the potential to influence miR-106b~25 expression. These results add to our understanding of the factors regulating NSPC activity and suggest that oncogenic miRNAs could have physiological functions in adult stem cells.

miR-106b~25 and NSPC proliferation

The effects of miR-106b~25 on adult NSPC proliferation are modest: miR-106b~25 or miR-25 overexpression increased NSPC proliferation by about 1.2-fold, miR-25 knockdown reduced proliferation by about 1.4-fold, and individual miR-106b and miR-93 knockdowns did not affect NSPC proliferation. While these proliferation changes are somewhat smaller than those seen by miR-106b~25 manipulation in carcinoma cells (up to 1.8-fold in similar assays) [42,44], a modest effect of miR-106b~25 on NSPC proliferation could still be important physiologically. While an enforced large increase in NSC proliferation rate could result in tumor initiation [15,17], a weaker increase in proliferation rate could lead to premature stem cell exhaustion [12,13]. On the other hand, fewer divisions could reduce NSC number and neurogenesis [32,92]. Therefore, it is possible that miR-106b~25 overexpression or underexpression, both of which alter NSPC proliferation in culture, could affect long-term NSC function in vivo.

Redundancy within the miR-17 family could dampen the influence of miR-106b~25 on NSPC proliferation in vitro. Knockdown of miR-106b or miR-93, which share the same mRNA-targeting seed sequence, did not affect proliferation, while knockdown of miR-25, which has a different seed sequence, reduced proliferation. miR-106b and miR-93 might be able to compensate for each other in NSPCs, which could be tested by inhibiting both miRNAs simultaneously. Furthermore, it is possible that NSPCs buffer miR-106b~25 alteration by expressing miR-17 family members from the other

paralogous clusters, thereby lessening the relative importance of one or even three miRNAs within this family, or allowing NSPCs to react to changes in miR-106b~25 expression with compensatory changes in miR-17~92 or miR-106a~363 expression. Our findings suggest the idea that compared to cancer cells, stem cells may be more resilient against oncogene perturbation, and therefore more tolerant of certain gene-specific anti-cancer therapies. This may be particularly true for miRNAs, which have been duplicated during animal evolution and tend to have overlapping targets and functions. Such redundancy may have evolved not only so that duplicated miRNAs can be controlled by distinct cis regulatory elements, but perhaps also so that stem cells can absorb fluctuations in gene expression.

miR-106b~25 in neuronal differentiation

We found that miR-106b~25 promotes both NSPC proliferation in self-renewal conditions and neuron production in differentiation conditions, whereas other miRNAs previously studied in adult NSCs seem to promote one function while inhibiting the other. The mechanism of this effect is still unknown: miR-106b~25 could affect NSPC tendency to produce neurons instead of glia, neuronal progenitor proliferation and survival, and neuron survival. Thus, it remains to be determined whether miR-106b~25 influences neurogenesis by directing cell fate or by regulating cell division and survival in specific cell types.

Adult NSCs decline in number and proliferation, neurogenesis, and self-renewal abilities during aging [5]. Activities that restore NSC activity, such as exercise or environmental enrichment, also restore cognitive performance in aged mice [93,94]. As NSC decline may contribute to cognitive aging, investigating how miR-106b~25 affects neurogenesis will improve our understanding of the molecular mechanisms involved in cognitive aging. While miR-106b~25 knockout mice have no apparent phenotype [52], neurogenesis and learning have not been examined in these mice. It would be worthwhile to investigate how NSCs lacking or overexpressing miR-106b~25 in vivo preserve their numbers and sustain neurogenesis throughout life.

Potential signaling pathways regulated by miR-25

Deciphering how stem cells sense and respond to tissue integrity and nutrient supply is key to understanding how stem cells maintain tissue homeostasis and how this function changes with age [83,95,96]. Analyzing candidate targets of miR-25 revealed that miR-25 might

modulate TGF β or insulin/IGF signaling at multiple points in each pathway. As TGF β signaling negatively regulates adult NSC proliferation and neurogenesis [78,79], one way miR-106b~25 might promote these behaviors is by repressing TGF β signaling in NSPCs. TGF β Receptor-2 is directly repressed by miR-106b in neuroblastoma cells [97] and by miR-106b and miR-93 in mouse embryonic fibroblasts [91]; thus, one enticing possibility is that TGF β Receptor-2 is targeted by all miR-106b~25 members in NSPCs. While inhibitory Smads (Smad6 and Smad7) are also predicted miR-25 targets, Smad7-deficient mice have increased adult NSPC proliferation and numbers, which may be due to TGF β -independent mechanisms [98]. The net functional effect of miR-25 regulation of TGF β signaling in NSPCs will depend on the relative expression, degree of miR-25 repression, and network connections of each member of the TGF β pathway in NSPCs.

Activation of the insulin/IGF pathway is sufficient to increase NSPC proliferation and self-renewal [80-83], while FoxO factors are necessary to prevent overproliferation, abnormal differentiation, and long-term depletion of NSCs [13,14]. Thus, another way miR-25 might increase NSPC proliferation is by de-repressing insulin/IGF signaling. Given that PTEN can be a major inhibitor of insulin/IGF signaling [99,100] and is a known target of miR-25 in prostate cancer cells [101], miR-25 may target PTEN to increase insulin/IGF signaling and repress FoxO activity. We cannot exclude the possibility, however, that miR-25 negatively regulates insulin/IGF signaling under some circumstances, such as by repressing Akt or PI3K.

There may even be crosstalk between the different pathways targeted by miR-25. In nematodes the TGF β pathway has been shown to genetically interact with the insulin/IGF-FoxO pathway to regulate lifespan [84]. In mammals TGF β promotes hematopoietic stem cell quiescence by downregulating Akt activity and upregulating FoxO3 activity [85], and in glioblastoma cells TGF β signaling induces Smad-FoxO transcriptional activation complexes that suppress proliferation [86]. In human keratinocytes, FoxO factors are required for the induction of a number of genes by TGF β , particularly cytostatic and stress response genes [102]. Thus, it is possible that miR-25 regulate NSPCs by coordinately modulating insulin/IGF and TGF β networks.

Regulation of miR-106b~25 by FoxO proteins

Our experiments suggest that FoxO3 regulates miR-106b~25 in a complex manner. FoxO3 binds to a site in the first intron of miR-106b~25/Mcm7 in NSPCs. In

FoxO3-null NSPCs, while Mcm7 mRNA abundance was decreased, the levels of mature miR-106b~25 members were not decreased, and were even slightly increased. Thus, FoxO3 might transcriptionally activate miR-106b~25/Mcm7, but act to repress miR-106b, miR-93, and miR-25 at a different promoter or at posttranscriptional steps like precursor cleavage, nuclear export, base editing, and degradation.

Other factors complicate our ability to define the regulation of miR-106b~25 by FoxO3. It is possible that in self-renewal culture conditions FoxO3 is bound near the promoter of miR-106b~25 but exerts control over miR-106b~25 expression only in other conditions such as differentiation, low nutrient levels, oxidative stress, or low oxygen tension. As NSPC cultures are heterogeneous, containing mixtures of stem cells, progenitor cells, and even some differentiated progeny [103,104], FoxO3 might also alter miR-106b~25 expression differently in different cell types. Such differential regulation would be consistent with FoxO3 and miR-106b~25 both promoting neuronal differentiation but having opposite effects on NSPC proliferation [13,14]. In these scenarios, FoxO3 would serve as one component of a “coincidence detector” regulating miR-106b~25, which in turn might indirectly influence FoxO activity.

CONCLUSION

This study shows that miR-106b~25 members modulate NSPC proliferation and differentiation and could potentially be regulated by the pro-longevity transcription factor FoxO3 under some circumstances. These results suggest a role for miR-106b~25 in normal adult stem cell function, in addition to a known role in cancer cells. Understanding how miR-106b~25 and FoxO3 function in NSPCs could reveal new strategies for preventing the loss of neurogenesis in adults, particularly during aging.

METHODS

Constructs. For miRNA overexpression, the 725-bp segment of the mouse Mcm7 gene containing the miR-106b, miR-93, and miR-25 precursors was cloned between the XhoI and PmeI sites of the MDH1-PGK-GFP 2.0 vector [105] using the primers F: 5'-AAACTCGAGCCTGCTGGCCATTCTCCGACTTTC C-3' and R: 5'-AAAGTTTAAACGGATCTTTCTTTGCTCCAGCTT CAAGC-3'. The 350-bp segment of the mouse Mcm7 gene containing the miR-25 precursor only was cloned between the XhoI and EcoRI sites of the MDH1-PGK-GFP 2.0 vector using the primers F: 5'-

AAACTCGAGCCCAGGACACAACCTCTGAT-3'
and R: 5'-
AAAGAATTCGAGGGGAATGAAGTCAAGGA-3'.

For luciferase assays, the 500-bp region of the mouse *Mcm7* intron containing the FoxO3 binding site was cloned between the KpnI and XhoI sites of the pGL3-SV40 vector (Promega) using the primers F: 5'-AAAGGTACCGCAGTGTTCCTTTTCACAAGTCCG-3' and R: 5'-AAACTCGAGCGTGTGTAAACAGTGTCTTCCGC-3'. Mutations in the FoxO binding sequence were made using the primers F: 5'-CCGCTCTTAATAGACAAAGAAGCACATGGGCC CAGATTCC-3' and R: 5'-GGAATCTGGGCCCATGTGCTTCTTTGTCTATTA AGAGCGG-3', and this mutated enhancer was subcloned into a new pGL3-SV40 backbone. The positive control plasmid, pGL3-SV40 containing three repeats of the FoxO3 binding site in the *FasL* promoter, and the FoxO3 expression plasmids were described previously [55].

Antibodies. For immunocytochemistry, the primary antibodies used were rat anti-BrdU (AbD Serotec; 1:500), goat anti-GFP (Rockland; 1:500), rabbit anti-Tuj1 (Covance; 1:1000), rat anti-GFAP (Calbiochem; 1:1000), and mouse anti-O4 (a gift from Ben Barres; 1:1000). Fluorescent secondary antibodies were from Jackson Immuno-Research and Molecular Probes (Invitrogen) and were used at 1:400 dilutions. The antibodies for ChIP were rabbit anti-FoxO3 "NFL" (Brunet laboratory) and rabbit IgG (Zymed).

NSPC isolation and culture. Each NSPC culture was generated from four to eight FVB/N mice (1:1 male-female ratio). Whole brain was extracted from each animal, and the olfactory bulbs, cerebellum, and brainstem were discarded. To dissociate the forebrain tissue, brains were diced, treated at 37°C for 30 min with HBSS (Invitrogen) containing 2.5 U/ml Papain (Worthington), 1 U/ml Dispase II (Roche), 250 U/ml DNase I (Sigma), and 1X penicillin-streptomycin-L-glutamine (PSQ; Invitrogen), and then mechanically dissociated in DMEM/F12 (Invitrogen) containing 10% fetal bovine serum (FBS; Invitrogen) and 1X PSQ. NSPCs were purified from myelin with a 22.5% Percoll gradient (GE Healthcare) and then from red blood cells with a 58.5% Percoll gradient. Freshly isolated NSPCs were considered "passage 1."

NSPCs were grown at 5% CO₂ in a 37°C incubator at 50,000 cells/ml in Neurobasal A Medium (NBA; Invitrogen) supplemented with 1X PSQ, 1X B-27 Supplement Minus Vitamin A (B27; Invitrogen), 20

ng/ml recombinant human bFGF (PeproTech), and 20 ng/ml recombinant human EGF (PeproTech). Cells were fed every 2 days by replacing half the media and replenishing bFGF and EGF; cells were transferred to a new plate every 4 days. NSPCs grew to full neurosphere colonies every 5-8 days, and were passaged using Accutase (Millipore) for dissociation.

miRNA overexpression by retroviral infection. HEK 293T cells were co-transfected with the expression vector MDH1-PGK-GFP 2.0 containing either miR-106b~25 or no insert (empty control) and the pCL-Eco viral packaging vector in a 2:1 ratio, using the calcium phosphate transfection method. The media was changed to NBA containing 1X PSQ and 1X B27 6-8 h later. The next day, NSPCs were dissociated and plated at 50,000 cells/ml on plates coated with 50 µg/ml poly-D-lysine (Sigma). The following day, NSPCs were infected by replacing half the media with 0.45 µm-filtered virus-containing supernatant from the 293T cultures and replenishing the growth factors. Sixteen hours later, the infection was stopped by replacing all the media with NSPC-conditioned media and fresh media in a 1:1 ratio and replenishing growth factors. NSPCs were fed every other day until they were 80% confluent, and then detached with Accutase and grown in suspension. After NSPCs had grown to full neurospheres, RNA and protein were collected, and cells were plated for proliferation assays.

miRNA knockdown. NSPCs were plated at 100,000 cells/ml in 0.5 ml NBA containing 1X L-glutamine (Invitrogen) and 1X B27 with growth factors in a poly-D-lysine-coated well of a 24-well plate. The next day, 45 nM locked nucleic acid (LNA) oligonucleotide (Exiqon) was diluted with 100 µl Opti-MEM (Invitrogen), incubated with 1 µl Lipofectamine PLUS reagent (Invitrogen) per 1 µg nucleic acid for 5 min, and then incubated with 6 µl Lipofectamine LTX reagent (Invitrogen) per 1 µg nucleic acid for 30 min before being added to cells. The media was changed to 1 ml NBA containing 1X PSQ and 1X B27 with growth factors 4-6 h later.

Proliferation assays. One week after retroviral infection (when NSPCs had grown to full neurospheres), NSPCs were dissociated and plated on nitric acid-treated glass coverslips (Bellco) coated with poly-D-lysine. Two days later, BrdU (EMD Biosciences) was added to a final concentration of 10 µM, or EdU (Invitrogen) was added to a 5 µM final concentration. One hour later, NSPCs were fixed in 4% paraformaldehyde in PBS for 12 min. The coverslips were blocked for 1 h with 10% donkey serum and 0.1% Triton in PBS and then incubated with goat anti-GFP antibody for 2 h. The

coverslips were then refixed with 4% paraformaldehyde for 10 min and incubated with 0.4% Triton for 30 min. DNA was denatured with 2 N HCl for 10 min. After 1 h of blocking, coverslips were incubated with rat anti-BrdU antibody for 2 h. The coverslips were incubated with Texas Red donkey anti-rat and FITC donkey anti-goat secondary antibodies for 1 h. The coverslips were mounted on slides using Vectashield with DAPI (Vector Labs).

Two days after transfection with LNA probes, EdU was added to a final concentration of 5 μ M. One hour later, NSPCs were fixed in 4% paraformaldehyde and 2% sucrose for 12 min. Cells were permeabilized with 0.4% Triton in PBS for 30 min and blocked with two 3% BSA (USB) rinses. Cells were then incubated in 1X Click-iT Reaction Buffer, 4 mM CuSO₄, 1:400 Alexa Fluor 594 azide, and 200 nM Click-iT EdU Buffer Additive (Invitrogen) for 30 min. Cells were then washed with 3% BSA, rinsed with PBS, and mounted on slides using Vectashield with DAPI.

Coverslips were examined using a Zeiss Axioskop 2 Plus microscope and digital camera with AxioVision 4 software. For quantification, 3-6 random fields (about 1000-2000 cells) were counted in a blinded manner, using Metamorph 7.0 software.

Differentiation assays. NSPCs were dissociated and plated on nitric acid-treated coverslips coated with poly-D-lysine at a density of 25,000 cells/ml. NSPCs were infected the next day, and the infection was stopped after 16 h. Two days later, NSPCs were differentiated by changing the media to NBA containing 1X PSQ, 1X B27, and 1% FBS. The media was replaced every other day. After 7 days of these differentiation conditions, NSPCs were stained for GFAP, Tuj1, or O4. For GFAP and Tuj1 staining, NSPCs were fixed in 4% paraformaldehyde and 2% sucrose. The coverslips were blocked for 1 h with 10% donkey serum and 0.1% Triton in PBS, and then incubated with rabbit anti-Tuj1 antibody for 2 h. After rinsing with PBS containing 0.01% Tween and blocking for another 15 min, coverslips were incubated with Texas Red donkey anti-rabbit or anti-rat secondary antibody for 1 h. For O4 staining, NSPCs were blocked with 5% goat serum and 7.5% BSA in PBS for 1 h and then incubated with mouse anti-O4 antibody (in 10% goat serum, 1% BSA, and 100 mM L-lysine in PBS) for 2 h. After rinsing with PBS, cells were fixed in 4% paraformaldehyde and 2% sucrose, blocked for another 15 min, and then incubated with Alexa Fluor 546 goat anti-mouse secondary antibody for 1 h. Coverslips were mounted on slides using Vectashield with DAPI. The total number of neurons on each coverslip was counted in a

blinded manner, and the total number of nuclei was estimated by counting 5 random fields (about 300-600 cells) in a blinded manner.

Target prediction. TargetScan (www.targetscan.org, version 5.1) was used to predict all conserved targets for mouse miR-25. This target list was analyzed using PANTHER (www.pantherdb.org, version 7) to compare Biological Process associations for genes in this list and the reference list, "NCBI: *M. musculus* genes," or analyzed with GSEA Molecular Signatures Database (www.broadinstitute.org/gsea/msigdb/, version 3.0) to compute overlaps for genes in this list and "CP" (Canonical Pathways) and "C5" (GO Gene Sets). The DIANA-miRPath program (<http://diana.cslab.ece.ntua.gr/pathways/>) using DIANA-microT-3.0-Strict was used to predict and analyze conserved targets for mouse miR-25 in the KEGG database.

RT-qPCR. Total RNA was extracted from NSPCs using the miRVana kit (Ambion). RNA was treated to remove genomic DNA in a reaction containing 100 ng/ μ l RNA, 1 U/ μ l RNase OUT (Invitrogen), and 10 U/ μ l DNase I (Invitrogen) at 37°C for 15 min and 75°C for 15 min.

miRNA expression was quantified using the miRCURY LNA miRNA PCR system or the miRCURY LNA Universal RT miRNA PCR system, according to the manufacturer's instructions (Exiqon). Samples were run in triplicate on a C1000 Thermal Cycler with the CFX96 Real-Time software (Bio-Rad), and miRNA expression was normalized to 5S RNA expression.

To quantify Mcm7 mRNA expression, RT was carried out using the High Capacity cDNA Reverse Transcription kit (Applied Biosystems). Each reaction contained 1X RT Buffer, 4 mM each dNTP, 1X Random Hexamers, 1 U/ μ l RNase OUT, 2.5 U/ μ l MultiScribe Reverse Transcriptase, and 45-90 ng/ μ l RNA. RT was performed at 25°C for 10 min, 37°C for 2 h, and 85°C for 5 min. Each 20- μ l qPCR reaction contained 0.25 μ M forward (F) Primer, 0.25 μ M reverse (R) Primer, 10 μ l iQ SYBR Green Supermix (Bio-Rad), and 0.625 μ l RT reaction. The program used was 95°C for 10 min; 40 cycles of 95°C for 20 sec, 55°C for 20 sec, and 72°C for 45 sec. Samples were run in triplicate, and Mcm7 expression was normalized to β -actin expression. The Mcm7 primers were F: 5'-TGAACACCGGCTGATGATGG-3' and R: 5'-GGCCTCGGAAATACAACCTCAA-3'. The β -actin primers were F: 5'-TGTTACCAACTGGGACGACA-3' and R: 5'-CTCTCAGCTGTGGTGGTGAA-3'.

Chromatin immunoprecipitation. ChIP was performed

as described [14] using IgG or FoxO3 antibodies. Immunoprecipitated chromatin was analyzed with qPCR: each 20- μ l reaction contained 2.5 μ l DNA, 10 μ l iQ SYBR Green Supermix, 0.25 μ M F primer, and 0.25 μ M R primer. Triplicate reactions were run with the following program: 94°C for 3 min; 40 cycles of 95°C for 20 sec, 57°C for 30 sec, and 72°C for 30 sec. The primers to amplify the region surrounding the FHRE FoxO3 binding site in the Mcm7 first intron were F: 5'-TAGGCCTCCTCTGCACTCAT-3' and R: 5'-AGGAATCCTGGGCTGTGAG-3'. The negative control primers to amplify an intergenic region lacking a Forkhead binding sequence were F: 5'-GGGGGATAATGATTGCAAAA-3' and R: 5'-GCGTGGACAGAGATCTAGGC-3'. For each chromatin sample, a standard curve using five 5-fold dilutions of input chromatin was used to quantify binding at each target site in the ChIPs: linear regression ($y=-ax+b$) was performed on Ct versus $\log_5(\text{input})$, and the amount of a site in the FoxO3 ChIP relative to the IgG ChIP was calculated as $5^{-\Delta C_t/a}$, with $\Delta C_t = C_{t\text{FoxO3}} - C_{t\text{IgG}}$.

Electrophoretic mobility shift assay. Complementary oligonucleotides (20 μ M) were annealed in 100 mM NaCl by heating at 80°C for 5 min and then cooling slowly to room temperature. Annealed probe (1 μ M) was labeled with 20 μ Ci/ μ l 32 P- γ ATP and 1 U/ μ l T4 PNK at 37°C for 1 h. Annealed probes were purified on 15% polyacrylamide and resuspended in 1X TE pH 8.

Each binding reaction was performed in Binding Buffer (200 mM Tris-HCl pH 7.5, 200 mM KCl, 200 mM MgCl₂, 2% NP-40, 10% glycerol, 5 mM DTT, and 500 ng/ μ l salmon sperm DNA) and contained 50 ng/ μ l GST or human FoxO3-GST, 1000 cpm/ μ l hot probe (5 nM FHRE probe; 3 nM positive control probe), and 0, 5, 50, or 500X competing cold probe. The reactions were incubated at room temperature for 20 min and then resolved on 4% non-denaturing PAGE at 4°C. The gels were dried and then autoradiographed for 4 days. The positive control oligonucleotides for a site bound by FoxO3 near its own promoter [106] were F: 5'-AAATAACACACACGTGTGCTGGTAAACAAGCGCGCCAGCC-3' and R: 5'-GGCTGGCGCGCTTGTTTACCAGCACACGTGTGTGTTATTT-3'. The oligonucleotides for the FHRE site within the Mcm7 intron region bound by FoxO3 in ChIP experiments were F: 5'-GGCCCATGTGCTTCTTTGTTTACTAAGAGCGGAAGCAG-3' and R: 5'-CTGCTTCCGCTCTTAGTAAACAAAGAAGCACATGGGCC-3'. The oligonucleotides for this FHRE site containing mutations in the FoxO consensus binding sequence were F: 5'-

GGCCCATGTGCTTCTGTGTCTATTAAGAGCGGAAGCAG-3' and R: 5'-CTGCTTCCGCTCTTAATAGACACAGAAGCACATGGGCC-3'.

Luciferase assays. HEK 293T cells were plated in 24-well plates at 150,000 cells/ml. The next day, they were transfected using the calcium phosphate method with 400 ng each of FoxO3 expression plasmid, pGL3-SV40 firefly luciferase plasmid, and pRL-null Renilla luciferase plasmid. Two days after transfection, cells were lysed with 0.5 ml Passive Lysis Buffer (Promega) and luciferase activity was measured with the Dual Luciferase Reporter Assay system. Triplicate transfections were averaged within each experiment, and firefly luciferase activity was normalized to Renilla luciferase activity.

Statistical analysis. Gene expression (RT-qPCR experiments) was analyzed using one-sample two-tailed t-tests. NSPC phenotype (proliferation and differentiation assays) was analyzed using paired two-tailed t-tests. The luciferase assay experiments were analyzed with unpaired two-tailed t-tests.

AUTHOR CONTRIBUTIONS

J.O.B. designed, performed, and analyzed all experiments. V.M.R. trained J.O.B., helped with experimental design and analysis, and generated FoxO3-null NSPC cultures. V.A.R. helped with experimental design and analysis and with neuronal differentiation experiments. A.E.W. performed the FoxO3 ChIP and helped with the bioinformatics analysis. A.B. helped with the design and analysis of the experiments. J.O.B. and A.B. wrote the paper.

CONFLICTS OF INTEREST STATEMENT

The authors have no conflicts of interest to declare.

ACKNOWLEDGEMENTS

We thank Chang-Zheng Chen for providing the MDH1-PGK-GFP plasmid and for his advice on miRNA expression and manipulation studies. We thank Dervish Salih for generating the FoxO3-GST and GST proteins for EMSA experiments. We thank Dena S. Leeman for critically reading the manuscript and for stimulating discussion, and Elizabeth Pollina for stimulating discussion. We thank Camille Guillerey, T. Richard Parenteau, and Christopher Itoh for conducting preliminary experiments. This work was funded by an NIH/NIA grant R01 AG026648, a CIRM grant, a Brain Tumor Society grant, a Klingenstein Fellowship, and an

AFAR grant (A.B.), a Stanford University Major Grant (J.O.B.), a Stanford University Dean's fellowship (V.M.R.), an NSF graduate fellowship (V.A.R.), and an NIH/NRSA 5T32 CA09302 (A.E.W.).

REFERENCES

1. Aboous DN, Koehl M, Le Moal M. Adult neurogenesis: From precursors to network and physiology. *Physiol Rev.* 2005; 85: 523-569.
2. Lledo PM, Merkle FT, Alvarez-Buylla A. Origin and function of olfactory bulb interneuron diversity. *Trends Neurosci.* 2008; 31: 392-400.
3. Leuner B, Gould E, Shors TJ. Is there a link between adult neurogenesis and learning? *Hippocampus.* 2006; 16: 216-224.
4. Knoth R, Singec I, Ditter M, Pantazis G, Capetian P, Meyer RP, Horvat V, Volk B, Kempermann G. Murine Features of neurogenesis in the human hippocampus across the lifespan from 0 to 100 years. *PLoS One.* 2010; 5: Article No.: e8809.
5. Molofsky AV, Slutsky SG, Joseph NM, He S, Pardal R, Krishnamurthy J, Sharpless NE, Morrison SJ. Increasing p16(INK4a) expression decreases forebrain progenitors and neurogenesis during ageing. *Nature.* 2006; 443: 448-452.
6. Pekcec A, Baumgartner W, Bankstahl JP, Stein VM, Potschka H. Effect of aging on neurogenesis in the canine brain. *Aging Cell.* 2008; 7: 368-374.
7. Mirescu C, Gould E. Stress and adult neurogenesis. *Hippocampus.* 2006; 16: 233-238.
8. Monje ML, Toda H, Palmer TD. Inflammatory blockade restores adult hippocampal neurogenesis. *Science.* 2003; 302: 1760-1765.
9. Kronenberg G, Bick-Sander A, Bunk E, Wolf C, Ehninger D, Kempermann G. Physical exercise prevents age-related decline in precursor cell activity in the mouse dentate gyrus. *Neurobiol Aging.* 2006; 27: 1505-1513.
10. Wu CW, Chang YT, Yu L, Chen HI, Jen CYJ, Wu SY, Lo CP, Kuo YM. Exercise enhances the proliferation of neural stem cells and neurite growth and survival of neuronal progenitor cells in dentate gyrus of middle-aged mice. *J Appl Physiol.* 2008; 105: 1585-1594.
11. Veena J, Srikumar BN, Raju TR, Rao BSS. Exposure to enriched environment restores the survival and differentiation of new born cells in the hippocampus and ameliorates depressive symptoms in chronically stressed rats. *Neurosci Lett.* 2009; 455: 178-182.
12. Kippin TE, Martens DJ, van der Kooy D. P21 loss compromises the relative quiescence of forebrain stem cell proliferation leading to exhaustion of their proliferation capacity. *Genes Dev.* 2005; 19: 756-767.
13. Paik JH, Ding ZH, Narurkar R, Ramkissoon S, Muller F, Kamoun WS, Chae SS, Zheng HW, Ying HQ, Mahoney J, Hiller D, Jiang S, Protopopov A, et al. FoxOs cooperatively regulate diverse pathways governing neural stem cell homeostasis. *Cell Stem Cell.* 2009; 5: 540-553.
14. Renault VM, Rafalski VA, Morgan AA, Salih DAM, Brett JO, Webb AE, Villeda SA, Thekkat PU, Guillerey C, Denko NC, Palmer TD, Buftte AJ, Brunet A. FoxO3 regulates neural stem cell homeostasis. *Cell Stem Cell.* 2009; 5: 527-539.
15. Alcantara Llaguno S, Chen J, Kwon C-H, Jackson EL, Li Y, Burns DK, Alvarez-Buylla A, Parada LF. Malignant astrocytomas originate from neural stem/progenitor cells in a somatic tumor suppressor mouse model. *Cancer Cell.* 2009; 15: 45-56.
16. Zhu Y, Guignard F, Zhao DW, Liu L, Burns DK, Mason RP, Messing A, Parada LF. Early inactivation of p53 tumor suppressor gene cooperating with NF1 loss induces malignant astrocytoma. *Cancer Cell.* 2005; 8: 119-130.
17. Liu HK, Wang Y, Belz T, Bock D, Takacs A, Radlwimmer B, Barbus S, Reifenberger G, Lichter P, Schutz G. The nuclear receptor tailless induces long-term neural stem cell expansion and brain tumor initiation. *Genes Dev.* 2010; 24: 683-695.
18. Molofsky AV, He SH, Bydon M, Morrison SJ, Pardal R. Bmi-1 promotes neural stem cell self-renewal and neural development but not mouse growth and survival by repressing the p16(Ink4a) and P19(Arf) senescence pathways. *Genes Dev.* 2005; 19: 1432-1437.
19. Molofsky AV, Pardal R, Iwashita T, Park IK, Clarke MF, Morrison SJ. Bmi-1 dependence distinguishes neural stem cell self-renewal from progenitor proliferation. *Nature.* 2003; 425: 962-967.
20. Fasano CA, Dimos JT, Ivanova NB, Lowry N, Lemischka IR, Temple S. shRNA knockdown of Bmi-1 reveals a critical role for p21-rb pathway in NSC self-renewal during development. *Cell Stem Cell.* 2007; 1: 87-99.
21. Zhang CL, Zou Y, He W, Gage FH, Evans RM. A role for adult TLX-positive neural stem cells in learning and behaviour. *Nature.* 2008; 451: 1004-1007.
22. Sun GQ, Yu RT, Evans RM, Shi YH. Orphan nuclear receptor TLX recruits histone deacetylases to repress transcription and regulate neural stem cell proliferation. *Proc Natl Acad Sci U S A.* 2007; 104: 15282-15287.
23. Gangaraju VK, Lin H. MicroRNAs: key regulators of stem cells. *Nat Rev Mol Cell Biol.* 2009; 10: 116-125.
24. Arnold C, Tan R, Zhou B, Yue S, Schaffert S, Biggs J, Doyonnas R, Lo M, Perry J, Renault V, Sacco A, Somervaille T, Viatour P, et al. MicroRNA programs in normal and aberrant stem and progenitor cells. *Genome Res.* 2011: In Press.
25. Kim VN, Han J, Siomi MC. Biogenesis of small RNAs in animals. *Nat Rev Mol Cell Biol.* 2009; 10: 126-139.
26. Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell.* 2009; 136: 215-233.
27. Landgraf P, Rusu M, Sheridan R, Sewer A, Iovino N, Aravin A, Pfeffer S, Rice A, Kamphorst AO, Landthaler M, Lin C, Socci ND, Hermida L, et al. A mammalian microRNA expression atlas based on small RNA library sequencing. *Cell.* 2007; 129: 1401-1414.
28. Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell.* 2005; 120: 15-20.

29. Stefani G, Slack FJ. Small non-coding RNAs in animal development. *Nat Rev Mol Cell Biol.* 2008; 9: 219-230.
30. Li XK, Jin P. Roles of small regulatory RNAs in determining neuronal identity. *Nat Rev Neurosci.* 2010; 11: 329-338.
31. Zhao C, Sun GQ, Li SX, Lang MF, Yang S, Li WD, Shi YH. MicroRNA let-7b regulates neural stem cell proliferation and differentiation by targeting nuclear receptor TLX signaling. *Proc Natl Acad Sci U S A.* 2010; 107: 1876-1881.
32. Nishino J, Kim I, Chada K, Morrison SJ. Hmga2 promotes neural stem cell self-renewal in young but not old mice by reducing p16(Ink4a) and p19(Arf) expression. *Cell.* 2008; 135: 227-239.
33. Zhao C, Sun GQ, Li SX, Shi YH. A feedback regulatory loop involving microRNA-9 and nuclear receptor TLX in neural stem cell fate determination. *Nat Struct Mol Biol.* 2009; 16: 365-371.
34. Cheng LC, Pastrana E, Tavazoie M, Doetsch F. miR-124 regulates adult neurogenesis in the subventricular zone stem cell niche. *Nat Neurosci.* 2009; 12: 399-408.
35. Liu CM, Teng ZQ, Santistevan NJ, Szulwach KE, Guo WX, Jin P, Zhao XY. Epigenetic regulation of miR-184 by MBD1 governs neural stem cell proliferation and differentiation. *Cell Stem Cell.* 2010; 6: 433-444.
36. Szulwach KE, Li X, Smrt RD, Li Y, Luo Y, Lin L, Santistevan NJ, Li W, Zhao X, Jin P. Cross talk between microRNA and epigenetic regulation in adult neurogenesis. *J Cell Biol.* 2010; 189: 127-141.
37. Ciafre SA, Galardi S, Mangiola A, Ferracin M, Liu CG, Sabatino G, Negrini M, Maira G, Croce CM, Farace MG. Extensive modulation of a set of microRNAs in primary glioblastoma. *Biochem Biophys Res Commun.* 2005; 334: 1351-1358.
38. Ernst A, Campos B, Meier J, Devens F, Liesenberg F, Wolter M, Reifenberger G, Herold-Mende C, Lichter P, Radlwimmer B. De-repression of CTGF via the miR-17-92 cluster upon differentiation of human glioblastoma spheroid cultures. *Oncogene.* 2010; 29: 3411-3422.
39. Fang L, Deng Z, Shatseva T, Yang J, Peng C, Du WW, Yee AJ, Ang LC, He C, Shan SW, Yang BB. MicroRNA miR-93 promotes tumor growth and angiogenesis by targeting integrin- β 8. *Oncogene.* 2011; 30:806-821.
40. Birks DK, Barton VN, Donson AM, Handler MH, Vibhakkar R, Foreman NK. Survey of MicroRNA expression in pediatric brain tumors. *Pediatr Blood Cancer.* 2011; 56: 211-216.
41. Malzkorn B, Wolter M, Liesenberg F, Grzendowski M, Stuhler K, Meyer HE, Reifenberger G. Identification and functional characterization of microRNAs involved in the malignant progression of gliomas. *Brain Pathol.* 2010; 20: 539-550.
42. Ivanovska I, Ball AS, Diaz RL, Magnus JF, Kibukawa M, Schelter JM, Kobayashi SV, Lim L, Burchard J, Jackson AL, Linsley PS, Cleary MA. MicroRNAs in the miR-106b family regulate p21/CDKN1A and promote cell cycle progression. *Mol Cell Biol.* 2008; 28: 2167-2174.
43. Petrocca F, Vecchione A, Croce CM. Emerging role of miR-106b-25/miR-17-92 clusters in the control of transforming growth factor beta signaling. *Cancer Res.* 2008; 68: 8191-8194.
44. Petrocca F, Visone R, Onelli MR, Shah MH, Nicoloso MS, de Martino I, Iliopoulos D, Pillozzi E, Liu CG, Negrini M, Cavazzini L, Volinia S, Alder H, et al. E2F1-regulated microRNAs impair TGF beta-dependent cell-cycle arrest and apoptosis in gastric cancer. *Cancer Cell.* 2008; 13: 272-286.
45. Pichiorri F, Suh SS, Ladetto M, Kuehl M, Palumbo T, Drandi D, Taccioli C, Zanesi N, Alder H, Hagan JP, Munker R, Volinia S, Boccadoro M, et al. MicroRNAs regulate critical genes associated with multiple myeloma pathogenesis. *Proc Natl Acad Sci U S A.* 2008; 105: 12885-12890.
46. Singh SK, Kagalwala MN, Parker-Thornburg J, Adams H, Majumder S. REST maintains self-renewal and pluripotency of embryonic stem cells. *Nature.* 2008; 453: 223-227.
47. Ballas N, Grunseich C, Lu DD, Speh JC, Mandel G. REST and its corepressors mediate plasticity of neuronal gene chromatin throughout neurogenesis. *Cell.* 2005; 121: 645-657.
48. Hebert SS, Horre K, Nicolai L, Bergmans B, Papadopoulou AS, Delacourte A, De Strooper B. MicroRNA regulation of Alzheimer's Amyloid precursor protein expression. *Neurobiol Dis.* 2009; 33: 422-428.
49. Caviness VS, Jr., Nowakowski RS, Bhide PG. Neocortical neurogenesis: morphogenetic gradients and beyond. *Trends Neurosci.* 2009; 32: 443-450.
50. Zhao CM, Deng W, Gage FH. Mechanisms and functional implications of adult neurogenesis. *Cell.* 2008; 132: 645-660.
51. Tanzer A, Stadler PF. Molecular evolution of a microRNA cluster. *J Mol Biol.* 2004; 339: 327-335.
52. Ventura A, Young AG, Winslow MM, Lintault L, Meissner A, Erkland SJ, Newman J, Bronson RT, Crowley D, Stone JR, Jaenisch R, Sharp PA, Jacks T. Targeted deletion reveals essential and overlapping functions of the miR-17-92 family of miRNA clusters. *Cell.* 2008; 132: 875-886.
53. Hewitt CA, Ling K-H, Merson TD, Simpson KM, Ritchie ME, King SL, Pritchard MA, Smyth GK, Thomas T, Scott HS, Voss AK. Gene network disruptions and neurogenesis defects in the adult Ts1Cje mouse model of Down syndrome. *PLoS One.* 2010; 5: Article No.: e11561.
54. Jacobs FM, van der Heide LP, Wijchers PJ, Burbach JP, Hoekman MF, Smidt MP. FoxO6, a novel member of the FoxO class of transcription factors with distinct shuttling dynamics. *J Biol Chem.* 2003; 278: 35959-35967.
55. Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, Hu LS, Anderson MJ, Arden KC, Blenis J, Greenberg ME. Akt promotes cell survival by phosphorylating and inhibiting a forkhead transcription factor. *Cell.* 1999; 96: 857-868.
56. Salih DA, Brunet A. FoxO transcription factors in the maintenance of cellular homeostasis during aging. *Curr Opin Cell Biol.* 2008; 20: 126-136.
57. Partridge L, Bruning JC. Forkhead transcription factors and ageing. *Oncogene.* 2008; 27: 2351-2363.
58. Hwangbo DS, Gershman B, Tu MP, Palmer M, Tatar M. Drosophila dFOXO controls lifespan and regulates insulin signalling in brain and fat body. *Nature.* 2004; 429: 562-566.

59. Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A *C. elegans* mutant that lives twice as long as wild type. *Nature*. 1993; 366: 461-464.
60. Boehm M, Slack F. A developmental timing microRNA and its target regulate life span in *C. elegans*. *Science*. 2005; 310: 1954-1957.
61. Anselmi CV, Malovini A, Roncarati R, Novelli V, Villa F, Condorelli G, Bellazzi R, Puca AA. Association of the FOXO3A locus with extreme longevity in a southern Italian centenarian study. *Rejuvenation Res*. 2009; 12: 95-103.
62. Flachsbarth F, Caliebec A, Kleindorp R, Blanche H, von Eller-Eberstein H, Nikolaus S, Schreiber S, Nebel A. Association of FOXO3A variation with human longevity confirmed in German centenarians. *Proc Natl Acad Sci U S A*. 2009; 106: 2700-2705.
63. Li Y, Wang WJ, Cao HQ, Lu JH, Wu C, Hu FY, Guo J, Zhao L, Yang F, Zhang YX, Li W, Zheng GY, Cui HB, et al. Genetic association of FOXO1A and FOXO3A with longevity trait in Han Chinese populations. *Hum Mol Genet*. 2009; 18: 4897-4904.
64. Pawlikowska L, Hu DL, Huntsman S, Sung A, Chu C, Chen J, Joyner AH, Schork NJ, Hsueh WC, Reiner AP, Psaty BM, Atzmon G, Barzilai N, et al. Association of common genetic variation in the insulin/IGF1 signaling pathway with human longevity. *Aging Cell*. 2009; 8: 460-472.
65. Willcox BJ, Donlon TA, He Q, Chen R, Grove JS, Yano K, Masaki KH, Willcox DC, Rodriguez B, Curb JD. FOXO3A genotype is strongly associated with human longevity. *Proc Natl Acad Sci U S A*. 2008; 105: 13987-13992.
66. Gritti A, Parati EA, Cova L, Frolichsthal P, Galii R, Wanke E, Faravelli L, Morassutti DJ, Roisen F, Nickel DD, Vescovi AL. Multipotential stem cells from the adult mouse brain proliferate and self-renew in response to basic fibroblast growth factor. *J Neurosci*. 1996; 16: 1091-1100.
67. Johe KK, Hazel TG, Muller T, Dugich-Djordjevic MM, McKay RDG. Single factors direct the differentiation of stem cells from the fetal and adult central nervous system. *Genes Dev*. 1996; 10: 3129-3140.
68. Lewis BP, Shih IH, Jones-Rhoades MW, Bartel DP, Burge CB. Prediction of mammalian microRNA targets. *Cell*. 2003; 115: 787-798.
69. Thomas PD, Campbell MJ, Kejariwal A, Mi HY, Karlak B, Daverman R, Diemer K, Muruganujan A, Narechania A. PANTHER: A library of protein families and subfamilies indexed by function. *Genome Res*. 2003; 13: 2129-2141.
70. Thomas PD, Kejariwal A, Guo N, Mi HY, Campbell MJ, Muruganujan A, Lazareva-Ilitsky B. Applications for protein sequence-function evolution data: mRNA/protein expression analysis and coding SNP scoring tools. *Nucleic Acids Res*. 2006; 34: W645-W650.
71. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005; 102: 15545-15550.
72. Papadopoulos GL, Alexiou P, Maragkakis M, Reczko M, Hatzigeorgiou AG. DIANA-mirPath: Integrating human and mouse microRNAs in pathways. *Bioinformatics*. 2009; 25: 1991-1993.
73. Maragkakis M, Reczko M, Simossis VA, Alexiou P, Papadopoulos GL, Dalamagas T, Giannopoulos G, Goumas G, Koukis E, Kourtis K, Vergoulis T, Koziris N, Sellis T, et al. DIANA-microT web server: elucidating microRNA functions through target prediction. *Nucleic Acids Res*. 2009; 37: W273-276.
74. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res*. 2000; 28: 27-30.
75. Panchision DM. The role of oxygen in regulating neural stem cells in development and disease. *J Cell Physiol*. 2009; 220: 562-568.
76. Meletis K, Wirta V, Hede SM, Nister M, Lundeberg J, Frisen J. p53 suppresses the self-renewal of adult neural stem cells. *Development*. 2006; 133: 363-369.
77. Luo CX, Jin X, Cao CC, Zhu MM, Wang B, Chang L, Zhou QG, Wu HY, Zhu DY. Bidirectional regulation of neurogenesis by neuronal nitric oxide synthase derived from neurons and neural stem cells. *Stem Cells*. 2010; 28: 2041-2052.
78. Wachs FP, Winner B, Couillard-Despres S, Schiller T, Aigner R, Winkler J, Bogdahn U, Aigner L. Transforming growth factor-beta1 is a negative modulator of adult neurogenesis. *J Neuropathol Exp Neurol*. 2006; 65: 358-370.
79. Buckwalter MS, Yamane M, Coleman BS, Ormerod BK, Chin JT, Palmer T, Wyss-Coray T. Chronically increased transforming growth factor-beta1 strongly inhibits hippocampal neurogenesis in aged mice. *Am J Pathol*. 2006; 169: 154-164.
80. Groszer M, Erickson R, Scripture-Adams DD, Lesche R, Trumpp A, Zack JA, Kornblum HI, Liu X, Wu H. Negative regulation of neural stem/progenitor cell proliferation by the Pten tumor suppressor gene in vivo. *Science*. 2001; 294: 2186-2189.
81. Groszer M, Erickson R, Scripture-Adams DD, Dougherty JD, Le Belle J, Zack JA, Geschwind DH, Liu X, Kornblum HI, Wu H. PTEN negatively regulates neural stem cell self-renewal by modulating G(0)-G(1) cell cycle entry. *Proc Natl Acad Sci U S A*. 2006; 103: 111-116.
82. Sinor AD, Lillien L. Akt-1 expression level regulates CNS precursors. *J Neurosci*. 2004; 24: 8531-8541.
83. Rafalski VA, Brunet A. Energy metabolism in adult neural stem cell fate. *Prog Neurobiol*. 2010; 93: 182-203.
84. Shaw WM, Luo S, Landis J, Ashraf J, Murphy CT. The *C. elegans* TGF-beta Dauer pathway regulates longevity via insulin signaling. *Curr Biol*. 2007; 17: 1635-1645.
85. Yamazaki S, Iwama A, Takayanagi S, Eto K, Ema H, Nakauchi H. TGF-beta as a candidate bone marrow niche signal to induce hematopoietic stem cell hibernation. *Blood*. 2009; 113: 1250-1256.
86. Seoane J, Le HV, Shen L, Anderson SA, Massague J. Integration of Smad and forkhead pathways in the control of neuroepithelial and glioblastoma cell proliferation. *Cell*. 2004; 117: 211-223.

87. Costa A, Onesti S. The MCM complex: (just) a replicative helicase? *Biochem Soc Trans.* 2008; 36: 136-140.
88. Biggs WH, Cavennee WK, Arden KC. Identification and characterization of members of the FKHR (FOX O) subclass of winged-helix transcription factors in the mouse. *Mamm Genome.* 2001; 12: 416-425.
89. Furuyama T, Nakazawa T, Nakano I, Mori N. Identification of the differential distribution patterns of mRNAs and consensus binding sequences for mouse DAF-16 homologues. *Biochem J.* 2000; 349: 629-634.
90. Foshay KM, Gallicano GI. miR-17 family miRNAs are expressed during early mammalian development and regulate stem cell differentiation. *Dev Biol.* 2009;326:431-443.
91. Li Z, Yang CS, Nakashima K, Rana TM. Small RNA-mediated regulation of iPS cell generation. *EMBO J.* 2011; In Press.
92. Balordi F, Fishell G. Mosaic removal of hedgehog signaling in the adult SVZ reveals that the residual wild-type stem cells have a limited capacity for self-renewal. *J Neurosci.* 2007; 27: 14248-14259.
93. van Praag H, Shubert T, Zhao CM, Gage FH. Exercise enhances learning and hippocampal neurogenesis in aged mice. *J Neurosci.* 2005; 25: 8680-8685.
94. Kempermann G, Kuhn HG, Gage FH. Experience-induced neurogenesis in the senescent dentate gyrus. *J Neurosci.* 1998; 18: 3206-3212.
95. Jasper H, Jones DL. Metabolic regulation of stem cell behavior and implications for aging. *Cell Metab.* 2010; 12: 561-565.
96. Rando TA. Stem cells, ageing and the quest for immortality. *Nature.* 2006; 441: 1080-1086.
97. Wang H, Liu J, Zong Y, Xu Y, Deng W, Zhu H, Liu Y, Ma C, Huang L, Zhang L, Qin C. miR-106b aberrantly expressed in a double transgenic mouse model for Alzheimer's disease targets TGF-beta type II receptor. *Brain Res.* 2010; 1357: 166-174.
98. Krampert M, Chirasani SR, Wachs FP, Aigner R, Bogdahn U, Yingling JM, Heldin CH, Aigner L, Heuchel R. Smad7 regulates the adult neural stem/progenitor cell pool in a transforming growth factor beta- and bone morphogenetic protein-independent manner. *Mol Cell Biol.* 2010; 30:3685-3694.
99. Oldham S, Stocker H, Laffargue M, Wittwer F, Wymann M, Hafen E. The Drosophila insulin/IGF receptor controls growth and size by modulating PtdInsP(3) levels. *Development.* 2002; 129: 4103-4109.
100. Hill R, Wu H. PTEN, stem cells, and cancer stem cells. *J Biol Chem.* 2009; 284: 11755-11759.
101. Poliseno L, Salmena L, Riccardi L, Fornari A, Song MS, Hobbs RM, Sportoletti P, Varmeh S, Egia A, Fedele G, Rameh L, Loda M, Pandolfi PP. Identification of the miR-106b~25 microRNA cluster as a proto-oncogenic PTEN-targeting intron that cooperates with its host gene MCM7 in transformation. *Sci Signal.* 2010; 3: ra29.
102. Gomis RR, Alarcon C, He W, Wang Q, Seoane J, Lash A, Massague J. A FoxO-Smad synexpression group in human keratinocytes. *Proc Natl Acad Sci U S A.* 2006; 103:12747-12752.
103. Pevny L, Rao MS. The stem-cell menagerie. *Trends Neurosci.* 2003; 26: 351-359.
104. Brazel CY, Limke TL, Osborne JK, Miura T, Cai JL, Pevny L, Rao MS. Sox2 expression defines a heterogeneous population of neurosphere-forming cells in the adult murine brain. *Aging Cell.* 2005; 4: 197-207.
105. Chen CZ, Li L, Lodish HF, Bartel DP. MicroRNAs modulate hematopoietic lineage differentiation. *Science.* 2004; 303: 83-86.
106. Greer EL, Oskoui PR, Banko MR, Maniar JM, Gygi MP, Gygi SP, Brunet A. The energy sensor AMP-activated protein kinase directly regulates the mammalian FOXO3 transcription factor. *J Biol Chem.* 2007; 282: 30107-30119.