

# An AMPK-FOXO Pathway Mediates Longevity Induced by a Novel Method of Dietary Restriction in *C. elegans*

Eric L. Greer,<sup>1,2</sup> Dara Dowlatshahi,<sup>3</sup> Max R. Banko,<sup>1</sup> Judit Villen,<sup>5</sup> Kimmi Hoang,<sup>1</sup> Daniel Blanchard,<sup>1</sup> Steven P. Gygi,<sup>5</sup> and Anne Brunet<sup>1,2,4,\*</sup>

<sup>1</sup>Department of Genetics

<sup>2</sup>Cancer Biology Program

<sup>3</sup>Biological Sciences Graduate Program

<sup>4</sup>Neurosciences Program

Stanford University

Stanford, California 94305

<sup>5</sup>Department of Cell Biology

Harvard Medical School

Boston, Massachusetts 02115

## Summary

**Background:** Dietary restriction (DR) is the most effective environmental intervention to extend lifespan in a wide range of species. However, the molecular mechanisms underlying the benefits of DR on longevity are still poorly characterized. AMP-activated protein kinase (AMPK) is activated by a decrease in energy levels, raising the possibility that AMPK might mediate lifespan extension by DR.

**Results:** By using a novel DR assay that we developed and validated in *C. elegans*, we find that AMPK is required for this DR method to extend lifespan and delay age-dependent decline. We find that AMPK exerts its effects in part via the FOXO transcription factor DAF-16. FOXO/DAF-16 is necessary for the beneficial effects of this DR method on lifespan. Expression of an active version of AMPK in worms increases stress resistance and extends longevity in a FOXO/DAF-16-dependent manner. Lastly, we find that AMPK activates FOXO/DAF-16-dependent transcription and phosphorylates FOXO/DAF-16 at previously unidentified sites, suggesting a possible direct mechanism of regulation of FOXO/DAF-16 by AMPK.

**Conclusions:** Our study shows that an energy-sensing AMPK-FOXO pathway mediates the lifespan extension induced by a novel method of dietary restriction in *C. elegans*.

## Introduction

Dietary restriction (DR)—restriction in food intake without malnutrition—increases both the mean and maximal lifespan in a wide range of species [1]. The longevity effects of DR are conserved in worms, flies, mice, and possibly primates, underscoring the importance of this environmental intervention on lifespan. In humans, DR decreases the incidence of age-dependent diseases, including atherosclerosis and cancer [2, 3]. Although numerous hypotheses have been proposed to explain the benefits of DR on longevity [4], the exact mechanisms by

which DR increases lifespan are still unclear. DR decreases available nutrients [5], raising the possibility that nutrient-sensing pathways may mediate DR's beneficial effects on aging and age-dependent diseases.

The energy-sensing AMP-activated protein kinase (AMPK) is an attractive candidate to mediate DR's ability to prolong lifespan. AMPK is activated in response to decreased energy or glucose levels, which are consequences of DR [6], and in specific cases, by DR itself [7]. AMPK is a heterotrimeric protein kinase composed of a catalytic subunit ( $\alpha$ ) and two regulatory subunits ( $\beta$  and  $\gamma$ ) that sense low cellular energy levels by monitoring changes in the AMP:ATP ratio [8]. AMP binding to the  $\gamma$  subunit induces a conformational change that allows the catalytic  $\alpha$  subunit to be phosphorylated and activated by the AMPK-activating protein kinases LKB1 and CAMKK $\beta$  [9–13]. AMP also contributes to AMPK activation by reducing the dephosphorylation of the catalytic  $\alpha$  subunit [14]. AMPK coordinates the adaptation of cellular metabolism and growth to decreased energy levels [6]. One of the two  $\alpha$  catalytic subunits of AMPK (AMPK $\alpha$ 2 or AAK-2) has recently been found to increase worm longevity [15]. However, AAK-2 is not necessary for longevity extension in response to a mutation in the *eat-2* gene [16], a mutation that is thought to genetically mimic DR but might have pleiotropic effects. Therefore, whether AMPK mediates lifespan extension by DR is still unknown and the molecular mechanisms by which AMPK controls lifespan remain unclear.

FOXO transcription factors are good candidates to mediate AMPK's ability to extend lifespan. These transcriptional regulators play an important conserved role in stress resistance by upregulating a series of target genes, including genes involved in detoxification of reactive oxygen species (ROS) [17–21]. FOXO factors are also activated in response to decreased activity of the insulin signaling pathway, an alternate nutrient-sensing pathway in cells [22, 23]. In *C. elegans*, the FOXO transcription factor DAF-16 mediates the longevity extension resulting from a mutation in the insulin receptor *daf-2* [17, 18]. Although the lifespan extension resulting from the *eat-2* mutation and to several DR regimens has been shown to be independent of FOXO/DAF-16 [24–29], FOXO may still mediate lifespan extension under other DR experimental conditions.

Based on these observations, we hypothesized that AMPK and FOXO may be important mediators of the beneficial effects of DR on longevity. By using a novel method of DR in worms, we show that AMPK and FOXO/DAF-16 are necessary to mediate a large part of the lifespan extension in response to DR. In addition, we show that constitutively active AMPK increases worm stress resistance and longevity in a FOXO/DAF-16-dependent manner. Lastly, we find that FOXO/DAF-16 is a direct *in vitro* substrate of AMPK and that AMPK increases FOXO-dependent transcription. Together, these results place the AMPK-FOXO module as a central pathway in the regulation of lifespan by DR.

\*Correspondence: [anne.brunet@stanford.edu](mailto:anne.brunet@stanford.edu)

## Results

### Development of a New DR Method, sDR, in the Nematode *C. elegans*

We developed a new DR method in worms by transferring adult worms every 2 days onto freshly seeded plates with restrictive amounts of feeding bacteria (from  $5 \times 10^{11}$  to  $5 \times 10^8$  bacteria/ml) at day 4 of adulthood (Figure 1A). We verified that worms ingested fewer bacteria under these conditions by using bacteria expressing the red fluorescent reporter dsRed (Figure 1B). This restriction in bacterial intake significantly increased worm lifespan (Figure 1C; Table S1 in the Supplemental Data available online) and reduced worm mortality rate (Figure 1D), indicating that this regimen delays aging. Our regimen also increased resistance to the oxidative stress agent Paraquat (Figure 1E; Table S2A) and delayed the age-dependent decline in worm locomotor activity (Figure 1F), consistent with the reported ability of DR to delay aging and age-related phenotypes in a variety of organisms [1]. The increase in lifespan was not simply a result of reduced pathogenicity or toxicity of the bacteria because (1) an increase in bacterial concentration to  $5 \times 10^{12}$  bacteria/ml did not shorten worm lifespan (Figure 1C) and (2) dilution of UV-killed bacteria also extended worm lifespan, albeit to a lesser extent (Figure 1G; Table S2B). Taken together, these results indicate that the food-restriction regimen that we have developed in *C. elegans* increases lifespan in a manner that is similar to DR in other species. In the rest of the manuscript, ad libitum will be  $5 \times 10^{11}$  bacteria/ml and dietary restriction will be  $5 \times 10^8$  bacteria/ml. We will refer to this method as sDR for “solid DR,” as, contrary to other methods, the diluted bacteria are placed on solid plates.

### sDR Increases the AMP:ATP Ratio and Is Dependent on AMPK

Because AMPK is activated by low energy levels, we asked whether sDR's ability to extend lifespan was mediated by AMPK in *C. elegans*. We first examined whether the sDR regimen affected energy levels in worms by measuring the AMP:ATP ratio in worm extracts by using reverse-phase high-performance liquid chromatography (HPLC). We found that restricting the amount of seeded bacteria from  $5 \times 10^{11}$  bacteria/ml (ad libitum) to  $5 \times 10^8$  bacteria/ml (sDR) at day 4 of adulthood significantly increased the AMP:ATP ratio in worms (Figure 2A). These results indicate that sDR leads to decreased energy levels in worms, as expected for a diet that restricts available nutrients.

To determine whether AMPK mediated the ability of sDR to extend lifespan in worms, we used a worm strain carrying a deletion in the gene encoding one of the catalytic subunits of AMPK (AMPK  $\alpha 2$  or *aak-2*). The lifespan extension resulting from sDR was suppressed in the *aak-2* mutant (Figure 2B; Table S3), indicating that AMPK is necessary for the lifespan extension resulting from sDR. In addition, *aak-2* was also necessary for sDR to prevent the age-dependent decline in worm locomotor activity (Figure 2C). Together, these data show that AMPK is necessary for sDR to extend lifespan and to slow age-dependent decline.

### The FOXO Transcription Factor DAF-16 Mediates sDR-Induced Extension of Lifespan

Although FOXO transcription factors are attractive candidates to mediate the actions of environmental interventions controlling stress resistance and longevity, several alternate methods of DR in worms have been shown to be independent of FOXO/DAF-16 [24–29]. To test whether DAF-16 played a role in lifespan extension in response to sDR, we assessed the lifespan of wild-type or *daf-16* mutant worms in ad libitum or food-restriction conditions. We found that the ability of sDR to extend worm lifespan was diminished in *daf-16* mutants (Figure 3A; Table S4A). These results indicate that DAF-16, like AMPK, mediates a portion of DR's ability to extend lifespan. The ability of DR to delay the age-dependent decline in locomotor activity was also diminished in *daf-16* mutant worms (Figure 3B), indicating that DAF-16 is necessary for the ability of DR to prevent the age-dependent decline in fitness.

FOXO's role in sDR-induced increase in lifespan may be dependent on the insulin signaling pathway, a well-known regulator of FOXO activity and a sensor of food levels [17, 18]. To test this possibility, we asked whether the insulin receptor *daf-2* also mediated the beneficial effects of sDR. In contrast to *aak-2* and *daf-16*, the long lifespan of *daf-2* insulin receptor mutants was further extended by sDR (Figure 3C; Table S4B), suggesting that sDR may regulate DAF-16 in a manner that is not dependent on the insulin receptor pathway. However, because the *daf-2* mutation is not a null mutation, it is still possible that part of the sDR-induced increase in lifespan might depend on the insulin receptor pathway.

Several known regulators of FOXO/DAF-16, including mutation in the insulin receptor or complete starvation, trigger the nuclear localization of FOXO/DAF-16 [30]. We therefore examined whether sDR triggers nuclear localization of DAF-16. Unlike starvation, sDR did not stimulate DAF-16 translocation to the nucleus (Figure 3D). Taken together, these results indicate that sDR regulates DAF-16 in a way that is distinct from starvation or mutation in the insulin receptor. These observations suggest that FOXO/DAF-16 may be regulated by AMPK in response to sDR.

### FOXO/DAF-16 Mediates the Ability of AMPK to Induce Stress Resistance and Longevity in Worms

AMPK function in worms was previously suggested to be independent of FOXO/DAF-16 [15], but the importance of FOXO as a mediator of AMPK action on lifespan has not been directly addressed. To test the possibility that FOXO/DAF-16 might mediate AMPK's role in controlling lifespan, we generated transgenic worms expressing a constitutively active (CA) mutant of worm AMPK  $\gamma 2$  (R81Q). The corresponding mutation in the AMP binding pocket of AMPK  $\gamma 2$  in humans causes the constitutive activation of AMPK [31]. To determine whether CA AMPK  $\gamma 2$  was sufficient to promote organismal resistance to oxidative stress, a phenotype tightly associated with increase in lifespan, we performed oxidative stress resistance assays, via the ROS generator Paraquat. Three independent CA AMPK  $\gamma 2$  transgenic worm lines (CA 1-3) were more resistant than control worm lines (EV 1-3) to increasing concentrations of Paraquat (Figure 4A) or to increasing times of treatment




Figure 1. sDR: A Novel Dietary Restriction Method in *C. elegans*

(A) Description of our dietary restriction method (sDR) in *C. elegans*. Populations of worms obtained by synchronized egg laying (EL) were scored and switched to freshly seeded plates every other day (indicated by asterisk). sDR was initiated at day 4 of adulthood. AL, ad libitum.

(B) Representative pictures of worms fed different concentrations of bacteria expressing DsRed, a red fluorescent protein. Right: quantification of DsRed fluorescence of wild-type (N2) worms incubated with different bacterial dilutions. Results represent the average and SEM of 3 independent experiments performed with 30 worms per condition at day 7. \*\*\* $p < 0.001$  in one-way ANOVA. AL,  $5 \times 10^{11}$  bacteria/ml; sDR,  $5 \times 10^8$  bacteria/ml.

(C) Serial dilution of OP50-1 bacteria increases worm lifespan ( $5 \times 10^8$  bacteria/ml = 28.9% increase over  $5 \times 10^{11}$  bacteria/ml). Mean and standard errors for this experiment done in triplicate are presented in Table S1.

(D) Gompertz analysis shows that sDR ( $5 \times 10^8$  bacteria/ml) decreases the worm mortality rate.

(E) sDR-treated worms displayed a 36% increase in survival compared with AL-fed worms at 200 mM Paraquat ( $p < 0.0001$ ). The mean and SEM for duplicate experiments done in sextuplicate is presented in Table S2A.




Figure 2. AMPK Is Necessary for sDR's Beneficial Effects on Lifespan and Age-Dependent Decline

(A) The AMP:ATP ratio, measured by reverse-phase HPLC, increases when worms are dietarily restricted at day 7. The results presented are the mean and SEM of two independent experiments of 5 samples each. \*\**p* = 0.0079 by Mann-Whitney test.

(B) AMPK mediates sDR's ability to extend lifespan. *aak-2* mutant worms displayed a mild decrease in lifespan compared to WT (N2) worms in ad libitum (AL) conditions ( $5 \times 10^{11}$  bacteria/ml) (10.5%; *p* < 0.0001). Under sDR conditions ( $5 \times 10^8$  bacteria/ml), *aak-2* mutant worms had a 48% shorter lifespan compared to WT (N2) worms (*p* < 0.0001). Comparison of five independent experiments performed in triplicate is presented in Table S3.

(C) sDR delays the age-dependent decline in locomotor activity in an *aak-2*-dependent manner. The experiment was performed as described in the legend of Figure 1F. \**p* < 0.05; \*\**p* < 0.01; \*\*\**p* < 0.001 in one-way ANOVA. Note that N2 worms are also represented in Figure 1F.

with Paraquat (Figure 4B; Table S5A). The stress protective effects of CA AMPK  $\gamma 2$  were reverted in worms in which *daf-16* was knocked-down by RNAi (Figures 4C and 4D; Table S5A), indicating that the ability of this constitutive form of AMPK to induce stress resistance is DAF-16 dependent.

To examine whether DAF-16 was also required for AMPK's ability to regulate lifespan, we examined the lifespan of CA AMPK  $\gamma 2$  transgenic worms in the absence or presence of *daf-16*. Consistent with the previous finding that AAK-2 (AMPK  $\alpha 2$ ) overexpression slightly extends worm lifespan [15], two independent CA AMPK  $\gamma 2$  transgenic lines (CA2 and CA3) displayed a modest, but statistically significant, increase in lifespan (Figure 4E; Table S5B). The increased longevity of CA AMPK  $\gamma 2$  transgenic worms (CA3) was reverted when DAF-16 expression was inhibited by RNAi (Figure 4F; Table S5B), indicating that AMPK's ability to extend lifespan requires DAF-16. Together, these results show that the worm FOXO factor DAF-16 is necessary to mediate the effects of AMPK on oxidative stress resistance and longevity.

#### AMPK Activates FOXO/DAF-16-Dependent Transcription

To gain insight into the mechanism of AMPK's action on FOXO/DAF-16 transcription factor, we examined the effect of AMPK activity on the expression of *sod-3*, a known DAF-16 target gene involved in both stress resistance and longevity [20, 32]. By using quantitative RT-PCR, we found that *sod-3* mRNA was increased in worms expressing the constitutively active form of AMPK (CA3) compared to control worms (EV1) and that this increase was dependent on the expression of DAF-16 (Figure 5A). Consistent with this observation,

*sod-3* mRNA expression was also decreased in AMPK  $\alpha 2$  (*aak-2*) mutant worms compared to wild-type worms (Figure 5B). These findings support the notion that AMPK enhances the ability of DAF-16 to transactivate *sod-3*.

#### AMPK Directly Phosphorylates DAF-16 In Vitro

AMPK could regulate FOXO/DAF-16 activity indirectly, via the insulin receptor, or directly, via phosphorylation. Given that sDR's beneficial effects on longevity appear to be independent of the insulin receptor, we asked whether AMPK could directly phosphorylate FOXO/DAF-16. To this end, we performed in vitro kinase assays with purified AMPK and a bacterially expressed form of DAF-16 as a substrate. We found that both worm and human AMPK phosphorylated worm DAF-16 (Figure 6A). DAF-16 phosphorylation by AMPK was greatly enhanced by the presence of AMP (Figure 6A), confirming that the purified kinase responsible for DAF-16 phosphorylation is AMP dependent. The stoichiometry of phosphorylation of DAF-16 by human and worm AMPK was higher than that of the SAMS peptide, a peptide containing a known AMPK phosphorylation site [13] (Figure 6B), suggesting that DAF-16 is a good in vitro substrate of AMPK.

We sought to determine the residues of FOXO/DAF-16 that were phosphorylated by AMPK. By using a combination of tandem mass spectrometry and site-directed mutagenesis, we found that AMPK phosphorylated DAF-16 in vitro at least at six residues—T166, S202, S314, S321, T463, and S466—that conform to varying degrees to the consensus motif phosphorylated by AMPK (amphipathic helix- $\beta\phi X\beta\phi X S/TXX\beta\phi\phi$ ;  $\phi$ , hydrophobic residues;  $\beta$ , basic residues) [33] (Figure 6C). The concomitant mutation of T166, S314, and S466

(F) sDR delays the age-dependent decline in worm locomotor activity. Quantification of the length of worm tracks after 1 hr on fresh ad libitum lawns of OP50-1 bacteria was performed with Image J. Each point represents the average movement and SEM of at least 20 worms. \**p* < 0.05; \*\**p* < 0.01; \*\*\**p* < 0.001 in one-way ANOVA.

(G) sDR induced by dilution of UV-killed bacteria results in a 15.9% increase in lifespan of N2 worms (statistically different from ad libitum fed worms with a *p* = 0.0004). The mean and SEM of two independent experiments conducted in triplicate is presented in Table S2B.






Figure 3. FOXO/DAF-16 Is Necessary for sDR's Beneficial Effects on Lifespan and Age-Dependent Decline

(A) DAF-16 is necessary for sDR to extend lifespan. sDR leads to a 30.6% increase ( $p < 0.0001$ ) in longevity in wild-type (N2) worms but does not significantly affect the lifespan of *daf-16* mutant worms. The mean and SEM of three independent experiments conducted in triplicates of 30 worms each is presented in Table S4A.

(B) sDR delays the age-dependent decline in locomotor activity in a *daf-16*-dependent manner. The experiment was performed as described in the legend of Figure 1F. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$  in one-way ANOVA. Note that N2 worms are also represented in Figure 1F.

(C) sDR is not dependent on the insulin receptor *daf-2*. sDR induced a 35.2% increase ( $p < 0.0001$ ) in wild-type (N2) worms and a 31.0% increase ( $p < 0.0001$ ) in *daf-2* mutant worms whereas sDR did not increase the lifespan of *aak-2* mutant worms. The mean and SEM of two independent experiments conducted in triplicates of 30 worms each is presented in Table S4B.

(D) sDR does not cause DAF-16 nuclear localization. DAF16::GFP-expressing worms were placed on ad libitum or sDR diet for 6 hr at day 7 of life.

significantly reduced the phosphorylation of DAF-16 by AMPK, indicating that these three sites are the main residues phosphorylated by AMPK in vitro (Figure 6D). The replacement of all six residues (T166, S202, S314, S321, T463, and S466) by alanine further reduced the phosphorylation of DAF-16 by AMPK, suggesting that S202, S321, and T463 are also phosphorylated by AMPK (Figure 6D). Taken together with our experiments on the effect of AMPK on *sod-3*, these findings suggest that AMPK could activate FOXO/DAF-16-dependent transcription via direct phosphorylation at these previously unidentified sites. However, our results do not exclude the possibility that AMPK may also regulate FOXO/DAF-16 indirectly. We have recently found that AMPK phosphorylates the human FOXO transcription factor FOXO3 at multiple sites in vitro and in cells [34], raising the possibility that the direct connection between AMPK and FOXO may be conserved throughout evolution.

## Discussion

### sDR: A Novel Method of DR that Extends Lifespan in Worms

We have developed a novel method of dietary restriction in *C. elegans* that we termed sDR. sDR has hallmarks of

dietary restriction in many species in that it (1) reduces food intake; (2) reproducibly extends lifespan; (3) delays the age-dependent decline in physical fitness; and (4) induces resistance to oxidative stress. Another well-known hallmark of dietary restriction is a reduction in fertility. sDR is initiated at day 4 of adulthood, which is past the reproductive period in worms. We found that if sDR was initiated earlier, it indeed reduced fertility, but also had deleterious effects on survival (E.L.G. and A.B., unpublished data). The fact that sDR extends lifespan at a postreproductive age suggests that sDR extends lifespan independently of changes in fertility.

The other methods that have been used in *C. elegans* to induce or mimic DR include the *eat-2* mutation, which reduces pharyngeal pumping [29], the dilution of feeding bacteria in liquid culture (bDR or IDR) [26–28, 35], the total absence of bacteria in liquid culture (axenic medium) [26], a chemically defined liquid media [36], the dilution of peptone in the plates [37], and the total absence of food on plates (DD for dietary deprivation) [24, 25] (Figure 7). Compared to these other methods, sDR is relatively simple to implement because it is done in the medium commonly used for worms in the laboratory and it can be performed with any mutant strain of worm. sDR is also similar to regimens used in other species, including




Figure 4. FOXO/DAF-16 Is Necessary for AMPK-Induced Increase in Stress Resistance and Lifespan

(A) Constitutively active (CA) AMPK  $\gamma$ 2 transgenic worms display an increased resistance to increasing concentrations of the reactive oxygen species generator Paraquat compared to control transgenic worms injected with an empty vector (EV) after 7 hr of incubation. Three independent lines of EV (1-3) and CA (1-3) were analyzed.

(B) CA3 displayed an 86.0% increased resistance to 300 mM Paraquat over control EV1 ( $p < 0.0001$ ). The mean and SEM of two independent experiments conducted in sextuplicate is presented in Table S5A.

(C) Stress resistance of CA AMPK  $\gamma$ 2 transgenic worms is dependent on the presence of DAF-16. CA3 worms were fed bacteria expressing RNAi to *daf-16* or control bacteria (ctl) and submitted to increased concentrations of Paraquat for 8 hr.

(D) CA3 displayed a 48.9% increased stress resistance compared to EV1 ( $p = 0.0005$ ) but did not significantly increase stress resistance when treated with RNAi to *daf-16*. The mean and SEM of two independent experiments conducted in sextuplicate is presented in Table S5A.

(E) CA AMPK  $\gamma$ 2 transgenic worms (CA2 and CA3) displayed a statistically significant increase in median lifespan (9.5% and 17.7%, respectively,  $p < 0.001$ ) compared to EV1 control worms. A comparison experiment done in triplicate is presented in Table S5B.

(F) CA AMPK  $\gamma$ 2 lifespan extension is DAF-16 dependent. CA AMPK  $\gamma$ 2 transgenic worms (CA3) show a statistically significant increase in median lifespan (11%;  $p = 0.0006$ ) compared to control transgenic worms (EV1) but did not display a significant lifespan extension compared to EV1 when treated with RNAi to DAF-16. Comparison of two independent experiments done in triplicate is presented in Table S5B.

mammals, in which DR is achieved by a reduction of the food normally provided to the animals.

#### sDR Reveals the Importance of AMPK and FOXO in Dietary Restriction-Induced Longevity

Our findings suggest the following sequence of events. sDR activates AMPK by increasing the AMP:ATP ratio. Activated AMPK enhances FOXO/DAF-16-dependent transcription, perhaps via direct phosphorylation, leading to resistance to oxidative stress and longevity

extension (Figure 7). The finding that sDR's ability to extend lifespan is dependent on the presence of AMPK/AAK-2 and FOXO/DAF16 is in contrast to numerous studies in worms that indicate that *aak-2* and *daf-16* do not mediate the lifespan extension observed in *eat-2* mutant [16, 29] and that *daf-16* does not mediate the lifespan extension by IDR [27], axenic medium [26], DD [24, 25], or bDR, particularly when bacteria are diluted less than 10-fold from ad libitum conditions [26, 28]. However, our results are consistent with the observation




Figure 5. AMPK Activates FOXO/DAF-16-Dependent Transcription of *sod-3*

(A) *sod-3* mRNA is increased in CA AMPK  $\gamma 2$  transgenic worms (CA3) compared with empty vector control transgenic line (EV1), and this increase is *daf-16* dependent. The results presented correspond to the mean and SEM of three independent quantitative RT-PCR experiments. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$  in a one-way ANOVA statistical test.

(B) *sod-3* mRNA was decreased in *aak-2* mutant worms compared with N2 wild-type worms. The results presented correspond to the mean and SEM of two independent experiments: \* $p < 0.05$  in a paired t test.

that DR elicited by a chemically defined liquid media resulted in the upregulation of *daf-16* and *daf-16*-dependent gene expression [36] and the observation that the lifespan extension by bDR when bacteria are diluted more than 10-fold from ad libitum conditions shows a partial *daf-16* dependency [26, 28].

One possible interpretation of these contradictory findings is that various methods of DR elicit distinct signaling pathways because they restrict food intake in different manners. For example, the degree of food restriction and whether food restriction is chronic or acute could influence the mechanisms underlying longevity caused by DR. *eat-2* mutants, which have a germline mutation in the acetylcholine receptor that reduces their pharyngeal pumping rate, likely induce a form of DR that is chronic [38], whereas sDR, by diluting the bacteria 1000-fold at a specific adult stage, may elicit a more acute form of DR. Different DR methods may also alter proteins, carbohydrates, or lipids in different proportions, which may engage distinct mechanisms of nutrient sensing.

The differences in the mechanisms relating sDR versus other liquid DR regimens may also be due to other

parameters that differ between these two methods, including movement, energy expenditure, food sensing, oxygen levels, bacterial amounts, or osmotic conditions. Finally, some DR regimens may also extend lifespan by reducing some kind of food-related toxicity. However, sDR is unlikely to operate solely via reduced food toxicity, because an excess of bacteria is not detrimental for lifespan and because a dilution of UV-killed bacteria also extends lifespan. Dissecting the mechanisms by which different methods of DR extend lifespan and understanding the parameters that vary between the DR regimens will be important for designing efficient interventions to extend lifespan.

#### Molecular Mechanisms Mediating DR in Worms

Interestingly, recent evidence indicates that the ability of *eat-2* mutants and bDR to extend lifespan is mediated by a member of the Forkhead family, the FOXA/PHA-4 transcription factor [28]. IDR has also been recently shown to be sensed specifically in sensory neurons by the transcription factor SKN-1 [27]. Finally, the *sir-2* gene has been shown to be important to mediate certain forms of DR in a variety of species from yeast to flies [39–42], though this observation is still subject to debate [24, 25, 43, 44]. SIR-2, an NAD-dependent protein deacetylase, may sense dietary restriction signals via alterations of the NAD:NADH ratio whereas AMPK may sense DR through changes in the AMP:ATP ratio. It will be interesting to determine whether SIR-2, PHA-4, and SKN-1 are important to mediate lifespan extension by sDR. Uncovering the genetic interactions between the SIR-2, the AMPK-FOXO, the FOXA, and the SKN-1 pathways will give important insight into the network of longevity regulation by DR (Figure 7).

#### AMPK Increases Lifespan and Stress Resistance via FOXO-Dependent and FOXO-Independent Mechanisms

We have shown that AMPK regulates lifespan in a *daf-16*-dependent manner. However, Apfeld et al. reported that mutating one of the AMPK catalytic subunits, *aak-2*, in conjunction with mutating *daf-16* decreased worm lifespan further than mutating each gene individually [15]. This experiment raises the possibility that DAF-16 may not be the sole contributor of AMPK's effect on lifespan. AMPK is known to regulate the TOR pathway in mammals [45], and mTOR/LET-363 and S6 kinase mutations have been found to extend lifespan in yeast, worms, and flies [44, 46–49]. Thus, the mTOR-S6 kinase pathway may also mediate part of AMPK's ability to extend lifespan (Figure 7). Changes in transcription (via FOXO) and translation (via TOR) may be both orchestrated by AMPK to allow the organism to adapt its longevity to low energy levels. Lastly, AMPK may also regulate the other proteins recently identified as important mediators of DR, including SIR-2, PHA-4, and SKN-1.

#### AMPK Phosphorylation of DAF-16: A Mechanism of Regulation?

The direct phosphorylation of DAF-16 by AMPK may be one of the mechanisms by which DR regulates longevity in worms. DAF-16 is phosphorylated by AMPK at multiple novel sites that are located in the DNA binding




Figure 6. AMPK Directly Phosphorylates FOXO/DAF-16 In Vitro

(A) AMPK phosphorylates DAF-16 in an AMP-dependent manner. Results with human AMPK are presented, but both human and worm AMPK phosphorylate DAF-16 in an AMP-dependent manner.  
 (B) Stoichiometry of phosphorylation of DAF-16 by human AMPK. The graph represents the mean and standard deviation of normalized molar units of phosphate incorporated per mole of substrate of 3 independent experiments. Similar results were obtained with the worm AMPK.  
 (C) Schematic of DAF-16 showing the location of the residues of DAF-16 phosphorylated by AMPK in vitro. DBD, DNA-binding domain. Alignment of the AMPK consensus phosphorylation motif with phosphorylation sites in DAF-16.  $\phi$ , hydrophobic residues;  $\beta$ , basic residues. The sequence in parentheses represents an amphipathic helix. The sites in italics were identified as being phosphorylated in vitro by AMPK by tandem mass spectrometry. All the sites listed here were confirmed as being phosphorylated in vitro by AMPK by site-directed mutagenesis.  
 (D) DAF-16 mutants display a significant reduction in phosphorylation by AMPK compared to WT DAF-16 in an in vitro kinase assay. 3A, T166A/S314A/ S466A mutant; 6A, T166A/S202A/S314A/S321A/T463A/S466A.

domain and the C-terminal transactivation domain, and these sites differ from AKT phosphorylation sites. Therefore, AMPK may regulate DAF-16 independently of its subcellular localization, consistent with our observation that sDR, which activates AMPK, does not stimulate DAF-16 nuclear translocation. Our experiments do not rule out, however, the possibility that AMPK regulates DAF-16 via an indirect mechanism or that AMPK and DAF-16 act in different tissues to regulate longevity in response to DR.

### An AMPK-FOXO Pathway in Other Species

Whether AMPK and FOXO play a role in DR-induced longevity in other species is not known. We have recently found that AMPK phosphorylates human FOXO3 in mammalian cells at novel regulatory sites that are distinct from the AKT sites [34]. Similar to our results in worms, AMPK also activates FOXO3-dependent transcription in mammalian cells [34], raising the possibility that the mechanism of regulation of FOXO by AMPK may be conserved.

A number of compounds that activate AMPK, including metformin, phenformin, and resveratrol, have been proposed to act as potential “caloric restriction mimetics” in mammals [50, 51]. It will be interesting to test whether these compounds affect lifespan in an AMPK-dependent manner. Understanding the mechanisms underlying the conserved extension of lifespan by DR will help harness the power of this intervention to slow aging and the onset of age-related diseases.

### Experimental Procedures

#### Worm Strains and RNA Interference

N2 and *daf-16 (mu86)* strains were a kind gift from M.-W. Tan. The *zls356 IV (TJ356) (daf-16::GFP)* strain was a kind gift from T. Johnson. The *aak-2 (ok524)* and *daf-2 (e1370)* strains were generously provided by T. Stiernagle at the *Caenorhabditis* Genetics Center. HT115 (DE3) bacteria transformed with vectors expressing dsRNA of the genes of interest were obtained from the Ahringer library (a gift from M.-W. Tan) and were grown at 37°C and seeded onto standard nematode growth medium (NGM) plates [52] containing ampicillin (100  $\mu$ g/ml) and IPTG (0.4 mM). Adult worms were placed on these plates and removed after 4–6 hr to obtain synchronized populations of worms. Each vector was sequenced to verify the presence of the appropriate gene of interest.

#### Worm Transgenic Lines

The PRKAG2 (AMPK $\gamma$ 2) gene (WBGene00020633/WP:CE29350) and regulatory regions were amplified by PCR from wild-type N2 worm genomic DNA and was subcloned into the L3691 pPD117.01 vector (a kind gift from A. Fire). To generate a constitutively active mutant, we replaced Arg81 by Glu (R81Q mutant) [31] by site-directed mutagenesis. Purified plasmids of interest (the L3691 vector without the coding region of AMPK $\gamma$ 2, L3691 vector encoding WT AMPK $\gamma$ 2, or L3691 vector encoding the R81Q mutant of AMPK $\gamma$ 2) were coinjected with *pha-1* encoding plasmid into *pha-1* mutant worms. Worms were placed at 24°C after injection and were allowed to lay eggs to rescue the *pha-1* phenotype. Individual F1 strains were selected from various injections and were clonally expanded. Incorporation of extrachromosomal arrays was verified by PCR of genomic DNA isolated from worms.

#### Lifespan in *C. elegans*

Worm lifespan assays were performed at 20°C. Worm populations were synchronized by placing young adult worms on NGM plates






Figure 7. Proposed Model for the Molecular Networks Integrating Different DR Methods

Different methods of dietary restriction may activate distinct signaling pathways or activate them to varying degrees. The table below the figure summarizes which pathways have been tested. The red question mark indicates an interaction identified in mammals, but not yet confirmed in *C. elegans*. sDR, solid plate DR; bDR, liquid DR described in [28, 35]; IDR, liquid DR described in [27]; DD, dietary deprivation [24, 25]; axenic, absence of bacteria in liquid culture [26]; peptone, dilution of peptone in plates [37]; CDLM, chemically defined liquid media [36]. D, dependency; I, independency; D/I?, dependency or independency based on different reports in the literature [42, 44, 53]; NT, not tested. Single asterisk: note that the lifespan extension by bDR when bacteria are diluted more than 10-fold from ad libitum conditions shows a partial *daf-16* dependency [26, 28]. Double asterisk: note that the *daf-2* mutation is not a null mutation, which could affect the interpretation of the results. *clk-1*, mutant of a gene necessary for ubiquinone biosynthesis [29].

seeded with the *E. coli* strain OP50-1 (unless otherwise noted) for 4–6 hr and then removed. The hatching day was counted as day one for all lifespan measurements. Worms were changed every other day to new plates to eliminate confounding progeny and were marked as dead or alive. Worms were scored as dead if they did not respond to repeated prods with a platinum pick. Worms were censored if they crawled off the plate or died from vulval bursting. For each lifespan assay, 90 worms were used in 3 plates (30 worms/plate). The data were plotted with the Kaplan-Meier Survival curves, and statistical significance was determined by Logrank (Mantel-Cox) tests. Lifespan assays were repeated at least once and showed similar trends in relative lifespan effects. Representative Kaplan-Meier survival curves are shown in the figures. Means, SEM, and p values are shown in Tables S1–S5.

#### Dietary Restriction in *C. elegans*

OP50-1 bacteria were serially diluted from  $5 \times 10^{12}$  bacteria/ml to  $5 \times 10^4$  bacteria/ml. Bacteria were resuspended in S Medium to inhibit bacterial growth. 150  $\mu$ l of these solutions were plated on each 35 mm plate on the day of transfer. 30 adult worms were placed on each plate containing various concentrations of bacteria starting at day 7 of life (day 4 of adulthood). Worms were switched to fresh

plates every other day. Worms placed on bacterial concentrations ranging from  $5 \times 10^4$  bacteria/ml to  $5 \times 10^7$  bacteria/ml died within 2 days after being placed on these extreme dietary restriction diets. Dietary restriction (sDR) was considered  $5 \times 10^8$  bacteria/ml and ad libitum (AL) was  $5 \times 10^{11}$  bacteria/ml. OP50-1 grown to saturation overnight normally resulted in a concentration of bacteria of  $5 \times 10^{11}$  bacteria/ml. The lifespan assays were conducted in the absence of FUDR. For lifespan experiments with UV-killed bacteria, OP50-1 bacteria were UV-irradiated for 4 min in a UV Stratalinker 2400 (0.6 Joules). This type of UV treatment completely prevented bacterial growth in solid and liquid cultures.

#### AMP:ATP Ratio Quantification

100 day 7 hand-picked *C. elegans* were placed in ad libitum or sDR conditions for 12 hr under the same conditions as for lifespan assays. Worms were then picked onto bacteria-free plates, pooled, and washed twice with M9 buffer (KH<sub>2</sub>PO<sub>4</sub>, 22 mM; K<sub>2</sub>HPO<sub>4</sub>, 34 mM; NaCl, 86 mM; MgSO<sub>4</sub>, 1 mM). All but 20  $\mu$ l of M9 buffer was removed from worm pellet. 80  $\mu$ l of ice-cold 8% (v/v) perchloric acid was added to worm samples. The samples were then sonicated and 1N KHCO<sub>3</sub> was added to neutralize perchloric acid. The samples were briefly centrifuged and supernatants were filtered through

a .22  $\mu\text{m}$  filter, and then reverse-phase HPLC was performed on a LC-18 250  $\times$  4.6 mm 5  $\mu\text{m}$  column. Samples were separated by addition of buffer A ( $\text{KH}_2\text{PO}_4$  [pH 6.0], 0.1 M). Nucleotides were detected at 254 nm with a Varian Pro Star detector. Peak areas were measured with Varian Star Workstation software. Nucleotide identities were confirmed by comigration with AMP and ATP standards (Sigma).

#### AMPK In Vitro Kinase Assay

Purified AMPK (Upstate Biotechnology or purified from 293T cells) was incubated with various substrates (1  $\mu\text{g}$ ) in the kinase reaction buffer (HEPES [pH 7.0], 15 mM; DTT, 450  $\mu\text{M}$ ;  $\text{MgCl}_2$ , 18.75 mM;  $\beta$ -glycero-phosphate, 6.25 mM; EGTA, 1.25 mM; ATP, 125  $\mu\text{M}$ ) and 12.5  $\mu\text{Ci}$  radiolabeled ATP, with or without 150  $\mu\text{M}$  AMP, at 30°C for 15 min. Phosphorylation was detected by incorporation of radiolabeled [ $\gamma$ - $^{32}\text{P}$ ]ATP. Quantifications were performed with the Amersham Storm phospho-imager and the Image Quant 5.2 software. The stoichiometry of phosphorylation was measured by excising Coomassie-stained bands of substrate and by quantifying the amount of radioactivity incorporated with a scintillation counter. The molar incorporation of phosphate was calculated and was divided by the molar amount of substrate. To compare independent experiments, the average moles of phosphate/molar amount of substrate was used to normalize the data.

#### Liquid Chromatography Followed by Tandem Mass Spectrometry

A Coomassie-stained band corresponding to GST-DAF-16 phosphorylated by AMPK was excised from an SDS-PAGE gel, divided in half, reduced with DTT, alkylated with iodoacetamide, and digested with either trypsin or chymotrypsin. Peptide mixtures were separated by microcapillary reverse-phase chromatography and online analyzed in a hybrid linear ion trap-orbitrap (LTQ-Orbitrap, Thermo Electron) mass spectrometer. Mass spectra were database searched with the SEQUEST algorithm. All peptide matches were initially filtered based on enzyme specificity, mass measurement error, and Xcorr and  $\Delta\text{Corr}$  scores and further manually validated for peptide identification and site localization.

#### Worm RNA Extraction and Reverse Transcription Followed by Quantitative PCR

RNA was extracted by addition of 1 ml of Trizol (Invitrogen) for 100 ml of worm pellet, followed by six freeze-thaw cycles in liquid nitrogen. The RNA extraction was performed according to the Trizol protocol. The expression of *sod-3* was determined by reverse transcription of 1  $\mu\text{g}$  total RNA with the Superscript II kit (Invitrogen) followed by quantitative PCR analysis on a Bio-rad iCycler with iQ SYBR green (Bio-rad) with the following primers: *sod-3* F: CAACTTGGCTAAG GATGGTGGAG; *sod-3* R: GCATTGGCAAATCTCTCGCTG; *pan-actin* F: TCGGTATGGACAGAAGGAC; *pan-actin* R: CATCCCAGTTGGT GACGATA.

The experiments were conducted in triplicate and the results were expressed as  $2^{-(\text{Gene of interest number of cycles} - \text{actin number of cycles})}$ . Control PCR reactions were also performed on total RNA that had not been reverse-transcribed to test for the presence of genomic DNA in the RNA preparation.

Constructs, oxidative stress resistance in worms, quantification of bacterial ingestion with DsRed, DAF-16 localization assays, worm locomotor activity, and worm and human AMPK purification were performed as described in detail in the [Supplemental Experimental Procedures](#).

#### Supplemental Data

Four tables and Experimental Procedures are available at <http://www.current-biology.com/cgi/content/full/17/19/1646/DC1/>.

#### Acknowledgments

We thank A. Fire and M.-W. Tan for helpful discussion, for helping us set up all *C. elegans* assays, and for kindly providing reagents and strains. We thank J. Chen for his help with the AMP:ATP ratio measurement by HPLC. We thank K. Shen for helpful discussion and for his help in the cloning of the worm AMPK subunits. We thank L. Avery for helpful discussion regarding *eat-2* mutants. We thank Z.

Serber for helping us with stoichiometry calculations. We thank L. Attardi, G. Barsh, J. Lieberman, M. Greenberg, S. Paradis, T. Rando, J. Sage, M.-W. Tan, and the members of the Brunet laboratory for helpful discussion and for critically reading the manuscript. This research was supported by an NIH grant (NIA AG026648-01), an American Institute for Cancer Research grant, and a Pfizer/AFAR Innovations in Aging Research grant to A.B. E.L.G. was supported by a PHS grant (CA 09302) from the NCI and by an NSF graduate fellowship.

Received: July 11, 2007

Revised: August 15, 2007

Accepted: August 16, 2007

Published online: September 27, 2007

#### References

1. Masoro, E.J. (2005). Overview of caloric restriction and ageing. *Mech. Ageing Dev.* 126, 913–922.
2. Fontana, L., Meyer, T.E., Klein, S., and Holloszy, J.O. (2004). Long-term caloric restriction is highly effective in reducing the risk for atherosclerosis in humans. *Proc. Natl. Acad. Sci. USA* 101, 6659–6663.
3. Michels, K.B., and Ekblom, A. (2004). Caloric restriction and incidence of breast cancer. *JAMA* 291, 1226–1230.
4. Sinclair, D.A. (2005). Toward a unified theory of caloric restriction and longevity regulation. *Mech. Ageing Dev.* 126, 987–1002.
5. Kalant, N., Stewart, J., and Kaplan, R. (1988). Effect of diet restriction on glucose metabolism and insulin responsiveness in aging rats. *Mech. Ageing Dev.* 46, 89–104.
6. Kahn, B.B., Alquier, T., Carling, D., and Hardie, D.G. (2005). AMP-activated protein kinase: ancient energy gauge provides clues to modern understanding of metabolism. *Cell Metab.* 1, 15–25.
7. Pallottini, V., Montanari, L., Cavallini, G., Bergamini, E., Gori, Z., and Trentalance, A. (2004). Mechanisms underlying the impaired regulation of 3-hydroxy-3-methylglutaryl coenzyme A reductase in aged rat liver. *Mech. Ageing Dev.* 125, 633–639.
8. Cheung, P.C., Salt, I.P., Davies, S.P., Hardie, D.G., and Carling, D. (2000). Characterization of AMP-activated protein kinase gamma-subunit isoforms and their role in AMP binding. *Biochem. J.* 346, 659–669.
9. Shaw, R.J., Kosmatka, M., Bardeesy, N., Hurley, R.L., Witters, L.A., DePinho, R.A., and Cantley, L.C. (2004). The tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates apoptosis in response to energy stress. *Proc. Natl. Acad. Sci. USA* 101, 3329–3335.
10. Hawley, S.A., Pan, D.A., Mustard, K.J., Ross, L., Bain, J., Edelman, A.M., Frenguelli, B.G., and Hardie, D.G. (2005). Calmodulin-dependent protein kinase kinase-beta is an alternative upstream kinase for AMP-activated protein kinase. *Cell Metab.* 2, 9–19.
11. Woods, A., Dickerson, K., Heath, R., Hong, S.P., Momcilovic, M., Johnstone, S.R., Carlson, M., and Carling, D. (2005).  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase kinase-beta acts upstream of AMP-activated protein kinase in mammalian cells. *Cell Metab.* 2, 21–33.
12. Hurley, R.L., Anderson, K.A., Franzone, J.M., Kemp, B.E., Means, A.R., and Witters, L.A. (2005). The  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase kinases are AMP-activated protein kinase kinases. *J. Biol. Chem.* 280, 29060–29066.
13. Hong, S.P., Leiper, F.C., Woods, A., Carling, D., and Carlson, M. (2003). Activation of yeast Snf1 and mammalian AMP-activated protein kinase by upstream kinases. *Proc. Natl. Acad. Sci. USA* 100, 8839–8843.
14. Sanders, M.J., Grondin, P.O., Hegarty, B.D., Snowden, M.A., and Carling, D. (2007). Investigating the mechanism for AMP activation of the AMP-activated protein kinase cascade. *Biochem. J.* 403, 139–148.
15. Apfeld, J., O'Connor, G., McDonagh, T., DiStefano, P.S., and Curtis, R. (2004). The AMP-activated protein kinase AAK-2 links energy levels and insulin-like signals to lifespan in *C. elegans*. *Genes Dev.* 18, 3004–3009.

16. Curtis, R., O'Connor, G., and DiStefano, P.S. (2006). Aging networks in *Caenorhabditis elegans*: AMP-activated protein kinase (*aak-2*) links multiple aging and metabolism pathways. *Aging Cell* 5, 119–126.
17. Ogg, S., Paradis, S., Gottlieb, S., Patterson, G.I., Lee, L., Tissenbaum, H.A., and Ruvkun, G. (1997). The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in *C. elegans*. *Nature* 389, 994–999.
18. Lin, K., Dorman, J.B., Rodan, A., and Kenyon, C. (1997). *daf-16*: an HNF-3/forkhead family member that can function to double the life-span of *Caenorhabditis elegans*. *Science* 278, 1319–1322.
19. Lee, S.S., Kennedy, S., Tolonen, A.C., and Ruvkun, G. (2003). DAF-16 target genes that control *C. elegans* life-span and metabolism. *Science* 300, 644–647.
20. Murphy, C.T., McCarroll, S.A., Bargmann, C.I., Fraser, A., Kamath, R.S., Ahringer, J., Li, H., and Kenyon, C. (2003). Genes that act downstream of DAF-16 to influence the lifespan of *Caenorhabditis elegans*. *Nature* 424, 277–283.
21. Oh, S.W., Mukhopadhyay, A., Dixit, B.L., Raha, T., Green, M.R., and Tissenbaum, H.A. (2006). Identification of direct DAF-16 targets controlling longevity, metabolism and diapause by chromatin immunoprecipitation. *Nat. Genet.* 38, 251–257.
22. Brunet, A., Bonni, A., Zigmond, M.J., Lin, M.Z., Juo, P., Hu, L.S., Anderson, M.J., Arden, K.C., Blenis, J., and Greenberg, M.E. (1999). Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* 96, 857–868.
23. Kops, G.J., de Ruiter, N.D., De Vries-Smits, A.M., Powell, D.R., Bos, J.L., and Burgering, B.M. (1999). Direct control of the Forkhead transcription factor AFX by protein kinase B. *Nature* 398, 630–634.
24. Lee, G.D., Wilson, M.A., Zhu, M., Wolkow, C.A., de Cabo, R., Ingram, D.K., and Zou, S. (2006). Dietary deprivation extends lifespan in *Caenorhabditis elegans*. *Aging Cell* 5, 515–524.
25. Kaeberlein, T.L., Smith, E.D., Tsuchiya, M., Welton, K.L., Thomas, J.H., Fields, S., Kennedy, B.K., and Kaeberlein, M. (2006). Lifespan extension in *Caenorhabditis elegans* by complete removal of food. *Aging Cell* 5, 487–494.
26. Houthoofd, K., Braeckman, B.P., Johnson, T.E., and Vanfleteren, J.R. (2003). Life extension via dietary restriction is independent of the Ins/IGF-1 signalling pathway in *Caenorhabditis elegans*. *Exp. Gerontol.* 38, 947–954.
27. Bishop, N.A., and Guarente, L. (2007). Two neurons mediate diet-restriction-induced longevity in *C. elegans*. *Nature* 447, 545–549.
28. Panowski, S.H., Wolff, S., Aguilaniu, H., Durieux, J., and Dillin, A. (2007). PHA-4/Foxa mediates diet-restriction-induced longevity of *C. elegans*. *Nature* 447, 550–555.
29. Lakowski, B., and Hekimi, S. (1998). The genetics of caloric restriction in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* 95, 13091–13096.
30. Henderson, S.T., and Johnson, T.E. (2001). *daf-16* integrates developmental and environmental inputs to mediate aging in the nematode *Caenorhabditis elegans*. *Curr. Biol.* 11, 1975–1980.
31. Arad, M., Benson, D.W., Perez-Atayde, A.R., McKenna, W.J., Sparks, E.A., Kanter, R.J., McGarry, K., Seidman, J.G., and Seidman, C.E. (2002). Constitutively active AMP kinase mutations cause glycogen storage disease mimicking hypertrophic cardiomyopathy. *J. Clin. Invest.* 109, 357–362.
32. Honda, Y., and Honda, S. (1999). The *daf-2* gene network for longevity regulates oxidative stress resistance and Mn-superoxide dismutase gene expression in *Caenorhabditis elegans*. *FASEB J.* 13, 1385–1393.
33. Michell, B.J., Stapleton, D., Mitchellhill, K.I., House, C.M., Katsis, F., Witters, L.A., and Kemp, B.E. (1996). Isoform-specific purification and substrate specificity of the 5'-AMP-activated protein kinase. *J. Biol. Chem.* 271, 28445–28450.
34. Greer, E.L., Oskoui, P.R., Banko, M.R., Maniar, J.M., Gygi, M.P., Gygi, S.P., and Brunet, A. (2007). The energy sensor AMP-activated protein kinase directly regulates the mammalian FOXO3 transcription factor. *J. Biol. Chem.* Published online August 20, 2007. 10.1074/jbc.M705325200.
35. Klass, M.R. (1977). Aging in the nematode *Caenorhabditis elegans*: major biological and environmental factors influencing life span. *Mech. Ageing Dev.* 6, 413–429.
36. Szewczyk, N.J., Udranszky, I.A., Kozak, E., Sunga, J., Kim, S.K., Jacobson, L.A., and Conley, C.A. (2006). Delayed development and lifespan extension as features of metabolic lifestyle alteration in *C. elegans* under dietary restriction. *J. Exp. Biol.* 209, 4129–4139.
37. Hosono, R., Nishimoto, S., and Kuno, S. (1989). Alterations of life span in the nematode *Caenorhabditis elegans* under monoxenic culture conditions. *Exp. Gerontol.* 24, 251–264.
38. Avery, L. (1993). The genetics of feeding in *Caenorhabditis elegans*. *Genetics* 133, 897–917.
39. Lin, S.J., Kaeberlein, M., Andalis, A.A., Sturtz, L.A., Defossez, P.A., Culotta, V.C., Fink, G.R., and Guarente, L. (2002). Calorie restriction extends *Saccharomyces cerevisiae* lifespan by increasing respiration. *Nature* 418, 344–348.
40. Rogina, B., and Helfand, S.L. (2004). Sir2 mediates longevity in the fly through a pathway related to calorie restriction. *Proc. Natl. Acad. Sci. USA* 101, 15998–16003.
41. Wood, J.G., Rogina, B., Lavu, S., Howitz, K., Helfand, S.L., Tatar, M., and Sinclair, D. (2004). Sirtuin activators mimic caloric restriction and delay ageing in metazoans. *Nature* 430, 686–689.
42. Wang, Y., and Tissenbaum, H.A. (2006). Overlapping and distinct functions for a *Caenorhabditis elegans* SIR2 and DAF-16/FOXO. *Mech. Ageing Dev.* 127, 48–56.
43. Kaeberlein, M., Kirkland, K.T., Fields, S., and Kennedy, B.K. (2004). Sir2-independent life span extension by calorie restriction in yeast. *PLoS Biol.* 2, E296. 10.1371/journal.pbio.0020296.
44. Hansen, M., Taubert, S., Crawford, D., Libina, N., Lee, S.J., and Kenyon, C. (2007). Lifespan extension by conditions that inhibit translation in *Caenorhabditis elegans*. *Aging Cell* 6, 95–110.
45. Inoki, K., Zhu, T., and Guan, K.L. (2003). TSC2 mediates cellular energy response to control cell growth and survival. *Cell* 115, 577–590.
46. Kaeberlein, M., Powers, R.W., 3rd, Steffen, K.K., Westman, E.A., Hu, D., Dang, N., Kerr, E.O., Kirkland, K.T., Fields, S., and Kennedy, B.K. (2005). Regulation of yeast replicative life span by TOR and Sch9 in response to nutrients. *Science* 310, 1193–1196.
47. Vellai, T., Takacs-Vellai, K., Zhang, Y., Kovacs, A.L., Orosz, L., and Muller, F. (2003). Genetics: influence of TOR kinase on lifespan in *C. elegans*. *Nature* 426, 620.
48. Kapahi, P., Zid, B.M., Harper, T., Koslover, D., Sapin, V., and Benzer, S. (2004). Regulation of lifespan in *Drosophila* by modulation of genes in the TOR signaling pathway. *Curr. Biol.* 14, 885–890.
49. Pan, K.Z., Palter, J.E., Rogers, A.N., Olsen, A., Chen, D., Lithgow, G.J., and Kapahi, P. (2007). Inhibition of mRNA translation extends lifespan in *Caenorhabditis elegans*. *Aging Cell* 6, 111–119.
50. Baur, J.A., Pearson, K.J., Price, N.L., Jamieson, H.A., Lerin, C., Kalra, A., Prabhu, V.V., Allard, J.S., Lopez-Lluch, G., Lewis, K., et al. (2006). Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* 444, 337–342.
51. Anisimov, V.N., Berstein, L.M., Egormin, P.A., Piskunova, T.S., Popovich, I.G., Zabezhinski, M.A., Kovalenko, I.G., Poroshina, T.E., Semenchenko, A.V., Provinciali, M., et al. (2005). Effect of metformin on life span and on the development of spontaneous mammary tumors in HER-2/neu transgenic mice. *Exp. Gerontol.* 40, 685–693.
52. Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics* 77, 71–94.
53. Henderson, S.T., Bonafe, M., and Johnson, T.E. (2006). *daf-16* protects the nematode *Caenorhabditis elegans* during food deprivation. *J. Gerontol. A Biol. Sci. Med. Sci.* 61, 444–460.