

Emerging Policy Challenges for Integrated Assessment

Snowmass

Allen Fawcett

July 22, 2019

Paris Agreement

- Article 2

1. This Agreement, in enhancing the implementation of the Convention, including its objective, aims to strengthen the global response to the threat of climate change, in the context of sustainable development and efforts to eradicate poverty, including by:
 - a) Holding the increase in the global average temperature to well below 2 °C above pre-industrial levels and to pursue efforts to limit the temperature increase to 1.5 °C above pre-industrial levels, recognizing that this would significantly reduce the risks and impacts of climate change;

- Article 4

1. In order to achieve the long-term temperature goal set out in Article 2, Parties aim to reach global peaking of greenhouse gas emissions as soon as possible, recognizing that peaking will take longer for developing country Parties, and to undertake rapid reductions thereafter in accordance with best available science, so as to achieve a balance between anthropogenic emissions by sources and removals by sinks of greenhouse gases in the second half of this century, on the basis of equity, and in the context of sustainable development and efforts to eradicate poverty.
19. All Parties should strive to formulate and communicate long-term low greenhouse gas emission development strategies, mindful of Article 2 taking into account their common but differentiated responsibilities and respective capabilities, in the light of different national circumstances.


- Decision Text

21. Invites the Intergovernmental Panel on Climate Change to provide a special report in 2018 on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways;

Global Warming of 1.5°C

An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty

Summary for Policymakers



“We have 12 years to limit climate change catastrophe, warns UN”

<https://www.theguardian.com/environment/2018/oct/08/global-warming-must-not-exceed-15c-warns-landmark-un-report>

“Report: we have just 12 years to limit devastating global warming”

<https://www.vox.com/2018/10/8/17948832/climate-change-global-warming-un-ipcc-report>

“We only have 12 years to save the planet”

https://www.washingtonpost.com/opinions/we-only-have-12-years-to-stop-climate-change/2019/01/14/42704374-15d1-11e9-ab79-30cd4f7926f2_story.html?utm_term=.e0e23a6f9101


A. Understanding Global Warming of 1.5°C⁴

- A.1 Human activities are estimated to have caused approximately 1.0°C of global warming⁵ above pre-industrial levels, with a *likely* range of 0.8°C to 1.2°C. Global warming is *likely* to reach 1.5°C between 2030 and 2052 if it continues to increase at the current rate. (*high confidence*) (Figure SPM.1) {1.2}


Cumulative emissions of CO₂ and future non-CO₂ radiative forcing determine the probability of limiting warming to 1.5°C

a) Observed global temperature change and modeled responses to stylized anthropogenic emission and forcing pathways

Global warming relative to 1850-1900 (°C)




b) Stylized net global CO₂ emission pathways Billion tonnes CO₂ per year (GtCO₂/yr)




Faster immediate CO₂ emission reductions limit cumulative CO₂ emissions shown in panel (c).

c) Cumulative net CO₂ emissions Billion tonnes CO₂ (GtCO₂)



Maximum temperature rise is determined by cumulative net CO₂ emissions and net non-CO₂ radiative forcing due to methane, nitrous oxide, aerosols and other anthropogenic forcing agents.

d) Non-CO₂ radiative forcing pathways Watts per square metre (W/m²)




Global emissions pathway characteristics

General characteristics of the evolution of anthropogenic net emissions of CO₂, and total emissions of methane, black carbon, and nitrous oxide in model pathways that limit global warming to 1.5°C with no or limited overshoot. Net emissions are defined as anthropogenic emissions reduced by anthropogenic removals. Reductions in net emissions can be achieved through different portfolios of mitigation measures illustrated in Figure SPM.3b.


Global total net CO₂ emissions

Billion tonnes of CO₂/yr



Timing of net zero CO₂


Line widths depict the 5-95th percentile and the 25-75th percentile of scenarios




Non-CO₂ emissions relative to 2010

Emissions of non-CO₂ forcers are also reduced or limited in pathways limiting global warming to 1.5°C with **no or limited overshoot**, but they do not reach zero globally.


Methane emissions



Black carbon emissions



Nitrous oxide emissions




Characteristics of four illustrative model pathways

Different mitigation strategies can achieve the net emissions reductions that would be required to follow a pathway that limits global warming to 1.5°C with no or limited overshoot. All pathways use Carbon Dioxide Removal (CDR), but the amount varies across pathways, as do the relative contributions of Bioenergy with Carbon Capture and Storage (BECCS) and removals in the Agriculture, Forestry and Other Land Use (AFOLU) sector. This has implications for emissions and several other pathway characteristics.

Breakdown of contributions to global net CO₂ emissions in four illustrative model pathways

● Fossil fuel and industry ● AFOLU ● BECCS



P1: A scenario in which social, business and technological innovations result in lower energy demand up to 2050 while living standards rise, especially in the global South. A downsized energy system enables rapid decarbonization of energy supply. Afforestation is the only CDR option considered; neither fossil fuels with CCS nor BECCS are used.

P2: A scenario with a broad focus on sustainability including energy intensity, human development, economic convergence and international cooperation, as well as shifts towards sustainable and healthy consumption patterns, low-carbon technology innovation, and well-managed land systems with limited societal acceptability for BECCS.

P3: A middle-of-the-road scenario in which societal as well as technological development follows historical patterns. Emissions reductions are mainly achieved by changing the way in which energy and products are produced, and to a lesser degree by reductions in demand.

P4: A resource- and energy-intensive scenario in which economic growth and globalization lead to widespread adoption of greenhouse-gas-intensive lifestyles, including high demand for transportation fuels and livestock products. Emissions reductions are mainly achieved through technological means, making strong use of CDR through the deployment of BECCS.


Global indicators	P1	P2	P3	P4	Interquartile range
<i>Pathway classification</i>	No or limited overshoot	No or limited overshoot	No or limited overshoot	Higher overshoot	No or limited overshoot
<i>CO₂ emission change in 2030 (% rel to 2010)</i>	-58	-47	-41	4	(-58,-40)
↳ <i>in 2050 (% rel to 2010)</i>	-93	-95	-91	-97	(-107,-94)
<i>Kyoto-GHG emissions* in 2030 (% rel to 2010)</i>	-50	-49	-35	-2	(-51,-39)
↳ <i>in 2050 (% rel to 2010)</i>	-82	-89	-78	-80	(-93,-81)

NOTE: Indicators have been selected to show global trends identified by the Chapter 2 assessment. National and sectoral characteristics can differ substantially from the global trends shown above.

* Kyoto-gas emissions are based on IPCC Second Assessment Report GWP-100
 ** Changes in energy demand are associated with improvements in energy efficiency and behaviour change


A.2.2 Reaching and sustaining net zero global anthropogenic CO₂ emissions and declining net non-CO₂ radiative forcing would halt anthropogenic global warming on multi-decadal time scales (*high confidence*). The maximum temperature reached is then determined by cumulative net global anthropogenic CO₂ emissions up to the time of net zero CO₂ emissions (*high confidence*) and the level of non-CO₂ radiative forcing in the decades prior to the time that maximum temperatures are reached (*medium confidence*). On longer time scales, sustained net negative global anthropogenic CO₂ emissions and/or further reductions in non-CO₂ radiative forcing may still be required to prevent further warming due to Earth system feedbacks and to reverse ocean acidification (*medium confidence*) and will be required to minimize sea level rise (*high confidence*). {Cross-Chapter Box 2 in Chapter 1, 1.2.3, 1.2.4, Figure 1.4, 2.2.1, 2.2.2, 3.4.4.8, 3.4.5.1, 3.6.3.2}

GLOBAL EMISSIONS TRAJECTORIES TO NET-ZERO GHG EMISSIONS AND PROBABILITY OF GLOBAL TEMPERATURE CHANGES



The United States MCS puts the nation on a path consistent with a successful global outcome. Achieving the Paris Agreement temperature goals will require increasing global ambition leading to 2030 and steep reductions to net-zero global GHG emissions following 2030. We show the probability of staying below 2°C and 1.5°C across global scenarios by 2100. While there could be an overshoot of the Paris Agreement temperature objectives before 2100, achieving net-zero GHG emissions globally could bring temperatures below peak levels in 2100 and beyond.

U.S. NET GHG EMISSIONS UNDER THREE MCS SCENARIOS



Multiple pathways to 80 percent GHG reductions by 2050 are achievable through large reductions in energy CO₂ emissions, smaller reductions in non-CO₂ emissions, and delivering negative emissions from land and CO₂ removal technologies. Note: "No CO₂ removal tech" assumes no availability of negative emissions technologies like BECCS.

-  Net GHG
-  CO₂
-  Non-CO₂
-  Land Sink
-  CO₂ Removal Technologies

US Context

