GEOMETRY TEST 2007 STANFORD MATH TOURNAMENT MARCH 4, 2007

- 1. An equilateral triangle has perimeter numerically equal to its area, which is not zero. Find its side length.
- 2. Two spheres of radius 2 pass through each other's center. Find the surface area of the regular octahedron inscribed within the space enclosed by both spheres.
- 3. Cumulation of a polyhedron means replacing each face with a pyramid of height h using the face as a base. There is a cumulation of the cube of side length s which (after removing unecessary edges) has twelve sides, each a congruent rhombus. What is the height h used in this cumulation?
- 4. Nathan is standing on vertex A of triangle ABC, with AB=3, BC=5, and CA=4. Nathan walks according to the following plan: He moves along the altitude-to-the-hypotenuse until he reaches the hypotenuse. He has now cut the original triangle into two triangles; he now walks along the altitude to the hypotenuse of the larger one. He repeats this process forever. What is the total distance that Nathan walks?
- 5. Given an octahedron with every edge of length s, what is the radius of the largest sphere that will fit in this octahedron?
- 6. Let TINA be a quadrilateral with IA = 8, IN = 4, $m \angle T = 30^{\circ}$, $m \angle NAT = 60^{\circ}$, and $m \angle TIA = m \angle INA$. Find NA.
- 7. Two regular tetrahedra of side length 2 are positioned such that the midpoint of each side of one coincides with the midpoint of a side of the other, and the tetrahedra themselves do not coincide. Find the volume of the region in which they overlap.
- 8. $\triangle ABC$ has $\overline{AB} = \overline{AC}$. Points M and N are midpoints of \overline{AB} and \overline{AC} , respectively. The medians \overline{MC} and \overline{NB} intersect at a right angle. Find $(\frac{\overline{AB}}{\overline{BC}})^2$.
- 9. Points P, Q, R, S, T lie in the pane with S on \overline{PR} and R on \overline{QT} . If PQ = 5, PS = 3, PR = 5, QS = 3, and $RT = 4/\sqrt{3}$, what is ST?
- 10. A car starts moving at constant speed at the origin facing in the positive y-direction. Its minimum turning radius is such that it the soonest it can return to the x-axis is after driving a distance d. Let Γ be the boundary of the region the car can reach by driving at most a distance d; find an x > 0 so that $\left(x, \frac{d}{3} + \frac{d\sqrt{3}}{2\pi}\right)$ is on Γ .