



The MAJORANA Demonstrator Neutrinoless Double-Beta Decay Experiment

Brandon White
Oak Ridge National Laboratory



On Behalf of The MAJORANA
Demonstrator Experiment



Outline

- Neutrinoless $\beta\beta$ decay
- MAJORANA Demonstrator
- Experiment Status



Outline


- Neutrinoless $\beta\beta$ decay
- MAJORANA Demonstrator Design
- Experiment Status




Neutrinoless $\beta\beta$ decay

This rare decay in which a nucleus (Z, A) decays to a new nucleus $(Z + 2, A)$.

2ν case.



0ν case.




Neutrinoless $\beta\beta$ decay

This rare decay in which a nucleus (Z, A) decays to a new nucleus $(Z + 2, A)$.


2ν case.




Measured In
several Isotopes



0ν case.




Only limits for
half-life of decay



Conditions for the zero neutrino case to occur include lepton number violation, massive neutrinos, and the neutrino must be its own antiparticle.

Neutrinoless $\beta\beta$ decay

$$(T_{1/2}^{0\nu})^{-1} = G_{0\nu}(Q_{\beta\beta}, Z) |M^{0\nu}|^2 \langle m_\nu \rangle^2$$



Neutrinoless $\beta\beta$ decay

Isotope	$2\nu \beta\beta$ decay half-life [yr]
^{48}Ca	$\sim 4.3 \times 10^{19}$
^{76}Ge	$\sim 1.5 \times 10^{21}$
^{82}Se	$\sim 9.2 \times 10^{19}$
^{96}Zr	$\sim 2.0 \times 10^{19}$
^{100}Mo	$\sim 7.0 \times 10^{18}$
^{116}Cd	$\sim 3.0 \times 10^{19}$
^{130}Te	$\sim 9.1 \times 10^{20}$
^{136}Xe	$\sim 2.1 \times 10^{21}$
^{150}Nd	$\sim 7.8 \times 10^{18}$


Some Considerations for Isotope Selection:

- Q value of the decay
- Nuclear matrix element
- Availability and abundance of isotope
- Detector design

Detection Requirements

Basic requirements for detecting such rare decay:

- Large source mass of desired isotope
- Low background to reach acceptable signal to background ratios
- High resolution to separate 0ν detection from tail of 2ν energy spectrum



Neutrinoless $\beta\beta$ decay

^{136}Xe



^{116}Cd



^{82}Se



^{130}Te




IGEX


^{76}Ge



Sensitivity, Background and Exposure



Discovery, Background and Exposure



Outline

- Neutrinoless $\beta\beta$ decay
- **MAJORANA Demonstrator**
- Experiment Status





The MAJORANA Collaboration



Black Hills State University, Spearfish, SD
Kara Keeter

Duke University, Durham, North Carolina, and TUNL
Matthew Busch

Institute for Theoretical and Experimental Physics, Moscow, Russia
Alexander Barabash, Sergey Konovalov, Vladimir Yumatov

Joint Institute for Nuclear Research, Dubna, Russia
Viktor Brudanin, M. Shirchenko, Sergey Vasilyev, E. Yakushev,
I. Zhitnikov

*Lawrence Berkeley National Laboratory, Berkeley, California and
the University of California - Berkeley*
Nicolas Abgrall, Mark Amman, Paul Barton, Adam Bradley,
Yuen-Dat Chan, Paul Luke, Susanne Mertens, Alan Poon,
Christopher Schmitt, Kai Vetter, Harold Yaver

Los Alamos National Laboratory, Los Alamos, New Mexico
Pinghan Chu, Steven Elliott, Johnny Goett, Ralph Massarczyk,
Keith Rielage, Larry Rodriguez, Harry Salazar, Wenqin Xu

Oak Ridge National Laboratory
Cristian Baldenegro-Barrera, Fred Bertrand, Kathy Carney,
Alfredo Galindo-Uribarri, Matthew P. Green, Monty Middlebrook,
David Radford, **Elisa Romero-Romero**, Robert Varner, Brandon White,
Timothy Williams, Chang-Hong Yu

Osaka University, Osaka, Japan
Hiroyasu Ejiri

Pacific Northwest National Laboratory, Richland, Washington
Isaac Arnquist, Eric Hoppe, Richard T. Kouzes, Brian LaFerriere, John Orrell

South Dakota School of Mines and Technology, Rapid City, South Dakota
Adam Caldwell, Cabot-Ann Christofferson, Stanley Howard,
Anne-Marie Suriano, Jared Thompson

Tennessee Tech University, Cookeville, Tennessee
Mary Kidd

University of North Carolina, Chapel Hill, North Carolina and TUNL
Tom Gilliss, **Graham K. Giovanetti**, Reyco Henning, **Jacqueline MacMullin**,
Samuel J. Meijer, **Benjamin Shanks**, Christopher O' Shaughnessy, **Jamin Rager**,
James Trimble, **Kris Vorren**, John F. Wilkerson

University of South Carolina, Columbia, South Carolina
Frank Avignone, Vince Guiseppe, David Tedeschi, **Clint Wiseman**

University of South Dakota, Vermillion, South Dakota
Dana Byram, **Ben Jasinski**, Ryan Martin, **Nathan Snyder**

University of Tennessee, Knoxville, Tennessee
Yuri Efremenko

University of Washington, Seattle, Washington
Tom Burritt, **Micah Buuck**, Clara Cuesta, Jason Detwiler, **Julieta Gruszko**,
Ian Guinn, Greg Harper, **Jonathan Leon**, David Peterson, R. G. Hamish Robertson,
Tim Van Wechel



The MAJORANA DEMONSTRATOR

Funded by DOE Office of Nuclear Physics and NSF Particle Astrophysics,
with additional contributions from international collaborators.



- Goals:
- Demonstrate backgrounds low enough to justify building a tonne scale experiment.
 - Establish feasibility to construct & field modular arrays of Ge detectors.
 - Searches for additional physics beyond the standard model.


- Located underground at 4850' Sanford Underground Research Facility
- Background Goal in the $0\nu\beta\beta$ peak region of interest (4 keV at 2039 keV)
3 counts/ROI/t/y (after analysis cuts) Assay U.L. currently ≤ 3.5
scales to 1 count/ROI/t/y for a tonne experiment
- 44-kg of Ge detectors
 - 29.6 kg of 87% enriched ^{76}Ge crystals
 - 15 kg of $^{\text{nat}}\text{Ge}$
 - Detector Technology: P-type, point-contact.
- 2 independent cryostats
 - ultra-clean, electroformed Cu
 - 20 kg of detectors per cryostat
 - naturally scalable
- Compact Shield
 - low-background passive Cu and Pb shield with active muon veto




Underground Laboratory




Located at the former Homestake Mine in Lead, SD.



Underground Laboratory



Apparatus Overview



The Shield



The Shield




Note keyed structure of shield



- Inner shield is made of underground electroformed copper surrounded by clean commercial copper.
- Cleaned Lead structure surrounds the inner copper shield.

Blank Monolith, when running only one cryostat of detectors




Shield Details




Side Veto

Upper Veto



Panel of Rn exclusion box



Prototype Module Cryostat


Keyed Pb Stack

Modules




A Module is:


- Cryostat
- Thermosyphon
- Vacuum
- Shield Section
- All resting on a movable bearing table



Modules





Top of the Cold Plate



Vespel connectors

HV cables are run from vacuum feed-through to detector.

Assembled Detector Unit and String



String Assembly


Electroformed Parts




All detector unit, string, and cryostat components are made from underground electroformed copper.



Front-End Board




Clean Au+Ti traces on fused silica, amorphous Ge resistor, FET mounted with silver epoxy, EFCu + low-BG Sn contact pin



Signal Connectors





Connectors reside on top of cold plate.
In-house machined from vespel. Axon' pico co-ax cable.
Low background solder and flux.





Detectors

P-type point contact (PPC) Ge detectors are used to maximize crystal size and allow for discrimination between multi-site and single-site interactions. This technique allows for the reduction of gamma backgrounds.





Hole v_{drift} (mm/ns), isocrones (grey) and paths (black)

Materials and Assay



- Significant R&D and advances made in improvement of ICP-MS sensitivity for U and Th in copper near 0.1 mBq/kg level.
- Monitoring U and Th in copper baths electrolyte.
- All plastic materials selected after high sensitivity NAA analysis. Assay complete.
- Significant progress made in development of low background front-end electronics.




DEMONSTRATOR Background Model



GEANT4 simulations to determine efficiencies for contamination to deposit energy in our detectors


50k CPU hours

8k+ runs, 40+ contaminants, 56 components, 21 materials



A. Schubert,
Univ. Washington,
PhD Dec. 2012

DEMONSTRATOR Background Model



Outline


- Neutrinoless $\beta\beta$ decay
- MAJORANA Demonstrator
- **Experiment Status**






Detectors

- Germanium was enriched in Russia and shipped by sea to Oak Ridge.
- Germanium was stored underground when not in detector production.





Cherokee Caverns outside of Oak Ridge



Detectors




- **ORTEC** selected for enriched detectors from early 2013 through June 2015.
- 35 Enriched detectors at SURF 29.7 kg, 87% ^{76}Ge
- 20 kg of modified natural-Ge BEGe (Canberra) detectors in hand (33 detectors UG).



- All detector related assembly performed in N_2 purged gloveboxes.
- All detectors' dimensions recorded by optical reader.


Detector Resolution



Detector Resolution

At 1332 keV, FWHM = 1.98 keV +/- 0.024 keV

At 2614 keV, FWHM = 2.72 keV +/- 0.15 keV




Pulse Shape Discrimination




Natural BEGe detector in Prototype Cryostat

B8466



Rising Edge for single and multi-site events



Electroforming



Temporary Clean Room at Ross Shops



Installation of mandrel in bath

Underground Electroformed copper growing since 2011.
Electroforming completed May 2015



NOV 20 15



Brandon White



MJD Status

Underground Clean Machine Shop operating since early 2012



Clean Lab Occupancy Since Summer 2012



The Shield



- Pb shield constructed
- Outer Cu shield layer installed
- Rn exclusion box installed
- Poly layers being installed
- Hovair in-use underground
- Most veto panels operational
- Calibration system demonstrated




MAJORANA DEMONSTRATOR Implementation

Three Steps


Prototype cryostat: 7.0 kg (10) ^{nat}Ge

Same design as Modules 1 and 2, but fabricated using OFHC Cu Components




June 2014

**Module 1: 16.8 kg (20) ^{enr}Ge
5.7 kg (9) ^{nat}Ge**





May 2015

**Module 2: 12.6 kg (14) ^{enr}Ge
9.4 kg (15) ^{nat}Ge**



End 2015



Modules



- Prototype cryostat operating, over 140 d live.
- Prototype was populated with two full strings and one single detector string of natural Ge detectors.
- Commercial copper components were used.

Modules




Module 1 currently in commissioning phase.



Module 2 construction and commissioning by the end of 2015

Sensitivity, Background and Exposure



Acknowledgements

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics. We acknowledge support from the Particle Astrophysics Program of the National Science Foundation. This research uses these US DOE Office of Science User Facilities: the National Energy Research Scientific Computing Center and the Oak Ridge Leadership Computing Facility. We acknowledge support from the Russian Foundation for Basic Research. We thank our hosts and colleagues at the Sanford Underground Research Facility for their support.



Thank you to the organizers of
the conference



Simulation: MJD 0-10 MeV




MJD, 5-year exposure




Backup

Scaling from MJD projects




Backup

Simulated spectra, 60 kg yrs, detector resolution applied




Backup

Simulated spectra, 60 kg yrs, detector resolution + all cuts applied




Backup



Backup

A. Dueck, W. Rodejohann and K. Zuber, Phys. Rev. D83 (2011)



Backup

H.V. Klapdor-Kleingrothaus And I. V. Krivosheina, Mod. Phys. LeN. A 21, 1547 (2006)

