

**LECTURE NOTES ON GEOMETRIC FEATURES OF THE  
ALLEN–CAHN EQUATION (UNIVERSITY OF CONNECTICUT,  
2018)**

OTIS CHODOSH

These are my notes from four lectures at a University of Connecticut summer school on geometric analysis in 2018. A “+” marks the exercises which are used subsequently in the text. The others can safely be skipped without subsequent confusion, but are probably more interesting/difficult. I am very grateful to be informed of any inaccuracies, typos, incorrect references, or other issues.

1. INTRODUCTION TO THE ALLEN–CAHN EQUATION

We will consider throughout  $(M^n, g)$  a complete Riemannian manifold.

**Definition 1.** We define the Allen–Cahn energy by

$$E_\varepsilon(u; \Omega) := \int_\Omega \left( \frac{\varepsilon}{2} |\nabla_g u|^2 + \frac{1}{\varepsilon} W(u) \right) d\mu_g.$$

Here  $W(\cdot)$  is a “double well potential,” which we will take as  $W(t) = \frac{1}{4}(1 - t^2)^2$  (more general functions are also possible). We will often drop  $\Omega$  (e.g. when  $M$  is compact).

It is clear that  $E_\varepsilon$  is well defined for  $u \in H^1(\Omega) \cap L^4(\Omega)$ . It is convenient to extend  $E_\varepsilon$  to functions  $u \notin H^1(\Omega) \cap L^4(\Omega)$  by  $E_\varepsilon(u) = \infty$ .




FIGURE 1. The double well potential  $W(t) = \frac{1}{4}(1 - t^2)^2$ .

**1.1. Brief history/heuristics.** Energy functionals of the form described above were first considered by Van der Waals (see the translated article [vdW79]) in 1893 and then rediscovered by Cahn–Hilliard [CH58] in 1958. They considered  $u(x)$  as describing what fraction of two phases/densities a system is at  $x$ . They argued that the “free energy” of such a system can be approximated by a functional of the form

$$\int \kappa |\nabla u|^2 + f(u)$$

for  $\kappa > 0$  and  $f(u)$  some function that represents the local energy density of a system that is entirely in the state  $u(x)$ . The “double well” we have considered above corresponds to the assumption that the minimum local energy occurs precisely when  $u(x) \in \{\pm 1\}$ . The gradient term (and scale factor  $\kappa$ ) influences how the two phases interact (and can be seen as a first-order correction to the energy that is formed simply by the total energy density of the system). Allen–Cahn [AC79] observed in 1978 that there is a basic link between the location of the interface between the two phases and the mean curvature of the interface. This link serves as the other main basis (besides the physical motivation) for mathematical interest in such energy functionals.

## 1.2. Critical points and the Allen–Cahn equation.

**Definition 2.** A function  $u : M \rightarrow \mathbb{R}$  is a *critical point* of  $E_\varepsilon$  if for any  $\varphi : M \rightarrow \mathbb{R}$  smooth with support compactly contained in a precompact open set  $\Omega \subset M$ , we have  $u \in H^1(\Omega) \cap L^4(\Omega)$  and

$$\left. \frac{d}{dt} \right|_{t=0} E_\varepsilon(u + t\varphi; \Omega) = 0.$$

**Exercise 1 (+).** Show that a function  $u : M \rightarrow \mathbb{R}$  is a critical point of  $E_\varepsilon$  if and only if  $u$  (weakly) solves the Allen–Cahn equation

$$\varepsilon \Delta_g u = \frac{1}{\varepsilon} W'(u) = \frac{1}{\varepsilon} u(u^2 - 1).$$

**Exercise 2.** This exercise requires some knowledge of elliptic regularity. It can be safely skipped.

- (a) Prove that if  $u : M \rightarrow \mathbb{R}$  is a critical point of the Allen–Cahn functional  $u$  with the additional property that  $u \in L^\infty(\Omega)$  for all precompact open sets  $\Omega$ , then  $u$  is smooth.
- (b) For a smooth critical point of the Allen–Cahn functional  $u$  on a closed manifold  $(M, g)$ , show that  $u \in [-1, 1]$ .

Note that (a) holds without the extra assumption that  $u \in L^\infty(\Omega)$ , by a somewhat more involved application of elliptic theory.

**Exercise 3 (+).** Check that  $u \equiv \pm 1$  and  $u = 0$  are all critical points for the Allen–Cahn equation. If  $(M, g)$  is compact, show that  $u = \pm 1$  are the unique global minimizers for  $E_\varepsilon$  in the sense that for any  $v : M \rightarrow \mathbb{R}$ , we have

$$E_\varepsilon(v) \geq E_\varepsilon(\pm 1) = 0,$$

with equality only for  $v \equiv \pm 1$ .

Are there other solutions to the Allen–Cahn equation?

**1.3. One dimensional solution.** Let us begin by considering the Allen–Cahn equation on  $\mathbb{R}$ . The Allen–Cahn equation becomes

$$(1) \quad \varepsilon u''(t) = \frac{1}{\varepsilon} W'(u(t)).$$

Rescaling by  $\varepsilon$  allows us (in this case) to study only  $\varepsilon = 1$ . Set  $\tilde{u}(t) = u(\varepsilon t)$ , so

$$\tilde{u}''(t) = \varepsilon^2 u''(\varepsilon t) = W'(u(\varepsilon t)) = W'(\tilde{u}(t)).$$

Thus, we will begin by considering  $\varepsilon = 1$  (and then rescale the coordinate function  $t$  to return to arbitrary  $\varepsilon$ ).

Dropping the tilde, we seek (other than  $u \equiv \pm 1$ ) solutions to

$$(2) \quad u''(t) = W'(u(t)) = u(t)^3 - u(t)$$

with finite energy on all of  $\mathbb{R}$ , i.e.,

$$\int_{-\infty}^{\infty} (u'(t)^2 + W(u(t))) dt < \infty.$$

**Exercise 4 (+).** (a) Show that the unique solutions to (2) with finite energy are given by  $u(t) \equiv \pm 1$  and  $u(t) = \pm \mathbb{H}(t - t_0)$ , where  $\mathbb{H}(t) = \tanh(t/\sqrt{2})$ . Hint: consider the quantity  $u'(t)^2 - 2W(u(t))$ .

(b) What are the solutions to (2) under the assumption  $u'(t) > 0$  (but not a priori assuming finite energy).




FIGURE 2. The heteroclinic solution  $\mathbb{H}(t) = \tanh(t/\sqrt{2})$ .

Thus, rescaling back to the general  $\varepsilon > 0$  equation we've seen

$$\mathbb{H}_\varepsilon(t) := \tanh\left(\frac{t}{\varepsilon\sqrt{2}}\right)$$

is the unique (up to sign and translation) non-trivial solution with finite energy to

$$\varepsilon\mathbb{H}_\varepsilon''(t) = \frac{1}{\varepsilon}W'(\mathbb{H}_\varepsilon(t)).$$

1.3.1. *First glimpse of the  $\varepsilon \rightarrow 0$  limit.* Note that for  $t > 0$ ,

$$\lim_{\varepsilon \rightarrow 0} \mathbb{H}_\varepsilon(t) = 1$$

and for  $t < 0$

$$\lim_{\varepsilon \rightarrow 0} \mathbb{H}_\varepsilon(t) = -1$$




FIGURE 3. The heteroclinic solution  $\mathbb{H}_\varepsilon(t)$  with  $\varepsilon = .01$  is converging to a step function.

Thus,  $\mathbb{H}_\varepsilon$  converges a.e., to the step function

$$\mathbb{H}_0(t) := \begin{cases} +1 & t > 0 \\ -1 & t < 0 \end{cases}.$$

Note that

$$\{0\} = \partial\{\mathbb{H}_0(t) = 1\}.$$

This somewhat trivial observation is the first hint of the connection between the singular limit  $\varepsilon \rightarrow 0$  for solutions to the Allen–Cahn equation and hypersurfaces (in this case, just a point).

1.4. **Solutions on  $\mathbb{R}^2$ .** We have seen that the set of (finite energy) solutions to the Allen–Cahn ODE on  $\mathbb{R}$  is rather simple (although the solution  $\mathbb{H}(t)$  is very important). We turn to solutions on  $\mathbb{R}^2$ . Observe that as above, if

$$(3) \quad \Delta u = W'(u),$$

then  $u_\varepsilon(x) := u(x/\varepsilon)$  solves

$$\varepsilon\Delta u_\varepsilon = \frac{1}{\varepsilon}W'(u_\varepsilon)$$

1.4.1. *The one-dimensional solution on  $\mathbb{R}^2$ .* We first observe that the one dimensional solution  $\mathbb{H}(t)$  we considered before provides a solution on  $\mathbb{R}^2$  as well. To that end, fix  $a \in \partial B_1(0) \subset \mathbb{R}^2$  and consider the function

$$u(x) = \mathbb{H}(\langle a, x \rangle).$$

It is clear that  $u$  solves (3). Note that this  $u$  has flat level sets and defining  $u_\varepsilon(x) = u(x/\varepsilon)$ ,

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon(x) = u_0(x) := \begin{cases} 1 & \langle a, x \rangle > 0 \\ -1 & \langle a, x \rangle < 0 \end{cases}$$

Note that  $\partial\{u_0 = 1\} = \{\langle a, x \rangle = 0\}$ , is a straight line.

**Exercise 5 (+).** Recall that  $\mathbb{H}(t) = \tanh(t/\sqrt{2})$  is the 1-dimensional solution. Show that

$$\sigma := \int_{-\infty}^{\infty} \frac{1}{2} \mathbb{H}'(t)^2 + W(\mathbb{H}(t)) = \int_{-\infty}^{\infty} \mathbb{H}'(t)^2 dt = \int_{-1}^1 \sqrt{2W(s)} ds$$

and compute the value of  $\sigma$ .

**Exercise 6.** For  $u_\varepsilon(x) = \mathbb{H}(\varepsilon^{-1} \langle a, x \rangle)$  on  $\mathbb{R}^2$  considered above, compute the value of  $\lim_{\varepsilon \rightarrow 0} E_\varepsilon(u_\varepsilon; B_1(0))$ . Hint: First compute the limit with  $B_1(0)$  replaced by an appropriately chosen square. Check that  $\mathbb{H}(t)$  is exponentially small as  $t \rightarrow \pm\infty$  to show that the associated error is small.

1.4.2. *The saddle solution and other four ended solutions.* On  $\mathbb{R}^2$ , there are other solutions to Allen–Cahn besides  $\mathbb{H}(\langle a, x \rangle)$ . The following saddle solution was first discovered by Dang–Fife–Peletier [DFP92]. It is an entire solution on  $\mathbb{R}^2$  with  $\{u = 0\} = \{xy = 0\}$ .

**Exercise 7.** Consider

$$\Omega_R := \{(x, y) \in \mathbb{R}^2 : x, y > 0, x^2 + y^2 < R^2\}.$$

Choose  $u_R$  a smooth function with Dirichlet boundary conditions minimizing  $E_1(\cdot)$  among functions in  $H_0^1(\Omega_R)$  (or equivalently smooth functions on  $\Omega_R$  that vanish on the boundary). Show that:

- (a) The function  $u_R$  exists, is smooth, satisfies the Allen–Cahn equation, and does not change sign. Argue that  $u_R$  is either identically zero or (possibly replacing  $u$  by  $-u$ )  $u \in (0, 1)$  in the interior of  $\Omega_R$ .
- (b) Show that  $E_1(u_R; \Omega_R) \leq CR$  for some  $C > 0$  independent of  $R$ . Conclude that  $u_R$  is strictly positive in the interior of  $\Omega_R$ , for  $R$  large.
- (c) Using odd reflections across the coordinate axes, construct  $\tilde{u}_R$  solving the Allen–Cahn equation on  $B_R(0) \subset \mathbb{R}^2$ . Using elliptic regularity, check that  $\tilde{u}_R$  is smooth across the axes and at 0 and has  $E_1(\tilde{u}_R; B_R(0)) \leq \tilde{C}R$ .
- (d) Using elliptic theory, take a subsequential limit as  $R \rightarrow \infty$  to find an entire solution  $u$  to Allen–Cahn whose nodal set is precisely  $\{xy = 0\}$ .

There are many other related solutions. For example:

**Theorem 3** (Kowalczyk–Liu–Pacard [KLP12]). *Given any two lines  $\ell_1, \ell_2 \subset \mathbb{R}^2$  intersecting precisely at the origin, there is a solution  $u$  on  $\mathbb{R}^2$  whose nodal set  $\{u = 0\}$  is asymptotic at infinity to  $\ell_1 \cup \ell_2$ .*

## 2. CONVERGENCE OF (LOCAL) MINIMIZERS OF THE ALLEN–CAHN FUNCTIONAL

The  $\varepsilon \searrow 0$  limit of the Allen–Cahn functional (and associated critical points) turns out to be intimately related with the area functional for hypersurfaces (and associated critical points, minimal surfaces). This relationship was first described in works of Modica and Mortola [MM77] based on the framework of “ $\Gamma$ -convergence” defined by De Giorgi [DGF75].

**Definition 4.** For  $\Omega \subset (M^n, g)$  an open set, a function  $u \in L^1(\Omega)$  is of *bounded variation*,  $u \in BV(\Omega)$ , if its distributional gradient is a Radon measure, i.e. if there is a  $TM$  valued Radon measure  $Du$  so that for any vector field  $X \in C_c^1(\Omega; TM)$

$$\int_{\Omega} u \operatorname{div}_g X \, d\mu_g = - \int_{\Omega} g(X, Du).$$

We write

$$\int_{\Omega} |Du| = \sup \left\{ \int_{\Omega} u \operatorname{div}_g X \, d\mu_g : X \in C_c^1(\Omega; TM), \|X\|_{L^\infty} \leq 1 \right\}$$

for the total variation norm.

**Proposition 5** (BV compactness, cf. [GT01, Theorem 7.22]). *If  $u_k \in BV(\Omega)$  satisfies*

$$\sup_k \left( \|u_k\|_{L^1(\Omega')} + \int_{\Omega'} |Du_k| \right) < \infty,$$

*for all  $\Omega'$  precompact open set in  $\Omega$ , then after passing to a subsequence, there is  $u \in BV_{loc}(\Omega)$ <sup>1</sup> so that  $u_k \rightarrow u$  in  $L^1_{loc}(\Omega)$ <sup>2</sup> and*

$$\int_{\Omega'} |Du| \leq \liminf_{k \rightarrow \infty} \int_{\Omega'} |Du_k|$$

*for all  $\Omega'$  precompact in  $\Omega$ .*

**Remark 6.** If  $\Omega$  has Lipschitz boundary, then we can drop the “loc,” i.e. we could replace  $\Omega'$  by  $\Omega$  throughout.

<sup>1</sup>That is,  $u \in BV(\Omega')$  for all  $\Omega'$  compactly contained in  $\Omega$ .

<sup>2</sup>That is,  $L^1$  convergence on precompact open sets.

**Definition 7.** For a Borel set  $E \subset \Omega$ , we say that  $E$  has *finite perimeter* if  $\chi_E \in BV(\Omega)$ . In this case, we define the *perimeter* of  $E$  by

$$P(E; \Omega) = \int_{\Omega} |D\chi_E|.$$

**Exercise 8 (+).** For  $E$  compact sets with smooth boundary, show that this agrees with the usual notion of perimeter.

Further references for BV functions and sets of finite perimeter include [Sim83, Giu84, AFP00, Mag12].

**2.1.  $\Gamma$ -convergence.** The following computation is rather simple but it underlies the theory of limits of minimizers. Define

$$\Phi(t) := \int_0^t \sqrt{2W(s)} ds.$$

Then, we compute, using AM-GM<sup>3</sup> and the chain rule:

$$\begin{aligned} (4) \quad E_{\varepsilon}(u; \Omega) &= \int_{\Omega} \left( \frac{\varepsilon}{2} |\nabla_g u|^2 + \frac{1}{\varepsilon} W(u) \right) d\mu_g \\ &\geq \int_{\Omega} \sqrt{2W(u)} |\nabla_g u| d\mu_g \\ &= \int_{\Omega} |\nabla_g(\Phi(u))| d\mu_g. \end{aligned}$$

Combined with BV compactness and some measure theoretic arguments we find the following result that loosely speaking says that the behavior of the Allen–Cahn energy is “controlled from below” as  $\varepsilon \rightarrow 0$  by the perimeter functional.

**Proposition 8** ([Mod85, MM77, Ste88, FT89]). *For  $\Omega \subset (M, g)$  a precompact open set, suppose that  $u_{\varepsilon}$  satisfy  $E_{\varepsilon}(u_{\varepsilon}; \Omega) \leq C$ . Then, there is a subsequence  $\varepsilon_k \rightarrow 0$  and  $u_0 \in BV_{loc}(\Omega)$  with  $u_0 \in \{\pm 1\}$  a.e., and*

$$u_{\varepsilon_k} \rightarrow u_0$$

in  $L^1_{loc}(\Omega)$ . Moreover,

$$\sigma P(\{u_0 = 1\}; \Omega') \leq \liminf_{k \rightarrow \infty} E_{\varepsilon_k}(u_{\varepsilon_k}; \Omega'),$$

where  $\sigma = \Phi(1) - \Phi(-1) = \int_{-1}^1 \sqrt{2W(s)} ds$ , for any  $\Omega'$  compactly contained in  $\Omega$ .

*Sketch of the proof.* We can check that  $|\Phi(t)| \leq \alpha + \beta W(t)$ ; thus, the uniform energy bounds  $E_{\varepsilon}(u_{\varepsilon}; \Omega) \leq C$  imply that  $\|\Phi(u_{\varepsilon})\|_{L^1(\Omega)} \leq C$ . Thus, we can use

<sup>3</sup>i.e.,  $2xy \leq x^2 + y^2$ .

(4) and BV compactness to find  $v_0 \in BV_{\text{loc}}(\Omega)$  so that a subsequence of  $\Phi(u_\varepsilon)$  converges in  $L^1_{\text{loc}}(\Omega)$  to  $v_0$  and

$$\int_{\Omega'} |Dv_0| \leq \liminf_{k \rightarrow \infty} E_{\varepsilon_k}(u_{\varepsilon_k}; \Omega').$$

The function  $\Phi$  is invertible, and indeed  $\Phi^{-1}$  is uniformly continuous. Moreover, because  $W(t) \geq ct^4$  for  $t$  sufficiently large, we see that  $\|u_\varepsilon\|_{L^4(\Omega)} \leq C$ . These facts suffice for us to find a further subsequence so that  $u_{\varepsilon_k} \rightarrow u_0 := \Phi^{-1}(v_0)$  in  $L^1_{\text{loc}}(\Omega)$  and that  $u_0 \in BV_{\text{loc}}(\Omega)$  with  $u_0 \in \{\pm 1\}$  a.e. in  $\Omega$ . The fact that  $u_0 \in \{\pm 1\}$  follows from:

$$\frac{4}{\delta^2} |\{x \in \Omega' : |u_{\varepsilon_k}(x)^2 - 1| > \delta\}| \leq \int_{\Omega'} W(u_\varepsilon) d\mu_g \leq C\varepsilon$$

for any  $\delta > 0$ .

Now, we compute

$$\begin{aligned} \Phi(u_0) &= \Phi(1)\chi_{\{u_0=1\}} + \Phi(-1)\chi_{\{u_0=-1\}} \\ &= (\Phi(1) - \Phi(-1))\chi_{\{u_0=1\}} + \Phi(-1)(\chi_{\{u_0\}=1} + \chi_{\{u_0=-1\}}) \\ &= (\Phi(1) - \Phi(-1))\chi_{\{u_0=1\}} + \Phi(-1)\chi_\Omega \end{aligned}$$

a.e. in  $\Omega$ . Hence,

$$\int_{\Omega'} |D\Phi(u_0)| = (\Phi(1) - \Phi(-1))P(\chi_{\{u_0=1\}}; \Omega') = (\Phi(1) - \Phi(-1)) \int_{\Omega'} |Du_0|.$$

This completes the proof.  $\square$

**Exercise 9.** Fill in the details missing in the previous sketch:

- Show that  $|\Phi(t)| \leq \alpha + \beta W(t)$ .
- Show that  $\Phi^{-1}$  exists and is uniformly continuous.
- Show that (after passing to a further subsequence)  $u_{\varepsilon_k}$  converges to  $u_0 := \Phi^{-1}(v_0)$  in measure on  $\Omega$ .<sup>4</sup>
- For  $(X, \mu)$  a measure space with  $\mu(X) < \infty$ , if  $f_i$  are measurable functions converging to  $f$  in measure, and  $\|f_i\|_{L^p(X)} \leq C$  for some  $C > 0, p > 1$ , show that  $f_i \rightarrow f$  in  $L^1(X)$ .
- Conclude that  $u_{\varepsilon_k}$  converges to  $u_0$  in  $L^1(\Omega')$  and thus (passing to a further subsequence via a diagonal argument) a.e. in  $\Omega$ .
- Check that  $u_0 \in BV_{\text{loc}}(\Omega)$  and  $u_0 \in \{\pm 1\}$  a.e. in  $\Omega$ .

The counterpart to the previous result is the following ‘‘recovery’’ result. It says that Proposition 8 is sharp along certain sequences. We emphasize that the sequences  $u_\varepsilon$  constructed below are not critical points (but more on this later).

<sup>4</sup>Recall that  $f_i \rightarrow f$  in measure if  $\lim_{i \rightarrow \infty} \mu(|f_i - f| \geq \delta) = 0$  for all  $\delta > 0$ .

**Proposition 9** ([Mod85, MM77, Ste88]). *If  $E \subset \Omega$  is a set of finite perimeter, then there is a sequence  $u_\varepsilon \in H^1(\Omega) \cap L^4(\Omega)$  with*

$$\sigma P(E; \Omega) = \lim_{\varepsilon \rightarrow 0} E_\varepsilon(u_\varepsilon; \Omega)$$

and  $u_\varepsilon \rightarrow \chi_E - \chi_{\Omega \setminus E}$  in  $L^1(\Omega)$ .

*Sketch of the proof.* Assume that  $\partial E, \partial\Omega$  are smooth and intersect transversally. In particular, the signed distance  $d_{\partial E}(\cdot)$  is smooth near  $\partial E$ . Recall that  $|\nabla d_{\partial E}| = 1$ . We consider  $u_\varepsilon = \varphi(\varepsilon^{-1}d_{\partial E}(\cdot))$  for  $\varphi$  to be chosen. We assume that  $\varphi \equiv \pm 1$  outside of  $[-K, K]$ . Then, writing  $\Sigma_t = \{d_{\partial E}(\cdot) = t\}$  for  $t$  sufficiently close to 0, we find that

$$\begin{aligned} E_\varepsilon(u_\varepsilon; \Omega) &= \int_{\Omega} \left( \frac{1}{2\varepsilon} \varphi'(\varepsilon^{-1}d_{\partial E}(x))^2 + \frac{1}{\varepsilon} W(\varphi(\varepsilon^{-1}d_{\partial E}(x))) \right) d\mu_g \\ &= \int_{-K\varepsilon}^{K\varepsilon} \int_{\Sigma_t} \left( \frac{1}{2\varepsilon} \varphi'(\varepsilon^{-1}t)^2 + \frac{1}{\varepsilon} W(\varphi(\varepsilon^{-1}t)) \right) d\mu_{\Sigma_t} dt \\ &\approx \text{area}(\partial E) \int_{-K\varepsilon}^{K\varepsilon} \left( \frac{1}{2\varepsilon} \varphi'(\varepsilon^{-1}t)^2 + \frac{1}{\varepsilon} W(\varphi(\varepsilon^{-1}t)) \right) dt \\ &\approx \text{area}(\partial E) \int_{-K}^K \left( \frac{1}{2} \varphi'(t)^2 + W(\varphi(t)) \right) dt. \end{aligned}$$

Choosing  $\varphi(t) = \mathbb{H}(t)$  (cut off to  $\pm 1$  outside of  $[-K, K]$ ), we find that

$$E_\varepsilon(u_\varepsilon; \Omega) \approx \text{area}(\partial E) \int_{-\infty}^{\infty} \left( \frac{1}{2} \mathbb{H}'(t)^2 + W(\mathbb{H}(t)) \right) dt = \sigma P(E; \Omega)$$

(see Exercise 5). □

**Remark 10.** The combination of the “lim-inf” lower bound from Proposition 8 for general sequences, with the “recovery” result from Proposition 9, means that “the Allen–Cahn functional  $\Gamma$ -converges to the perimeter functional (times  $\sigma$ ).” This is a rather general phenomenon (first suggested by De Giorgi [DGF75, DG79]) that is very powerful for the study of (local) minimizers of functionals (as we will briefly discuss below). The main downside to using  $\Gamma$ -convergence comes when considering more general critical points (in particular, it does not seem to handle well the issue of “multiplicity” that we will discuss later).

**2.2. Consequences for (local) minimizers.** We will consider  $(M, g)$  a closed Riemannian manifold.

**Definition 11.** We say that a function  $u \in H^1(M) \cap L^4(M)$  is a *strict local minimizer* of  $E_\varepsilon(\cdot)$  if there is  $\delta > 0$  so that  $E_\varepsilon(v) > E_\varepsilon(u)$  for any  $v \in L^1(M)$  with  $0 < \|u - v\|_{L^1(M)} \leq \delta$ . We will also write strict  $\delta$ -local minimizer to emphasize the size of  $\delta$ .

**Exercise 10** (+). A local minimizer is a critical point of  $E_\varepsilon(\cdot)$  and thus satisfies the Allen–Cahn equation.

**Definition 12.** For  $\Omega \subset (M, g)$  precompact, we say that a set  $E \subset \Omega$  *minimizes perimeter* in  $\Omega$  (we will also drop  $\Omega$  when  $\Omega = M$ ) if for any  $E' \subset \Omega$  with  $E\Delta E'$  compactly contained in  $\Omega$ , then

$$P(E; \Omega) \leq P(E'; \Omega).$$

We say that  $E$  is a (*strict*) *local minimizer* if there is  $\delta > 0$  so that the previous holds (with the strict inequality) for  $E'$  with  $0 < \|\chi_E - \chi_{E'}\|_{L^1(\Omega)} \leq \delta$ .

**Proposition 13.** *Suppose that  $u_\varepsilon$  is a sequence of  $\delta$ -local minimizers of  $E_\varepsilon(\cdot)$  in  $(M, g)$ . Assume that  $E_\varepsilon(u_\varepsilon) \leq C$ . Then, after passing to a subsequence  $\varepsilon_k \rightarrow 0$ ,  $u_{\varepsilon_k} \rightarrow u_0$  in  $L^1(M)$ , with  $u_0 \in BV(M)$  and  $u_0 \in \{\pm 1\}$  a.e. in  $M$ . The set  $E := \{u_0 = 1\}$  is a local minimizer of perimeter.*

*Proof.* We only need to prove that  $E$  is a local minimizer of perimeter. If not, there is  $\tilde{E}$  with

$$\|\chi_E - \chi_{\tilde{E}}\|_{L^1(M)} < \frac{\delta}{2}$$

and  $P(\tilde{E}) < P(E)$ . Using Proposition 9, we can find  $\tilde{u}_{\varepsilon_k}$  with  $\tilde{u}_{\varepsilon_k} \rightarrow \chi_{\tilde{E}}$  in  $L^1(M)$  and

$$\lim_{k \rightarrow \infty} E_{\varepsilon_k}(\tilde{u}_{\varepsilon_k}) = \sigma P(\tilde{E})$$

On the other hand, we compute

$$\begin{aligned} \|\tilde{u}_{\varepsilon_k} - u_{\varepsilon_k}\|_{L^1(M)} &\leq \|\tilde{u}_{\varepsilon_k} - (\chi_{\tilde{E}} - \chi_{M \setminus \tilde{E}})\|_{L^1(M)} + \|u_{\varepsilon_k} - (\chi_E - \chi_{M \setminus E})\|_{L^1(M)} \\ &\quad + \|(\chi_E - \chi_{M \setminus E}) - (\chi_{\tilde{E}} - \chi_{M \setminus \tilde{E}})\|_{L^1(M)} \\ &= \|\tilde{u}_{\varepsilon_k} - (\chi_{\tilde{E}} - \chi_{M \setminus \tilde{E}})\|_{L^1(M)} + \|u_{\varepsilon_k} - (\chi_E - \chi_{M \setminus E})\|_{L^1(M)} \\ &\quad + 2\|\chi_E - \chi_{\tilde{E}}\|_{L^1(M)} \\ &< \delta + o(1) \end{aligned}$$

as  $k \rightarrow \infty$ . Thus, for  $k$  sufficiently large, we find that

$$E_{\varepsilon_k}(\tilde{u}_{\varepsilon_k}) \geq E_{\varepsilon_k}(u_{\varepsilon_k})$$

Thus, we find that

$$\sigma P(E) \leq \liminf_{k \rightarrow \infty} E_{\varepsilon_k}(u_{\varepsilon_k}) \leq \liminf_{k \rightarrow \infty} E_{\varepsilon_k}(\tilde{u}_{\varepsilon_k}) = \sigma P(\tilde{E})$$

This is a contradiction. This completes the proof.  $\square$

The following result is a sort of strengthening of the “recovery” result in Proposition 9 in the sense that it finds (for local minimizers of perimeter) recovery sequences that are again themselves local minimizers.

**Proposition 14** (Kohn–Sternberg [KS89]). *Suppose that  $E \subset (M, g)$  is a local minimizer of perimeter. Then, for  $\varepsilon$  sufficiently small, there exists  $u_\varepsilon$  solving the Allen–Cahn equation and locally minimizing  $E_\varepsilon(\cdot)$  so that  $u_\varepsilon \rightarrow (\chi_E - \chi_{M \setminus E})$  in  $L^1(M)$  and  $\lim_{\varepsilon \rightarrow 0} E_\varepsilon(u_\varepsilon) = \sigma P(E)$ .*

**Exercise 11.** Prove this. Hint: minimize  $E_\varepsilon(\cdot)$  in a (closed)  $L^1$ -ball centered at  $\chi_E - \chi_{M \setminus E}$ . Is the minimizer in the interior of the ball or at the boundary?

### 2.3. Some minimal surface facts.

**Proposition 15** (De Giorgi, Fleming, Almgren, Federer, Simons). *If a set  $E \subset (M^n, g)$  is a local minimizer of perimeter for  $3 \leq n \leq 7$ , then, after changing  $E$  by a set of measure zero, the topological boundary of  $E$ ,  $\partial E$ , is a smooth hypersurface.*

The hypersurface  $\partial E$  is a minimal hypersurface in the sense that its *mean curvature* vanishes

$$H_{\partial E} = \text{tr}_{T\partial E} A_{\partial E}(\cdot, \cdot) = 0,$$

for  $A_{\partial E}$  the second fundamental form. This is equivalent to the following property: if  $\Sigma_t$  is smooth family of hypersurfaces for  $t \in (-\delta, \delta)$  with  $\Sigma_0 = \partial E$ , then

$$\left. \frac{d}{dt} \right|_{t=0} \text{area}_g(\Sigma_t) = 0.$$

Suppose that  $\Sigma_t$  is the image of  $F_t : \Sigma \rightarrow (M, g)$  with

$$\left. \frac{\partial F_t}{\partial t} \right|_{t=0} F_t = \varphi \nu$$

for  $\nu$  the unit normal to  $\Sigma_0$ . Then, we have

$$\left. \frac{d^2}{dt^2} \right|_{t=0} \text{area}_g(\Sigma_t) = \int_{\Sigma} \varphi J\varphi d\mu := \mathcal{Q}_{\Sigma}(\varphi, \varphi).$$

for  $J\varphi = -\Delta_{\Sigma}\varphi - (|A_{\Sigma}|^2 + \text{Ric}_g(\nu, \nu))\varphi$ . If  $E$  is a local minimizer of perimeter, then the usual calculus characterization of a local minimizer gives that

$$\int_{\Sigma} \varphi J\varphi d\mu_t \geq 0$$

for any such  $\varphi$ . In general,  $\Sigma$  satisfying this condition is *stable*.

## 3. THE PACARD–RITORÉ CONSTRUCTION

It turns out that solutions to Allen–Cahn exist near minimal surfaces  $\Sigma$  beyond just local minimizers, e.g. for unstable  $\Sigma$ . We say that  $\Sigma$  is *non-degenerate* if  $\ker J = \{0\}$ .

**Theorem 16** (Pacard–Ritoré [PR03], cf. [Pac12]). *If  $\Sigma^{n-1} \subset (M^n, g)$  is a smooth non-degenerate minimal hypersurface that divides  $M$  into two pieces, then for  $\varepsilon_0 = \varepsilon_0(\Sigma, M, g) > 0$  sufficiently small, there exists  $\{u_\varepsilon\}_{\varepsilon \in (0, \varepsilon_0)}$  solving*

$$\varepsilon^2 \Delta_g u_\varepsilon = W'(u_\varepsilon)$$

and so that  $u_\varepsilon$  approximates  $\Sigma$  in the sense that  $u_\varepsilon$  converges to 1 on one side of  $\Sigma$  and  $-1$  on the other side and so that

$$\lim_{\varepsilon \rightarrow 0} E_\varepsilon(u_\varepsilon) = \sigma \operatorname{area}_g(\Sigma).$$

*Idea of the proof.* When  $\Sigma$  is not a local minimizer, the  $\Gamma$ -convergence/minimization approach is no longer straightforward (however see [JS09]). Instead, the proof proceeds via an “infinite dimensional Liapunov–Schmidt reduction.”

The basic idea is to find a first approximation to  $u_\varepsilon$  built out of  $\Sigma$  and the 1-dimensional solution  $\mathbb{H}(t)$ . This will solve the Allen–Cahn equation up to a reasonably good error (here, minimality of  $\Sigma$  is used). Then, this ansatz is perturbed appropriately to solve away the error (this is the step where non-degeneracy of  $\Sigma$  is used).

We use Fermi coordinates around  $\Sigma$ , i.e., coordinates  $(y, z)$  on an fixed tubular neighborhood  $U$  of  $\Sigma$  where, for  $y \in \Sigma$ ,  $(y, z)$  corresponds to the point

$$Z_\Sigma(y, z) = \exp_y(z\nu(y)) \in U.$$

In the  $(y, z)$  coordinates, the Laplace–Beltrami operator associated to  $g$  becomes

$$(5) \quad \Delta = \Delta_{\Gamma_z} + \partial_z^2 + H_{\Gamma_z} \partial_z$$

where  $\Gamma_z$  is the hypersurface  $\{Z_\Sigma(y, z) : y \in \Sigma\}$  and  $H_{\Gamma_z}$  is the mean curvature of  $\Gamma_z$ .

Setting  $u_1(y, z) = \mathbb{H}(\varepsilon^{-1}z)$ , we compute

$$\begin{aligned} \varepsilon^2 \Delta_g u_1 &= \mathbb{H}''(\varepsilon^{-1}z) + \varepsilon H_{\Gamma_z} \mathbb{H}'(\varepsilon^{-1}z) \\ &= W'(\mathbb{H}(\varepsilon^{-1}z)) + \varepsilon H_{\Gamma_z} \mathbb{H}'(\varepsilon^{-1}z) \\ &= W'(u_1) + \varepsilon H_{\Gamma_z} \mathbb{H}'(\varepsilon^{-1}z). \end{aligned}$$

Thus, this choice of  $u$  solves the Allen–Cahn equation up to an error term  $\mathcal{E}_1 = \varepsilon H_{\Gamma_z} \mathbb{H}'(\varepsilon^{-1}z)$ . To estimate the size (we’ll estimate the  $C^0$ -norm, but similar estimates hold for higher derivatives) of  $\mathcal{E}_1$ , note that  $H_{\Gamma_z} = O(|z|)$  (because  $\Sigma$  is minimal), so writing  $t = \varepsilon^{-1}z$ , we have

$$|\mathcal{E}_1| \lesssim \varepsilon^2 \mathbb{H}'(t).$$

The general strategy is to consider  $u = u_1 + v$ , where  $v$  is an error term. In reality, we must consider a better ansatz (roughly) of the form  $u_2(y, z) = \mathbb{H}(\varepsilon^{-1}(z - \zeta(y)))$  for some function  $\zeta(y)$  chosen to cancel the influence of  $H_{\Gamma_z}$  to one higher order. This is the step where the non-degeneracy of  $\Sigma$  is used.

In any case, we would like to solve

$$0 = \varepsilon^2 \Delta_g u - W'(u) = \underbrace{\varepsilon^2 \Delta_g v + W'(u_1) - W'(u_1 + v)}_{(*)} + \mathcal{E}_1.$$

Note that the linearization of  $(*)$  around  $v = 0$  is

$$\mathcal{L}v := \varepsilon^2 \Delta_g v - W''(u_1)v$$

As such, it is crucial to understand the kernel of  $\mathcal{L}$  (we can then try to solve the above equation by using a contraction mapping fixed point argument). As such, a fundamental issue is to understand the kernel of  $\mathcal{L}$  (so that we can invert  $\mathcal{L}$  appropriately). See Proposition 17 below.  $\square$

As indicated above, the following result is key in the Pacard–Ritoré construction (and in many other related constructions). It also plays an important role in the Wang–Wei [WW17] regularity theory discussed later.

**Exercise 12.** Verify the expression for the Laplacian in Fermi coordinates in (5).

**Proposition 17** (cf. [PR03, Corollary 7.5] and [Pac12]). *Suppose that  $w \in L^\infty(\mathbb{R}^{n-1} \times \mathbb{R})$  satisfies*

$$L_* w := \Delta w - W''(\mathbb{H}(x^n))w = 0.$$

*Then  $w(x', x^n) = c\mathbb{H}'(x^n)$  for some  $c \in \mathbb{R}$ .*

**Exercise 13.** This exercise is related Proposition 17.

- (a) Check that  $\mathbb{H}'(x^n) \in L^\infty(\mathbb{R}^n)$  satisfies  $L_*(\mathbb{H}'(x^n)) = 0$ .
- (b) Prove Proposition 17 when  $n = 1$ . Hint, compute  $(\log \mathbb{H}'(t))''$  and multiply by  $u(t)^2$  (an arbitrary function with compact support). Conclude that

$$\int_{-\infty}^{\infty} u'(t)^2 + W''(\mathbb{H}(t))u(t)^2 dt = \int_{-\infty}^{\infty} (\mathbb{H}'(t)^{-1}\mathbb{H}''(t)u(t) - u'(t))^2 dt$$

Choose  $u(t) = w(t)\varphi_R$  where  $\varphi_R$  cuts off from  $R$  to  $2R$ . Letting  $R \rightarrow \infty$ , conclude that

$$w'(t) = w(t)\mathbb{H}'(t)^{-1}\mathbb{H}''(t)$$

and use this to complete the proof.

- (c) Using a logarithmic cutoff function, show that a similar proof works for  $n = 2$ .
- (d) Returning to  $n = 1$ , argue (by contradiction) that there is some  $\mu > 0$  so that if  $u(t)$  satisfies  $\int_{-\infty}^{\infty} u(t)\mathbb{H}'(t)dt = 0$ , then

$$\int_{-\infty}^{\infty} u'(t)^2 + W''(\mathbb{H}(t))u(t) dt \geq \mu \int_{-\infty}^{\infty} u(t)^2 dt.$$

- (e) For  $n > 2$ , write a solution to  $L_*w = 0$  as  $w(x', x^n) = c(x')\mathbb{H}'(x^n) + \bar{w}(x', x^n)$  where

$$\int_{-\infty}^{\infty} \bar{w}(x', t)\mathbb{H}'(t)dt = 0$$

for each  $x' \in \mathbb{R}^{n-1}$ . Show that  $c$  is bounded and harmonic (and thus constant by Liouville's theorem). Hint: write  $L_*w = 0$  in terms of  $c$  and  $\bar{w}$ , multiply by  $\mathbb{H}'(x^n)$ , and integrate with respect to  $x^n$ .

- (f) Assume that  $\bar{w}$  and its derivatives tend to zero as  $x^n \rightarrow \pm\infty$  rapidly enough to justify differentiating under the integral sign<sup>5</sup> for

$$V(x') := \int_{-\infty}^{\infty} \bar{w}(x', t)^2 dt.$$

Show that the equation satisfied by  $\bar{w}$  (given that  $c$  is constant) implies that

$$\Delta_{\mathbb{R}^{n-1}}V - \mu V \geq 2 \int_{-\infty}^{\infty} |\nabla \bar{w}(x', t)|^2 dt \geq 0.$$

- (g) Multiply  $V$  by a well chosen function and integrate by parts to conclude that  $V \equiv 0$ , finishing the proof of Proposition 17.

**3.1. The Morse index.** For  $\Sigma^{n-1} \subset (M^n, g)$  a (closed) minimal (two-sided<sup>6</sup>) surface, recall that

$$\left. \frac{d^2}{dt^2} \right|_{t=0} \text{area}_g(\Sigma_t) = \int_{\Sigma} \varphi J\varphi d\mu := \mathcal{Q}_{\Sigma}(\varphi, \varphi).$$

is the second variation for  $J\varphi = -\Delta_{\Sigma}\varphi - (|A_{\Sigma}|^2 + \text{Ric}_g(\nu, \nu))\varphi$ .

**Definition 18.** The *Morse index* of  $\Sigma$  is the largest dimension of a linear subspace  $W \subset C^{\infty}(\Sigma)$  so that for  $\varphi \in W \setminus \{0\}$ ,  $\mathcal{Q}_{\Sigma}(\varphi, \varphi) < 0$ .

**Exercise 14.** For  $M = \mathbb{S}^n = \{x \in \mathbb{R}^{n+1} : |x| = 1\}$  and  $\Sigma = \{x^n = 0\} \cap \mathbb{S}^n$ , check that  $\Sigma$  is minimal and has Morse index 1.

Similarly, for a function  $u_{\varepsilon}$  on  $(M^n, g)$  solving the Allen–Cahn equation  $\varepsilon^2 \Delta_g u_{\varepsilon} = W'(u_{\varepsilon})$ , we define

$$\left. \frac{d^2}{dt^2} \right|_{t=0} E_{\varepsilon}(u_{\varepsilon} + t\psi) := \mathcal{Q}_{u_{\varepsilon}}(\psi, \psi).$$

**Definition 19.** The *Morse index* of  $u_{\varepsilon}$  is the largest dimension of a linear subspace  $W \subset C^{\infty}(M)$  so that for  $\psi \in W \setminus \{0\}$ ,  $\mathcal{Q}_{u_{\varepsilon}}(\psi, \psi) < 0$ .

It turns out that in “multiplicity one” situations, the Allen–Cahn index and the limiting minimal surface index agree. See [CM18] (and also [Le11, Hie17, Gas17]).

<sup>5</sup>This can be justified by a barrier argument, cf. [Pac12, p. 18]

<sup>6</sup>Two-sided means that there is a consistent choice of unit normal. Compare to  $\mathbb{R}P^2 \subset \mathbb{R}P^3$ . Note that if  $\Sigma = \partial E$ , then  $\Sigma$  is necessarily two-sided.

## 4. STABLE/BOUNDED INDEX SOLUTIONS TO ALLEN–CAHN EQUATION

It turns out that some manifolds do not admit any stable minimal surfaces or stable solutions to Allen–Cahn.

**Exercise 15** (+). Suppose that  $u_\varepsilon$  is a solution to Allen–Cahn on  $(M^n, g)$ .

(a) Show that

$$\mathcal{Q}_{u_\varepsilon}(\psi, \psi) = \int_M \left( \varepsilon |\nabla \psi|^2 + \frac{1}{\varepsilon} W''(u_\varepsilon) \psi^2 \right) d\mu_g.$$

(b) Verify the Bochner formula

$$\frac{1}{2} \Delta_g |\nabla f|^2 = |D^2 f|^2 + g(\nabla \Delta_g f, \nabla f) + \text{Ric}_g(\nabla f, \nabla f),$$

and use it to show that<sup>7</sup>

$$\begin{aligned} & \mathcal{Q}_{u_\varepsilon}(\psi |\nabla u_\varepsilon|, \psi |\nabla u_\varepsilon|) \\ &= \int_M (|\nabla \psi|^2 |\nabla u_\varepsilon|^2 - (|D^2 u_\varepsilon|^2 - |\nabla |\nabla u_\varepsilon||^2) + \text{Ric}_g(\nabla u_\varepsilon, \nabla u_\varepsilon)) \psi^2 d\mu_g \end{aligned}$$

Suppose now that  $(M^n, g)$  has positive Ricci curvature.

(c) Show that there are no stable minimal (two-sided) hypersurfaces.

(d) Show that there are no stable solutions to Allen–Cahn.

Thus, we see that one must turn to unstable solutions (both for minimal surfaces and Allen–Cahn). Our general goal will be: “how can the Allen–Cahn equation be used to find (unstable) minimal surfaces? Moreover, what properties can be proved about the limiting surfaces and the limiting process?”

**4.1. Guaraco–Gaspar existence theory.** Consider  $(M^n, g)$  closed Riemannian manifold. Recall that  $\pm 1$  are the (unique) global minimizers for  $E_\varepsilon(\cdot)$  (and they both have the same energy 0). This calls for the mountain-pass theorem! Because  $E_\varepsilon(\cdot)$  is defined on  $H^1(M)$  an infinite dimensional space, one must check the so-called *Palais–Smale* condition. The idea of this (in reality, one must slightly modify this argument) is contained in the following exercise:

**Exercise 16.** For  $\varepsilon > 0$  fixed, consider  $u_k \in H^1(M)$  with  $|u_k| \leq 1$ ,

$$\|u_k\|_{H^1(M)} + E_\varepsilon(u_k) \leq C,$$

and  $DE_\varepsilon|_{u_k} \rightarrow 0$  weakly in  $H^1(M)$  in the sense that

$$DE_\varepsilon|_{u_k}(v) = \int_M \left( \varepsilon g(\nabla u_k, \nabla v) + \frac{1}{\varepsilon} W'(u_k)v \right) d\mu_g \rightarrow 0$$

<sup>7</sup>To justify plugging this into the stability operator, consider  $\psi \sqrt{|\nabla u_\varepsilon|^2 + \delta}$  for  $\delta > 0$  and send  $\delta \rightarrow 0$ .

for each  $v \in H^1(M)$ . Show that a subsequence of  $u_k$  converges *strongly* in  $H^1(M)$  to  $u$  which is a solution to the Allen–Cahn equation on  $(M, g)$ . Hint: for a weak subsequential limit  $u$  of the  $u_k$ , check that  $u$  solves the Allen–Cahn equation. Then, relate  $\int |\nabla(u_k - u)|^2 d\mu_g$  to  $DE_\varepsilon|_{u_k}(u_k - u)$  (up to other terms tending to zero).

This implies (when combined with energy bounds for paths between  $+1$  and  $-1$ ) the following result.

**Theorem 20** (Guaraco [Gua18]). *For  $(M^n, g)$  a closed Riemannian manifold and  $\varepsilon > 0$  sufficiently small there exists  $u_\varepsilon$  solving the Allen–Cahn equation with  $\text{index}(u_\varepsilon) \leq 1$  and  $C^{-1} \leq E_\varepsilon(u_\varepsilon) \leq C$ .*

In fact, higher index critical points exist as well.

**Theorem 21** (Gaspar–Guaraco [GG18a]). *For  $(M^n, g)$  a closed Riemannian manifold,  $p \in \{1, 2, \dots\}$  and  $\varepsilon > 0$  sufficiently small (depending on  $p$ ), there exists  $u_\varepsilon$  solving the Allen–Cahn equation with<sup>8</sup>  $\text{index}(u_\varepsilon) \leq p$  and  $E_\varepsilon(u_\varepsilon) \simeq p^{\frac{1}{n}}$ .*

These critical points are not local minimizers, so it is necessary to use a different theory to take the limit as  $\varepsilon \rightarrow 0$ . This is based on the regularity of limits of stable solutions to Allen–Cahn (via the following fact).

**Exercise 17.** Suppose that  $u_\varepsilon$  is as in Theorem 21. Show that there are at most  $p$  points  $\{x_1, \dots, x_p\} \subset M$  so that for  $x \in M \setminus \{x_1, \dots, x_p\}$ , there is a ball  $B_r(x) \subset M$  so that  $u_\varepsilon$  is stable for deformations supported in  $B_r(x)$ .

**4.2. The Hutchinson–Tonogawa–Wickramasekera regularity theory.** We have seen that on any Riemannian manifold there are many solutions to the Allen–Cahn equation with bounded index. It is thus natural to try to take the limit of these solutions as  $\varepsilon \rightarrow 0$ .

The following is the combination of several deep works: Hutchinson–Tonogawa [HT00] considered limits of critical points with bounded energy and proved the limiting object is a stationary integral varifold (a weak notion of submanifold). Later, Tonogawa–Wickramasekera [TW12] considered stable critical points and proved regularity of the limiting hypersurface (using earlier work of Wickramasekera [Wic14] and Schoen–Simon [SS81] on stable minimal hypersurfaces). Finally, this was generalized to bounded index solutions by Guaraco [Gua18].

**Theorem 22.** *Suppose that  $u_\varepsilon$  are solutions to the Allen–Cahn equation with  $E_\varepsilon(u_\varepsilon) \leq C$  and  $\text{index}(u_\varepsilon) \leq p$ . Then, there is  $\Sigma_1 \dots \Sigma_k$  embedded minimal hypersurfaces that are smooth (outside a singular set of codimension 7) and disjoint, as well as positive integers  $m_1, \dots, m_k$  so that a subsequence of  $u_\varepsilon$  converges to  $\Sigma = \cup_{i=1}^k \Sigma_i$  in the following sense:*

<sup>8</sup>There are exactly  $p$  (counted with multiplicity) non-positive eigenvalues of the stability inequality.

- (i)  $u_\varepsilon$  converges in  $L^1$  to  $u_0$  which is equal to one of  $-1$  or  $+1$  on the various components of  $M \setminus \Sigma$ .
- (ii) A subsequence of the measures  $\varepsilon|\nabla u_\varepsilon|^2 d\mu_g$  converge (weakly) to

$$\mu = \sigma \sum_{i=1}^k m_i \mathcal{H}^{n-1}|_{\Sigma_i}.$$

In fact, there is a varifold  $V_{u_\varepsilon}$  that converges to the varifold associated to  $\mu$ , but we will not discuss this here.

A key feature that is present for stable/bounded index surfaces that was not present for minimizers is the presence of multiplicity. For example:

**Exercise 18.** (a) Show that the following situation is not possible:  $u_\varepsilon$  are  $\delta$ -local minimizers of the Allen–Cahn energy and  $\varepsilon|\nabla u_\varepsilon|^2 d\mu_g$  converges weakly to the measure  $m\mathcal{H}^{n-1}|_\Sigma$  for some smooth minimal hypersurface  $\Sigma$  and integer  $m > 1$ . You can assume the following fact (see [HT00, Theorem 1(a)]: if this occurs, then

$$\mu = \sigma m \mathcal{H}^{n-1}|_\Sigma = \lim_{k \rightarrow \infty} \varepsilon_k |\nabla u_{\varepsilon_k}|^2 d\mu_g = \lim_{k \rightarrow \infty} \frac{2}{\varepsilon_k} W(u_{\varepsilon_k}) d\mu_g$$

where the limits are in the sense weak\* convergence of measures (everything besides the final equality here is already asserted in Theorem 22).<sup>9</sup>

- (b) Show that multiplicity can occur for limits of stable solutions. Hint: consider  $(M^n, g)$  a closed manifold containing a region isometric to a warped product on  $(-1, 1) \times \mathbb{S}^{n-1}$  with metric  $dt^2 + f(t)^2 g_{\mathbb{S}^{n-1}}$ . Choose  $f$  so that there is a sequence  $\tau_i \rightarrow 0$  with  $\{\pm\tau_i\} \times \mathbb{S}^{n-1}$  are non-degenerate stable minimal surfaces. Use Theorem 16. Alternatively, you can use the fact that non-degenerate stable surfaces are locally  $L^1$ -minimizing [Whi94, MR10] and apply Proposition 14.

The latter option shows that the resulting solutions are stable (why?), while if one applies Theorem 16, then to prove that the solutions are stable one can refer to e.g. [Hie17, Gas17, CM18].

- (c) Check that  $\{0\} \times \mathbb{S}^{n-1}$  is a degenerate minimal surface. Compare with Theorem 25 below.
- (d) What is the  $L^1$ -limit of the solutions  $u_\varepsilon$  constructed in (b)? How is this relevant for the  $\Gamma$ -convergence theory of non-minimizing stable solutions (convince yourself that the answer to this part shows that  $u_\varepsilon$  are not  $\delta$ -minimizing for  $\delta$  uniform as  $\varepsilon \rightarrow 0$ )?

The presence of multiplicity is rather undesirable from the point of view of using the Allen–Cahn equation to construct minimal hypersurfaces. For example,

<sup>9</sup>This fact is known as “equidistribution of energy,” i.e., both terms in the Allen–Cahn energy functional contribute the same amount in the (weak) limit.

Gaspar–Guaraco construct (cf. Theorem 21) Allen–Cahn solutions for each  $p$  which we might expect to give distinct minimal surfaces (perhaps of index  $p$ ) in the limit. However, a priori it could happen that each of these solutions limits to  $m_p \Sigma$  for some  $\Sigma$  fixed, but with multiplicity  $m_p$  depending on  $p$ .

**Definition 23.** A Riemannian manifold  $(M^n, g)$  is *bumpy* if no immersed minimal hypersurface is degenerate.

**Theorem 24** (White [Whi91, Whi17]). *Any metric  $g$  can be perturbed slightly to become bumpy.*<sup>10</sup>

In a joint work with Mantoulidis, we recently obtained the following result.

**Theorem 25** ([CM18]). *If  $(M^3, g)$  is bumpy and  $u_\varepsilon$  solves the Allen–Cahn equation with bounded energy and index, i.e.,  $E_\varepsilon(u_\varepsilon) \leq C$  and  $\text{index}(u_\varepsilon) \leq I_0$ , then the limiting minimal surface  $\Sigma$  occurs with multiplicity one (and is two-sided).*

This resolves (in the Allen–Cahn setting) the *multiplicity one* conjecture of Marques–Neves [MN16, Mar14, Nev14, MN18]. In particular, it has the following consequence.

**Corollary 26** ([CM18]). *If  $(M^3, g)$  is a Riemannian manifold with a bumpy metric, then for each positive integer  $p$ ,  $(M, g)$  contains a two-sided minimal surface  $\Sigma_p$  with  $\text{index}(\Sigma_p) = p$  and  $\text{area}(\Sigma_p) \simeq p^{\frac{1}{3}}$ .*

In particular such an  $(M^3, g)$  has infinitely many minimal surfaces.

**Remark 27.** It was conjectured by Yau [Yau82] that every 3-manifold contains infinitely many (immersed) minimal surfaces. This was originally proven for manifolds of positive Ricci curvature by Marques–Neves [MN17], and for generic metrics (with the conclusion that the set of minimal surfaces is dense) by Irie–Marques–Neves [IMN18] (see also [LMN18] and [GG18b]). Very recently, Song has proven Yau’s conjecture for all metrics (not just a generic set) [Son18]. These results hold in ambient dimensions  $3 \leq n \leq 7$ .

These proofs proceed via very clever arguments by contradiction (relying on the min-max construction of minimal surfaces due to Almgren–Pitts). As such, they do not seem to get any information concerning the area or index of the obtained surfaces as in Corollary 26. At the moment, the Allen–Cahn equation is the only known approach to prove such a result (and the results are presently limited to ambient dimension 3).

Theorem 25 builds on foundational work of Wang–Wei [WW17] who proved curvature estimates for stable solutions to Allen–Cahn on 2-dimensional surfaces and developed a general framework for understanding solutions to Allen–Cahn

---

<sup>10</sup>Actually, the set of bumpy metrics is generic in the Baire category sense, i.e., the countable intersection of open dense sets.

with Lipschitz level sets as  $\varepsilon \rightarrow 0$ . We will not discuss this in generality here, but will only mention one ingredient.

In analyzing solutions of Allen–Cahn as  $\varepsilon \rightarrow 0$  it is very useful to have a picture for the interface  $\{u = 0\}$  at scale  $O(\varepsilon)$ . We have seen numerous examples where the interface behaves in a complicated way at the scale of  $(M, g)$ , but sometimes the small-scale behavior of solutions is much simpler.

**Exercise 19 (+).** Suppose that  $u_\varepsilon$  is a sequence of functions solving the Allen–Cahn equation on  $(M, g)$ . Choose  $x_j \in M$  and  $\varepsilon_j \rightarrow 0$ , and consider for  $K$  fixed (large) the ball  $B_{K\varepsilon_j}(x_j) \subset M$ .

- (a) Make sense of what it means to “zoom in” by scale  $\varepsilon_j^{-1}$  by defining a new metric  $g_j$  on  $B_K$  and a rescaled function  $\tilde{u}_j$ .
- (b) Show that  $g_j$  converges smoothly to the flat Euclidean metric on  $B_K(0) \subset \mathbb{R}^n$  and (after passing to a subsequence)  $\tilde{u}_j$  converges smoothly to  $\tilde{u}$  solving the Allen–Cahn equation on  $B_K$  with  $\varepsilon = 1$ .
- (c) If  $u_{\varepsilon_j}$  was stable in  $B_\rho(x_j)$  for some  $\rho > 0$  fixed, show that  $\tilde{u}$  is stable in  $B_K(0)$  for compactly supported variations.
- (d) If  $\text{index}(u_{\varepsilon_j}; B_\rho(x_j)) \leq I_0$  show the same for  $\tilde{u}$ .
- (e) If  $u_{\varepsilon_j}$  was  $\delta$ -locally minimizing, what property does  $\tilde{u}$  satisfy?
- (f) Writing  $\tilde{u}_K$  to emphasize the choice of  $K$ , show that we can send  $K \rightarrow \infty$  to find an entire solution to the Allen–Cahn equation on  $\mathbb{R}^n$  with  $\varepsilon = 1$ . What happens in cases (c)-(e)?

## 5. ENTIRE SOLUTIONS TO THE ALLEN–CAHN EQUATION

Exercise 19 motivates the study of entire solutions to Allen–Cahn with  $\varepsilon = 1$  on  $\mathbb{R}^n$  with various additional conditions (e.g., stability, bounded index, minimizing). Interestingly, this is not the original motivation for the study of entire solutions to Allen–Cahn. Instead, a motivating problem in the study of the Allen–Cahn equation has been the following conjecture of De Giorgi made in 1978:

**Conjecture 28** (De Giorgi [DG79]). *Consider  $u \in C^2(\mathbb{R}^n)$  solving the Allen–Cahn equation*

$$\Delta u = W'(u) = u^3 - u$$

*so that  $|u| \leq 1$  and  $\frac{\partial u}{\partial x^n} > 0$ . At least for  $n \leq 8$ , is it true that  $u(x) = \mathbb{H}(\langle a, x \rangle)$  is the one-dimensional solution?*

This conjecture (and particularly the monotonicity  $\frac{\partial u}{\partial x^n} > 0$  condition) here is motivated by the classical Bernstein conjecture for minimal surfaces.

**Theorem 29** (Bernstein [Ber27], Fleming [Fle62], De Giorgi [DG65], Almgren [Alm66], Simons [Sim68], Bombieri–De Giorgi–Giusti [BDGG69]). *Suppose that*

$u : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  has the property that  $\text{graph}(u) \subset \mathbb{R}^n$  is a minimal surface. Equivalently,

$$\sum_{i=1}^{n-1} D_i \left( \frac{D_i u}{\sqrt{1 + |Du|^2}} \right) = 0.$$

Then, for  $n \leq 8$ ,  $u(x) = \langle x, a \rangle + b$  is an affine function. For  $n > 8$ , non-flat minimal graphs exist.

Unlike the Bernstein conjecture for minimal surfaces, the De Giorgi conjecture is not resolved in its entirety. It is completely understood in low dimensions

**Theorem 30** (Ghoussoub–Gui [GG98] ( $n = 2$ ), Ambrosio–Cabre [AC00] ( $n = 3$ )). For  $n = 2, 3$ , consider  $u \in C^2(\mathbb{R}^n)$  solving the Allen–Cahn equation with  $|u| \leq 1$  and  $\frac{\partial u}{\partial x^n} > 0$ . Then,  $u(x) = \mathbb{H}(\langle a, x \rangle)$ .

It is also completely understood in high dimensions (here, the dimensional restriction is expected to be sharp).

**Theorem 31** (del Pino–Kowalczyk–Wei [dPKW11]). For  $n \geq 9$ , there is  $u \in C^\infty(\mathbb{R}^n)$  solving the Allen–Cahn equation with  $|u| \leq 1$  and  $\frac{\partial u}{\partial x^n} > 0$  that does not have flat level sets.

For  $n \in \{4, 5, \dots, 8\}$ , the De Giorgi conjecture is still open. However, it can be solved with an additional hypothesis:

**Theorem 32** (Savin [Sav09]). For  $n \leq 8$ , consider  $u \in C^2(\mathbb{R}^n)$  solving the Allen–Cahn equation with  $|u| \leq 1$  and  $\frac{\partial u}{\partial x^n} > 0$ . Assume in addition that

$$\lim_{x^n \rightarrow \pm 1} u(x) = \pm 1.$$

Then,  $u(x) = \mathbb{H}(\langle a, x \rangle)$ .

See also [Wan17a].

**5.1. Stability and minimizing properties of monotone solutions.** Recall that  $u \in C^2(\mathbb{R}^n)$  solving the Allen–Cahn equation is *stable* if it is stable on compact sets, and *minimizing* if it minimizes  $E_1(\cdot)$  on compact sets.

**Exercise 20 (+).** Consider  $u$  solving the Allen–Cahn equation on  $\mathbb{R}^n$  with  $\frac{\partial u}{\partial x^n} > 0$ .

- (a) Set  $v = \frac{\partial u}{\partial x^n} > 0$ . Show that  $v$  satisfies the linearized equation  $\Delta v = W''(u)v$ .
- (b) Show that  $u$  is stable. Hint: consider the first eigenfunction associated to  $\mathcal{Q}_u(\cdot, \cdot)$  with Dirichlet boundary conditions on  $B_R(0)$ . Use the maximum principle combined with (a) to control the sign of the associated eigenvalue.
- (c) Conclude, in particular, that the 1-dimensional solution  $u(x) = \mathbb{H}(\langle a, x \rangle)$  is stable.

**Exercise 21.** (a) Suppose that  $f_1, f_2$  both solve the Allen–Cahn equation on  $B_R$ ,  $f_1 \leq f_2$  on  $B_R$  with  $f_1(x) = f_2(x)$  for some  $x$  with  $|x| < R$ . Show that  $f_1 \equiv f_2$  on  $B_R$ . Hint: write  $\hat{f} = f_2(x) - f_1(x)$  and show that

$$\Delta \hat{f} = \omega \hat{f}$$

where  $\omega := \int_0^1 W''((1-t)f_1(x) + tf_2(x))dt$ . is smooth.

(b) Consider  $u$  solving the Allen–Cahn equation on  $\mathbb{R}^n$  with  $\frac{\partial u}{\partial x^n} > 0$  and the condition from Theorem 32,  $\lim_{x^n \rightarrow \pm 1} u(x) = \pm 1$ . Show that  $u$  minimizes  $E_1(\cdot)$  on compact subsets of  $\mathbb{R}^n$ .

**5.2. Classifying stable entire solutions.** As such, we see that to solve De Giorgi’s conjecture (as well as to satisfy our original motivation: understanding stable solutions at scale  $O(\varepsilon)$ ) it makes sense to study entire stable solutions. The following results represent the state of affairs of the classification of stable solutions in  $\mathbb{R}^n$ .

**Theorem 33** (Ghoussoub–Gui [GG98]). *Consider  $u \in C^2(\mathbb{R}^2)$  a stable solution to the Allen–Cahn equation with  $|u| \leq 1$ . Then  $u(x) = \mathbb{H}(\langle a, x \rangle)$  is the 1-dimensional solution.*

See also [FMV13], who give a slightly different strategy of proof (this is the basis for the proof we give below).

**Theorem 34** (Ambrosio–Cabre [AC00]). *Consider  $u \in C^2(\mathbb{R}^3)$  a stable solution to the Allen–Cahn equation with  $|u| \leq 1$  and*

$$E_1(u; B_R) \leq CR^2$$

*for some  $C > 0$  independent of  $R$ . Then  $u(x) = \mathbb{H}(\langle a, x \rangle)$  is the 1-dimensional solution.*

**Theorem 35** (Pacard–Wei [PW13]). *For  $n \geq 8$ , there exists  $u \in C^\infty(\mathbb{R}^n)$  a stable solution to the Allen–Cahn equation with*

$$E_1(u; B_R) \leq CR^{n-1}$$

*but the level sets of  $u$  are not flat.*

Liu–Wang–Wei [LWW17] have recently extended this result to construct *minimizers* in  $\mathbb{R}^n$  for  $n \geq 8$ .

**5.3. Stable solutions in  $\mathbb{R}^2$ .** We prove Theorem 33. The beginning of the argument will work in all dimensions. We will indicate where we specialize to  $n = 2$  below. Assume that  $u$  is a stable solution to Allen–Cahn on  $\mathbb{R}^n$  with  $|u| \leq 1$ . By Exercise 15, stability implies that we have

$$\int_{\Sigma} |\nabla \varphi|^2 |\nabla u|^2 \geq \int_{\Sigma} (|D^2 u|^2 - |\nabla |\nabla u||^2) \varphi^2$$

for any compactly supported smooth function  $\varphi$ .

- Exercise 22 (+).** (a) Show that  $|D^2u|^2 - |\nabla|\nabla u||^2 \geq 0$ .
- (b) Suppose that  $|D^2u|^2 - |\nabla|\nabla u||^2 \equiv 0$  on  $\mathbb{R}^n$ . Show that  $\frac{\nabla u}{|\nabla u|}$  is a parallel vector field on  $\mathbb{R}^n$ . Hint: Compute  $|D(\nabla u/|\nabla u|)|^2$ .
- (c) Assumptions as in (b). Show that  $u(x) = \mathbb{H}(\langle a, x \rangle)$  for some  $|a| = 1$ .
- (d) The quantity  $|D^2u|^2 - |\nabla|\nabla u||^2$  is often thought of as the (square) norm of the “second fundamental form” of  $u$ . Justify this heuristic. Thus, this problem shows that the 1-dimensional solution is the unique solution on  $\mathbb{R}^n$  with vanishing second fundamental form.

**Exercise 23 (+).** Using interior Schauder estimates and  $|u| \leq 1$ , prove that  $|\nabla u| \leq C$  on  $\mathbb{R}^n$ .

Now, we specialize to  $n = 2$ . We would like to choose cutoff functions  $\varphi_i$  that tends to 1 pointwise on  $\mathbb{R}^2$  and so that

$$\int_{\mathbb{R}^2} |\nabla \varphi_i|^2 \rightarrow 0.$$

The function  $\psi_R$  cutting linearly (perhaps with a bit of smoothing) off between  $R$  and  $2R$  only gives

$$\int_{\mathbb{R}^2} |\nabla \psi_R|^2 \lesssim R^{-2} R^2 \leq C,$$

which is just barely failing what we want. It turns out that the solution is to use the *log-cutoff trick* which appears all over the place in similar problems (e.g. stable minimal surfaces in  $\mathbb{R}^3$ ). Motivated by the fundamental solution to the Laplacian on  $\mathbb{R}^2$ , we set

$$\varphi_R(x) := \begin{cases} 1 & |x| \leq R \\ 2 - \frac{\log|x|}{\log R} & R < |x| < R^2 \\ 0 & |x| \geq R^2 \end{cases}$$

for  $R > 1$ . As usual,  $\varphi$  is only Lipschitz, but we can justify plugging it into the stability inequality by an approximating argument. We thus find that

$$\int_{\mathbb{R}^2} |\nabla \varphi_R|^2 = \int_{B_{R^2}(0) \setminus B_R(0)} \frac{1}{|x|^2 \log^2 R} \lesssim \frac{1}{\log^2 R} \int_R^{R^2} r^{-1} dr = \frac{\log R}{\log^2 R} = \frac{1}{\log R} \rightarrow 0$$

as  $R \rightarrow \infty$ . We now use the previous two exercises: because  $|\nabla u| \leq C$  we get that

$$\liminf_{R \rightarrow \infty} \int_{\mathbb{R}^2} |\nabla \varphi_R|^2 |\nabla u|^2 \leq C^2 \liminf_{R \rightarrow \infty} \int_{\mathbb{R}^2} |\nabla \varphi_R|^2 = 0.$$

Moreover, combining  $\varphi \rightarrow 1$  pointwise and the fact that the right hand side of the stability inequality is non-negative, Fatou’s lemma implies that

$$\int_{\mathbb{R}^2} (|D^2u|^2 - |\nabla|\nabla u||^2) \leq 0.$$

Thus  $|D^2u|^2 - |\nabla|\nabla u||^2 = 0$ , so the proof is finished.

**5.4. Stable solutions in  $\mathbb{R}^3$ .** The strategy we used before has no hope of working if we try to repeat the steps verbatim.

**Exercise 24.** Show that if  $\varphi \equiv 1$  on  $B_R(0) \subset \mathbb{R}^3$  and  $\varphi$  has compact support, then  $\int_{\mathbb{R}^2} |\nabla\varphi|^2 \geq 4\pi R$ .

However, under the energy growth assumption  $E_1(u; B_R) \leq CR^2$ , we can do better by not using  $|\nabla u| \leq C$ . Using  $\varphi_R$  (the log-cutoff function), we find

$$\int_{\mathbb{R}^3} (|D^2u|^2 - |\nabla|\nabla u||^2)\varphi_R^2 \leq \int_{\mathbb{R}^3} |\nabla\varphi_R|^2 |\nabla u|^2 = \frac{1}{\log^2 R} \int_{B_{R^2}(0) \setminus B_R(0)} |x|^{-2} |\nabla u|^2.$$

For simplicity, assume that  $R = 2^k$  for some  $k = \frac{\log R}{\log 2} \in \mathbb{N}$ . Then, write

$$R_0 = 2^k, R_1 = 2^{k+1}, \dots, R_k = 2^{2k} = R^2.$$

so

$$\begin{aligned} \int_{B_{R^2}(0) \setminus B_R(0)} |x|^{-2} |\nabla u|^2 &= \sum_{j=0}^{k-1} \int_{B_{R_{j+1}} \setminus B_{R_j}} |x|^{-2} |\nabla u|^2 \\ &\leq \sum_{j=0}^{k-1} R_j^{-2} \int_{B_{R_{j+1}} \setminus B_{R_j}} |\nabla u|^2 \\ &\leq \sum_{j=0}^{k-1} R_j^{-2} \int_{B_{R_{j+1}}} |\nabla u|^2 \\ &\leq \sum_{j=0}^{k-1} R_j^{-2} R_{j+1}^2 \\ &\leq \sum_{j=0}^{k-1} 4 \\ &= 4k \\ &= \frac{4 \log R}{\log 2}. \end{aligned}$$

Thus, the remaining  $\log R$  in the denominator saves us (as before), so we find that

$$\int_{\mathbb{R}^3} |\nabla\varphi_R|^2 |\nabla u|^2 \rightarrow 0.$$

The proof is then completed as for  $n = 2$ .

**5.5. Area growth of monotone solutions in  $\mathbb{R}^3$ .** Finally, because we have developed most of the tools, we present the proof of De Giorgi's conjecture in  $\mathbb{R}^3$  by Ambrosio–Cabre (Theorem 30). Note that in  $n = 2$ , because monotone solutions are stable, the  $n = 2$  classification of stable solutions of Ghoussoub–Gui (Theorem 33) automatically resolves the problem. In  $\mathbb{R}^3$ , to apply the classification of stable solutions we must verify that monotone solutions have the quadratic area growth  $E_1(u; B_R) \leq CR^2$ .

Define  $u^t(x) = u(x_1, x_2, x_3 + t)$ . By monotonicity,

$$u^{\pm\infty}(x) := \lim_{t \rightarrow \pm\infty} u^t(x)$$

exists and is independent of  $x^3$ . Moreover, by using Schauder estimates, we see that the limit occurs smoothly on compact subsets of  $\mathbb{R}^3$ .

**Exercise 25 (+).** Show that if we drop  $x^3$ , then  $u^{\pm\infty}(x_1, x_2)$  is a stable solution to Allen–Cahn on  $\mathbb{R}^2$ . Thus, the classification of stable solutions on  $\mathbb{R}^2$  shows they are 1-dimensional. Use this to show that (as functions on  $\mathbb{R}^3$ ) we have

$$E_1(u^{\pm\infty}; B_R \subset \mathbb{R}^3) \leq CR^2$$

for some  $C > 0$  independent of  $R$ . Thus, conclude that

$$\lim_{t \rightarrow \pm\infty} E_1(u^t; B_R \subset \mathbb{R}^3) \leq CR^2$$

for some  $C > 0$  independent of  $R$ .

Note that because  $u$  is monotone,  $\partial_t u^t > 0$ . Now, consider

$$E_1(u^t; B_R) := \int_{B_R} \frac{1}{2} |\nabla u^t|^2 + W(u^t)$$

The idea is to differentiate this with respect to  $t$  and use the information just gained as  $t \rightarrow \infty$ . Recall that  $|\nabla u^t| \leq C$ . We now compute:

$$\begin{aligned} \partial_t E_1(u^t; B_R) &= \int_{B_R} \langle \nabla \partial_t u^t, \nabla u^t \rangle + W'(u^t) \partial_t u^t \\ &= \int_{B_R} -(\partial^t u^t) \Delta u^t + W'(u^t) \partial_t u^t + \int_{\partial B_R} \partial_t u^t \partial_\nu u^t \\ &= \int_{\partial B_R} \partial_t u^t \partial_\nu u^t \\ &\geq -C \int_{\partial B_R} \partial_t u^t. \end{aligned}$$

In the final inequality we crucially used (again) the monotonicity property of  $u$ .

Thus, integrating this with respect to  $t$ , we find that

$$E_1(u^{+\infty}; B_R) - E_1(u; B_R) \geq -C \int_{\partial B_R} (u^{+\infty} - u) \geq -CR^2.$$

In the final inequality, we used  $|u| \leq 1$  and  $|\partial B_R| = 4\pi R^2$ . Putting this together, we find that

$$E_1(u; B_R) \leq CR^2.$$

This completes the proof.

## 6. FURTHER READING

We give (a non-exhaustive) list of some references (in addition to those given above):

- Modica inequality and a monotonicity formula (an entire solution on  $\mathbb{R}^n$  satisfies the Modica inequality  $|\nabla u|^2 \leq 2W(u)$  with equality only for the 1-dimensional solution; this leads to a monotonicity formula that is fundamental for geometric applications): [Mod85, Ilm93, HT00]
- De Giorgi conjecture and classification stable solutions (besides those discussed above, there are several other related results): [GG03, JM04, FS17]
- Gibbons conjecture (the De Giorgi conjecture with a stronger condition as  $x^n \rightarrow \pm\infty$  is proven in all dimensions): [Far99, BBG00, BHM00]
- Existence/classification of solutions on  $\mathbb{R}^2$  (our understanding of entire solutions is best in dimension 2; certain uniqueness questions are still open): [KL11, Gui12, KLP13, dPKP13, KLPW15, GLW16, Wan17b, WW17]
- Other entire solutions in  $\mathbb{R}^n$  (in higher dimensions there are many interesting entire solutions; very few classification results are known): [dP10, dPMP12, dPKW13, AdPW15]
- The Toda system and the interaction between interfaces (a surprising feature that we did not discuss is the the interaction between level sets of solutions to the Allen–Cahn equation; this interaction is governed by a nonlinear system of PDE’s known as the Toda system): [Kow05, dPKW08, dPKWY10, WW17]
- The Allen–Cahn equation on manifolds (further existence and qualitative results not discussed above): [Man17, GG18b]

## REFERENCES

- [AC79] Samuel M Allen and John W Cahn, *A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening*, *Acta Metallurgica* **27** (1979), no. 6, 1085–1095.
- [AC00] Luigi Ambrosio and Xavier Cabré, *Entire solutions of semilinear elliptic equations in  $\mathbf{R}^3$  and a conjecture of De Giorgi*, *J. Amer. Math. Soc.* **13** (2000), no. 4, 725–739. MR 1775735
- [AdPW15] Oscar Agudelo, Manuel del Pino, and Juncheng Wei, *Solutions with multiple catenoidal ends to the Allen–Cahn equation in  $\mathbf{R}^3$* , *J. Math. Pures Appl.* (9) **103** (2015), no. 1, 142–218. MR 3281950

- [AFP00] Luigi Ambrosio, Nicola Fusco, and Diego Pallara, *Functions of bounded variation and free discontinuity problems*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 2000. MR 1857292
- [Alm66] F. J. Almgren, Jr., *Some interior regularity theorems for minimal surfaces and an extension of Bernstein's theorem*, Ann. of Math. (2) **84** (1966), 277–292. MR 0200816
- [BBG00] Martin T. Barlow, Richard F. Bass, and Changfeng Gui, *The Liouville property and a conjecture of De Giorgi*, Comm. Pure Appl. Math. **53** (2000), no. 8, 1007–1038. MR 1755949
- [BDGG69] E. Bombieri, E. De Giorgi, and E. Giusti, *Minimal cones and the Bernstein problem*, Invent. Math. **7** (1969), 243–268. MR 0250205
- [Ber27] Serge Bernstein, *über ein geometrisches Theorem und seine Anwendung auf die partiellen Differentialgleichungen vom elliptischen Typus*, Math. Z. **26** (1927), no. 1, 551–558. MR 1544873
- [BHM00] H. Berestycki, F. Hamel, and R. Monneau, *One-dimensional symmetry of bounded entire solutions of some elliptic equations*, Duke Math. J. **103** (2000), no. 3, 375–396. MR 1763653
- [CH58] John W Cahn and John E Hilliard, *Free energy of a nonuniform system. i. interfacial free energy*, The Journal of chemical physics **28** (1958), no. 2, 258–267.
- [CM18] Otis Chodosh and Christos Mantoulidis, *Minimal surfaces and the Allen-Cahn equation on 3-manifolds: index, multiplicity, and curvature estimates*, <https://arxiv.org/abs/1803.02716> (2018).
- [DFP92] Ha Dang, Paul C. Fife, and L. A. Peletier, *Saddle solutions of the bistable diffusion equation*, Z. Angew. Math. Phys. **43** (1992), no. 6, 984–998. MR 1198672
- [DG65] Ennio De Giorgi, *Una estensione del teorema di Bernstein*, Ann. Scuola Norm. Sup. Pisa (3) **19** (1965), 79–85. MR 0178385
- [DG79] ———, *Convergence problems for functionals and operators*, Proceedings of the International Meeting on Recent Methods in Nonlinear Analysis (Rome, 1978), Pitagora, Bologna, 1979, pp. 131–188. MR 533166
- [DGF75] Ennio De Giorgi and Tullio Franzoni, *Su un tipo di convergenza variazionale*, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8) **58** (1975), no. 6, 842–850. MR 0448194
- [dP10] Manuel del Pino, *Minimal surfaces and entire solutions of the Allen-Cahn equation*, <http://math.stanford.edu/~ryzhik/BANFF/delpino.pdf> (2010).
- [dPKP13] Manuel del Pino, MichałKowalczyk, and Frank Pacard, *Moduli space theory for the Allen-Cahn equation in the plane*, Trans. Amer. Math. Soc. **365** (2013), no. 2, 721–766. MR 2995371
- [dPKW08] Manuel del Pino, MichałKowalczyk, and Juncheng Wei, *The Toda system and clustering interfaces in the Allen-Cahn equation*, Arch. Ration. Mech. Anal. **190** (2008), no. 1, 141–187. MR 2434903
- [dPKW11] Manuel del Pino, Michał Kowalczyk, and Juncheng Wei, *On De Giorgi's conjecture in dimension  $N \geq 9$* , Ann. of Math. (2) **174** (2011), no. 3, 1485–1569. MR 2846486
- [dPKW13] ———, *Entire solutions of the Allen-Cahn equation and complete embedded minimal surfaces of finite total curvature in  $\mathbb{R}^3$* , J. Differential Geom. **93** (2013), no. 1, 67–131. MR 3019512
- [dPKWY10] Manuel del Pino, Michał Kowalczyk, Juncheng Wei, and Jun Yang, *Interface foliation near minimal submanifolds in Riemannian manifolds with positive Ricci curvature*, Geom. Funct. Anal. **20** (2010), no. 4, 918–957. MR 2729281

- [dPMP12] Manuel del Pino, Monica Musso, and Frank Pacard, *Solutions of the Allen-Cahn equation which are invariant under screw-motion*, Manuscripta Math. **138** (2012), no. 3-4, 273–286. MR 2916313
- [Far99] Alberto Farina, *Symmetry for solutions of semilinear elliptic equations in  $\mathbf{R}^N$  and related conjectures*, Ricerche Mat. **48** (1999), no. suppl., 129–154, Papers in memory of Ennio De Giorgi (Italian). MR 1765681
- [Fle62] Wendell H. Fleming, *On the oriented Plateau problem*, Rend. Circ. Mat. Palermo (2) **11** (1962), 69–90. MR 0157263
- [FMV13] Alberto Farina, Luciano Mari, and Enrico Valdinoci, *Splitting theorems, symmetry results and overdetermined problems for Riemannian manifolds*, Comm. Partial Differential Equations **38** (2013), no. 10, 1818–1862. MR 3169764
- [FS17] Alessio Figalli and Joaquim Serra, *On stable solutions for boundary reactions: a De Giorgi-type result in dimension  $4+1$* , <https://arxiv.org/abs/1705.02781> (2017).
- [FT89] Irene Fonseca and Luc Tartar, *The gradient theory of phase transitions for systems with two potential wells*, Proc. Roy. Soc. Edinburgh Sect. A **111** (1989), no. 1-2, 89–102. MR 985992
- [Gas17] Pedro Gaspar, *The second inner variation of energy and the Morse index of limit interfaces*, <https://arxiv.org/abs/1710.04719> (2017).
- [GG98] N. Ghoussoub and C. Gui, *On a conjecture of De Giorgi and some related problems*, Math. Ann. **311** (1998), no. 3, 481–491. MR 1637919
- [GG03] Nassif Ghoussoub and Changfeng Gui, *On De Giorgi’s conjecture in dimensions 4 and 5*, Ann. of Math. (2) **157** (2003), no. 1, 313–334. MR 1954269
- [GG18a] Pedro Gaspar and Marco A. M. Guaraco, *The Allen–Cahn equation on closed manifolds*, Calc. Var. Partial Differential Equations **57** (2018), no. 4, 57:101. MR 3814054
- [GG18b] Marco A. M. Guaraco and Pedro Gaspar, *The Weyl Law for the phase transition spectrum and density of limit interfaces*, <https://arxiv.org/abs/1804.04243> (2018).
- [Giu84] Enrico Giusti, *Minimal surfaces and functions of bounded variation*, Monographs in Mathematics, vol. 80, Birkhäuser Verlag, Basel, 1984. MR 775682
- [GLW16] Changfeng Gui, Yong Liu, and Juncheng Wei, *On variational characterization of four-end solutions of the Allen-Cahn equation in the plane*, J. Funct. Anal. **271** (2016), no. 10, 2673–2700. MR 3548276
- [GT01] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, Reprint of the 1998 edition. MR 1814364
- [Gua18] Marco A. M. Guaraco, *Min–max for phase transitions and the existence of embedded minimal hypersurfaces*, J. Differential Geom. **108** (2018), no. 1, 91–133. MR 3743704
- [Gui12] Changfeng Gui, *Symmetry of some entire solutions to the Allen-Cahn equation in two dimensions*, J. Differential Equations **252** (2012), no. 11, 5853–5874. MR 2911416
- [Hie17] Fritz Hiesmayr, *Spectrum and index of two-sided Allen-Cahn minimal hypersurfaces*, <https://arxiv.org/abs/1704.07738> (2017).
- [HT00] John E. Hutchinson and Yoshihiro Tonegawa, *Convergence of phase interfaces in the van der Waals-Cahn-Hilliard theory*, Calc. Var. Partial Differential Equations **10** (2000), no. 1, 49–84. MR 1803974

- [Ilm93] Tom Ilmanen, *Convergence of the Allen-Cahn equation to Brakke's motion by mean curvature*, J. Differential Geom. **38** (1993), no. 2, 417–461. MR 1237490
- [IMN18] Kei Irie, Fernando Marques, and André Neves, *Density of minimal hypersurfaces for generic metrics*, Ann. of Math. (2) **187** (2018), no. 3, 963–972. MR 3779962
- [JM04] David Jerison and Régis Monneau, *Towards a counter-example to a conjecture of De Giorgi in high dimensions*, Ann. Mat. Pura Appl. (4) **183** (2004), no. 4, 439–467. MR 2140525
- [JS09] Robert L. Jerrard and Peter Sternberg, *Critical points via  $\Gamma$ -convergence: general theory and applications*, J. Eur. Math. Soc. (JEMS) **11** (2009), no. 4, 705–753. MR 2538502
- [KL11] Michał Kowalczyk and Yong Liu, *Nondegeneracy of the saddle solution of the Allen-Cahn equation*, Proc. Amer. Math. Soc. **139** (2011), no. 12, 4319–4329. MR 2823077
- [KLP12] Michał Kowalczyk, Yong Liu, and Frank Pacard, *The space of 4-ended solutions to the Allen-Cahn equation in the plane*, Ann. Inst. H. Poincaré Anal. Non Linéaire **29** (2012), no. 5, 761–781. MR 2971030
- [KLP13] Michał Kowalczyk, Yong Liu, and Frank Pacard, *The classification of four-end solutions to the Allen-Cahn equation on the plane*, Anal. PDE **6** (2013), no. 7, 1675–1718. MR 3148064
- [KLPW15] Michał Kowalczyk, Yong Liu, Frank Pacard, and Juncheng Wei, *End-to-end construction for the Allen-Cahn equation in the plane*, Calc. Var. Partial Differential Equations **52** (2015), no. 1-2, 281–302. MR 3299182
- [Kow05] Michał Kowalczyk, *On the existence and Morse index of solutions to the Allen-Cahn equation in two dimensions*, Ann. Mat. Pura Appl. (4) **184** (2005), no. 1, 17–52. MR 2128093
- [KS89] Robert V. Kohn and Peter Sternberg, *Local minimisers and singular perturbations*, Proc. Roy. Soc. Edinburgh Sect. A **111** (1989), no. 1-2, 69–84. MR 985990
- [Le11] Nam Q. Le, *On the second inner variation of the Allen-Cahn functional and its applications*, Indiana Univ. Math. J. **60** (2011), no. 6, 1843–1856. MR 3008253
- [LMN18] Yevgeny Liokumovich, Fernando Marques, and André Neves, *Weyl law for the volume spectrum*, Ann. of Math. (2) **187** (2018), no. 3, 933–961. MR 3779961
- [LWW17] Yong Liu, Kelei Wang, and Juncheng Wei, *Global minimizers of the Allen-Cahn equation in dimension  $n \geq 8$* , J. Math. Pures Appl. (9) **108** (2017), no. 6, 818–840. MR 3723158
- [Mag12] Francesco Maggi, *Sets of finite perimeter and geometric variational problems*, Cambridge Studies in Advanced Mathematics, vol. 135, Cambridge University Press, Cambridge, 2012, An introduction to geometric measure theory. MR 2976521
- [Man17] Christos Mantoulidis, *Allen-Cahn min-max on surfaces*, <https://arxiv.org/abs/1706.05946> (2017).
- [Mar14] Fernando C. Marques, *Minimal surfaces - variational theory and applications*, Proceedings of the International Congress of Mathematicians, <https://arxiv.org/abs/1409.7648> (2014).
- [MM77] Luciano Modica and Stefano Mortola, *Un esempio di  $\Gamma^-$ -convergenza*, Boll. Un. Mat. Ital. B (5) **14** (1977), no. 1, 285–299. MR 0445362
- [MN16] Fernando C. Marques and André Neves, *Morse index and multiplicity of min-max minimal hypersurfaces*, Camb. J. Math. **4** (2016), no. 4, 463–511. MR 3572636
- [MN17] ———, *Existence of infinitely many minimal hypersurfaces in positive Ricci curvature*, Invent. Math. **209** (2017), no. 2, 577–616. MR 3674223

- [MN18] ———, *Morse index of multiplicity one min-max minimal hypersurfaces*, <https://arxiv.org/abs/1803.04273> (2018).
- [Mod85] Luciano Modica, *A gradient bound and a Liouville theorem for nonlinear Poisson equations*, *Comm. Pure Appl. Math.* **38** (1985), no. 5, 679–684. MR 803255
- [MR10] Frank Morgan and Antonio Ros, *Stable constant-mean-curvature hypersurfaces are area minimizing in small  $L^1$  neighborhoods*, *Interfaces Free Bound.* **12** (2010), no. 2, 151–155. MR 2652015
- [Nev14] André Neves, *New applications of min-max theory*, *Proceedings of International Congress of Mathematics*, <https://arxiv.org/abs/1409.7537> (2014), 939–957.
- [Pac12] Frank Pacard, *The role of minimal surfaces in the study of the Allen-Cahn equation*, *Geometric analysis: partial differential equations and surfaces*, *Contemp. Math.*, vol. 570, Amer. Math. Soc., Providence, RI, 2012, pp. 137–163. MR 2963598
- [PR03] Frank Pacard and Manuel Ritoré, *From constant mean curvature hypersurfaces to the gradient theory of phase transitions*, *J. Differential Geom.* **64** (2003), no. 3, 359–423. MR 2032110
- [PW13] Frank Pacard and Juncheng Wei, *Stable solutions of the Allen-Cahn equation in dimension 8 and minimal cones*, *J. Funct. Anal.* **264** (2013), no. 5, 1131–1167. MR 3010017
- [Sav09] Ovidiu Savin, *Regularity of flat level sets in phase transitions*, *Ann. of Math. (2)* **169** (2009), no. 1, 41–78. MR 2480601
- [Sim68] James Simons, *Minimal varieties in riemannian manifolds*, *Ann. of Math. (2)* **88** (1968), 62–105. MR 0233295
- [Sim83] Leon Simon, *Lectures on geometric measure theory*, *Proceedings of the Centre for Mathematical Analysis, Australian National University*, vol. 3, Australian National University, Centre for Mathematical Analysis, Canberra, 1983. MR 756417
- [Son18] Antoine Song, *Existence of infinitely many minimal hypersurfaces in closed manifolds*, <https://arxiv.org/abs/1806.08816> (2018).
- [SS81] Richard Schoen and Leon Simon, *Regularity of stable minimal hypersurfaces*, *Comm. Pure Appl. Math.* **34** (1981), no. 6, 741–797. MR 634285
- [Ste88] Peter Sternberg, *The effect of a singular perturbation on nonconvex variational problems*, *Arch. Rational Mech. Anal.* **101** (1988), no. 3, 209–260. MR 930124
- [TW12] Yoshihiro Tonegawa and Neshan Wickramasekera, *Stable phase interfaces in the van der Waals–Cahn–Hilliard theory*, *J. Reine Angew. Math.* **668** (2012), 191–210. MR 2948876
- [vdW79] Johannes Diederik van der Waals, *The thermodynamic theory of capillarity under the hypothesis of a continuous variation of density*, *Journal of Statistical Physics* **20** (1979), no. 2, 200–244.
- [Wan17a] Kelei Wang, *A new proof of Savin’s theorem on Allen-Cahn equations*, *J. Eur. Math. Soc. (JEMS)* **19** (2017), no. 10, 2997–3051. MR 3713000
- [Wan17b] ———, *Some remarks on the structure of finite Morse index solutions to the Allen-Cahn equation in  $\mathbb{R}^2$* , *NoDEA Nonlinear Differential Equations Appl.* **24** (2017), no. 5, Art. 58, 17. MR 3690656
- [Whi91] Brian White, *The space of minimal submanifolds for varying Riemannian metrics*, *Indiana Univ. Math. J.* **40** (1991), no. 1, 161–200. MR 1101226
- [Whi94] ———, *A strong minimax property of nondegenerate minimal submanifolds*, *J. Reine Angew. Math.* **457** (1994), 203–218. MR 1305283
- [Whi17] ———, *On the bumpy metrics theorem for minimal submanifolds*, *Amer. J. Math.* **139** (2017), no. 4, 1149–1155. MR 3689325

- [Wic14] Neshan Wickramasekera, *A general regularity theory for stable codimension 1 integral varifolds*, Ann. of Math. (2) **179** (2014), no. 3, 843–1007. MR 3171756
- [WW17] Kelei Wang and Juncheng Wei, *Finite Morse index implies finite ends*, to appear in Comm. Pure Appl. Math., <https://arxiv.org/abs/1705.06831> (2017).
- [Yau82] Shing Tung Yau, *Problem section*, Seminar on Differential Geometry, Ann. of Math. Stud., vol. 102, Princeton Univ. Press, Princeton, N.J., 1982, pp. 669–706. MR 645762