

Program Number: 363.8**Day / Time:** Monday, Nov. 14, 9:45 AM - 10:00 AM

Complex patterns of motor cortex activity during reaches at different speeds

M.M.Churchland^{*}; K.V.Shenoy

1. Dept of Elec Eng, 2. Neurosci. Prog., Stanford Univ., Stanford, CA, USA,

Is neural activity in the motor cortices translated fairly directly into muscle activity, or does it embody a representation of kinematic parameters such as hand velocity? Prior work has addressed this question by varying posture, load or initial hand position. We chose a similar approach, but varied reach speed. Two monkeys were trained to reach to radially arranged targets while keeping within speed 'windows' instructed by target color. 'Fast' reaches (red targets) were near the limit of the monkeys' abilities (~70-150 cm/s peak speed depending on distance; ~100-200 ms duration). 'Moderate' reaches (green targets) were ~2/3 as fast (~40-105 cm/s; ~150-300 ms duration). For both reach speeds, EMG activity recorded from the major muscles of the arm/shoulder differed considerably from the profile of hand velocity. Hand velocity was single-peaked, roughly bell-shaped, and scaled in the expected manner for fast vs. moderate reaches. EMG patterns were more complex, often multi-peaked, and did not simply scale for fast vs. moderate reaches. We asked whether the temporal patterns of neural activity, recorded from motor and dorsal premotor cortex, more closely resembled those of hand velocity or of the muscles. We found that the response patterns of individual neurons were strikingly complex, often containing multiple activity peaks during the reach. The number, amplitude and phase of these peaks varied with reach direction, distance and speed, often in rather idiosyncratic ways. In particular, patterns of neural activity for the two reach speeds were rarely scaled versions of one another. Thus, neural responses were inconsistent with a code dominated by hand velocity signals. On the other hand, the response patterns of individual neurons typically did not match the activity of any recorded muscle. It may be that the responses of many neurons combine to produce the response of a given muscle, but the activity of individual muscles was clearly not represented by the activity of individual neurons.

Support Contributed By: Helen Hay Whitney, NIH, BWF, NSF, ONR, Sloan, Whitaker

Citation: M.M.Churchland, K.V.Shenoy. Complex patterns of motor cortex activity during reaches at different speeds. Program No. 363.8. *2005 Abstract Viewer/Itinerary Planner*. Washington, DC: Society for Neuroscience

2005 Copyright by the Society for Neuroscience all rights reserved. Permission to republish any abstract or part of an abstract in any form must be obtained in writing from the SfN Office prior to publication

Application Design and Programming Copyright ScholarOne, Inc. All Rights Reserved. Patent Pending.