


# Functions


What is a function?

# Functions, High-School Edition



$$f(x) = x^4 - 5x^2 + 4$$

$$f(x) = \frac{x^2 + 4x - 9}{x^2 + 10x + 21}$$



# Functions, High-School Edition

In high school, functions are usually given as objects of the form

$$f(x) = \frac{x^3 + 3x^2 + 15x + 7}{1 - x^{137}}$$

What does a function do?

It takes in as input a real number.

It outputs a real number

... except when there are vertical asymptotes or other discontinuities, in which case the function doesn't output anything.

# Functions, CS Edition

```
int flipUntil(int n) {  
    int numHeads = 0;  
    int numTries = 0;  
  
    while (numHeads < n) {  
        if (randomBoolean()) numHeads++;  
  
        numTries++;  
    }  
  
    return numTries;  
}
```



# Functions, CS Edition

In programming, functions

- might take in inputs,
- might return values,
- might have side effects,
- might never return anything,
- might crash, and
- might return different values when called multiple times.

# What's Common?


Although high-school math functions and CS functions are pretty different, they have two key aspects in common:

- They take in inputs.
- They produce outputs.

In math, we like to keep things easy, so that's pretty much how we're going to define a function.

## ***Rough Idea of a Function:***

A function is an object  $f$  that takes in an input and produces exactly one output.



(This is not a complete definition – we'll revisit this in a bit.)

# High School versus CS Functions



In high school, functions usually were given by a rule:


$$f(x) = 4x + 15$$

In CS, functions are usually given by code:

```
int factorial(int n) {  
    int result = 1;  
    for (int i = 1; i <= n; i++) {  
        result *= i;  
    }  
    return result;  
}
```

What sorts of functions are we going to allow from a mathematical perspective?





... but also ...

$$f(x) = x^2 + 3x - 15$$



$$f(n) = \begin{cases} -n/2 & \text{if } n \text{ is even} \\ (n+1)/2 & \text{otherwise} \end{cases}$$

Functions like these  
are called ***piecewise  
functions.***

To define a function, you will typically either

- draw a picture, or
- give a rule for determining the output.

In mathematics, functions are ***deterministic***.

That is, given the same input, a function must always produce the same output.

The following is a perfectly valid piece of C++ code, but it's not a valid function under our definition:

```
int randomNumber(int numOutcomes)
{   return rand() % numOutcomes; }
```

One Challenge

$$f(x) = x^2 + 2x + 5$$

$$f(x) = x^2 + 2x + 5$$

$$f(3) = 3^2 + 3 \cdot 2 + 5 = 20$$

$$f(x) = x^2 + 2x + 5$$

$$f(3) = 3^2 + 3 \cdot 2 + 5 = 20$$

$$f(0) = 0^2 + 0 \cdot 2 + 5 = 5$$


$$f(x) = x^2 + 2x + 5$$

$$f(3) = 3^2 + 3 \cdot 2 + 5 = 20$$

$$f(0) = 0^2 + 0 \cdot 2 + 5 = 5$$

$$f(\text{Pikachu}) = \dots ?$$






$$f(\text{Pikachu}) = \text{Flying Pikachu}$$

$$f(137) = \dots?$$

We need to make sure we can't apply functions to meaningless inputs.

# Domains and Codomains

- Every function  $f$  has two sets associated with it: its **domain** and its **codomain**.
- A function  $f$  can only be applied to elements of its domain. For any  $x$  in the domain,  $f(x)$  belongs to the codomain.



# Domains and Codomains

- Every function  $f$  has two sets associated with it: its **domain** and its **codomain**.
- A function  $f$  can only be applied to elements of its domain. For any  $x$  in the domain,  $f(x)$  belongs to the codomain.



The **domain** of this function is  $\mathbb{R}$ . Any real number can be provided as input.

The **codomain** of this function is  $\mathbb{R}$ . Everything produced is a real number, but not all real numbers can be produced.

```
double absoluteValueOf(double x) {  
    if (x >= 0) {  
        return x;  
    } else {  
        return -x;  
    }  
}
```

# Domains and Codomains

- If  $f$  is a function whose domain is  $A$  and whose codomain is  $B$ , we write  $f : A \rightarrow B$ .
- Think of this like a “function prototype” in C++.



# The Official Rules for Functions

Formally speaking, we say that  $f : A \rightarrow B$  if the following two rules hold.

First,  $f$  must obey its domain/codomain rules:

**$\forall a \in A. \exists b \in B. f(a) = b$**   
(*“Every input in  $A$  maps to some output in  $B$ .”*)

Second,  $f$  must be deterministic:

**$\forall a_1 \in A. \forall a_2 \in A. (a_1 = a_2 \rightarrow f(a_1) = f(a_2))$**   
(*“Equal inputs produce equal outputs.”*)

If you’re ever curious about whether something is a function, look back at these rules and check! For example:

Can a function have an empty domain?

Can a function with a nonempty domain have an empty codomain?

# Defining Functions

Typically, we specify a function by describing a rule that maps every element of the domain to some element of the codomain.

Examples:

- $f(n) = n + 1$ , where  $f : \mathbb{Z} \rightarrow \mathbb{Z}$
- $f(x) = \sin x$ , where  $f : \mathbb{R} \rightarrow \mathbb{R}$
- $f(x) = \lfloor x \rfloor$ , where  $f : \mathbb{R} \rightarrow \mathbb{Z}$

Notice that we're giving both a rule and the domain/codomain.

# Defining Functions

Typically, we specify a function by describing a rule that maps every element of the domain to some element of the codomain.

Examples:

- $f(n) = n + 1$ , where  $f : \mathbb{Z} \rightarrow \mathbb{Z}$
- $f(x) = \sin x$ , where  $f : \mathbb{R} \rightarrow \mathbb{R}$
- $f(x) = \lfloor x \rfloor$ , where  $f : \mathbb{R} \rightarrow \mathbb{Z}$

Notice that we're giving both a **rule** and the **domain/codomain**.



# Defining Functions

Typically, we specify a function by describing a rule that maps every element of the domain to some element of the codomain.

This is the ceiling function – the smallest integer greater than or equal to  $x$ . For example,  $\lceil 1 \rceil = 1$ ,  $\lceil 1.37 \rceil = 2$ , and  $\lceil 3.14 \rceil = 4$ .

Examples:


$$f(n) = n + 1, \text{ where } f: \mathbb{Z} \rightarrow \mathbb{Z}$$

$$f(x) = \sin x, \text{ where } f: \mathbb{R} \rightarrow \mathbb{R}$$


$$f(x) = \lceil x \rceil, \text{ where } f: \mathbb{R} \rightarrow \mathbb{Z}$$

Notice that we're giving both a rule and the domain/codomain.




# Is This a Function From $A$ to $B$ ?



# Is This a Function From $A$ to $B$ ?




# Is This a Function From $A$ to $B$ ?



# Combining Functions

***f : People → Places***

***g : Places → Prices***



*People*

*Places*

*Prices*


***h : People → Prices***

***h(x) = g(f(x))***

# Function Composition

Suppose that we have two functions  $f : A \rightarrow B$  and  $g : B \rightarrow C$ .

Notice that the codomain of  $f$  is the domain of  $g$ . This means that we can use outputs from  $f$  as inputs to  $g$ .



# Function Composition

- Suppose that we have two functions  $f : A \rightarrow B$  and  $g : B \rightarrow C$ .
- The **composition of  $f$  and  $g$** , denoted  **$g \circ f$** , is a function where
  - $g \circ f : A \rightarrow C$ , and
  - $(g \circ f)(x) = g(f(x))$ .
- A few things to notice:
  - The domain of  $g \circ f$  is the domain of  $f$ . Its codomain is the codomain of  $g$ .
  - Even though the composition is written  $g \circ f$ , when evaluating  $(g \circ f)(x)$ , the function  $f$  is evaluated first.

The name of the function is  $g \circ f$ . When we apply it to an input  $x$ , we write  $(g \circ f)(x)$ . I don't know why, but that's what we do.



**Time-Out for Announcements!**

# Problem Set One Feedback


- Hopefully you have all seen problem set 1 feedback.
- If you haven't already, please review the feedback we've left for you as soon as possible, as well as the solution set.
- We're happy to answer any questions about specific comments in office hours or on Campuswire.
- If you believe we've made a grading error, see the Regrade Policies handout for instructions on how to submit a regrade.

# Problem Set Three

- Problem Set Three is due on Thursday at 11:59pm.
- Play around with binary relations, functions, their properties, and their applications!
- As usual, ***feel free to ask questions!***
- Ask on Campuswire!
- Stop by office hours!
- Pseudobreak from psets next week.


Back to CS103!

# Special Types of Functions






- Mercury
- Venus
- Earth
- Mars
- Jupiter
- Saturn
- Uranus
- Neptune
- Pluto














# Injective Functions

A function  $f : A \rightarrow B$  is called **injective** (or **one-to-one**) if the following statement is true about  $f$ :

$$\forall a_1 \in A. \forall a_2 \in A. (a_1 \neq a_2 \rightarrow f(a_1) \neq f(a_2))$$

*(“If the inputs are different, the outputs are different.”)*

The following first-order definition is equivalent and is often useful in proofs.

$$\forall a_1 \in A. \forall a_2 \in A. (f(a_1) = f(a_2) \rightarrow a_1 = a_2)$$

*(“If the outputs are the same, the inputs are the same.”)*

A function with this property is called an **injection**.

How does this compare to our second rule for functions?

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:**

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:**

*How many of the following are correct ways of starting off this proof?*

Consider any  $n_1, n_2 \in \mathbb{N}$  where  $n_1 = n_2$ . We will prove that  $f(n_1) = f(n_2)$ .

Consider any  $n_1, n_2 \in \mathbb{N}$  where  $n_1 \neq n_2$ . We will prove that  $f(n_1) \neq f(n_2)$ .

Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ . We will prove that  $n_1 = n_2$ .

Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) \neq f(n_2)$ . We will prove that  $n_1 \neq n_2$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:**

What does it mean for the function  $f$  to be injective?



# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:**

What does it mean for the function  $f$  to be injective?

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( f(n_1) = f(n_2) \rightarrow n_1 = n_2 )$$

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( n_1 \neq n_2 \rightarrow f(n_1) \neq f(n_2) )$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:**

What does it mean for the function  $f$  to be injective?

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( f(n_1) = f(n_2) \rightarrow n_1 = n_2 )$$

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( n_1 \neq n_2 \rightarrow f(n_1) \neq f(n_2) )$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:**

What does it mean for the function  $f$  to be injective?

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( f(n_1) = f(n_2) \rightarrow n_1 = n_2 )$$

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( n_1 \neq n_2 \rightarrow f(n_1) \neq f(n_2) )$$

Therefore, we'll pick arbitrary  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ , then prove that  $n_1 = n_2$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:**

What does it mean for the function  $f$  to be injective?

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( f(n_1) = f(n_2) \rightarrow n_1 = n_2 )$$

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( n_1 \neq n_2 \rightarrow f(n_1) \neq f(n_2) )$$

Therefore, we'll pick arbitrary  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ , then prove that  $n_1 = n_2$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:**

What does it mean for the function  $f$  to be injective?

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( f(n_1) = f(n_2) \rightarrow n_1 = n_2 )$$

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( n_1 \neq n_2 \rightarrow f(n_1) \neq f(n_2) )$$

Therefore, we'll pick arbitrary  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ , then prove that  $n_1 = n_2$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:**

What does it mean for the function  $f$  to be injective?

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( f(n_1) = f(n_2) \rightarrow n_1 = n_2 )$$

$$\forall n_1 \in \mathbb{N}. \forall n_2 \in \mathbb{N}. ( n_1 \neq n_2 \rightarrow f(n_1) \neq f(n_2) )$$

Therefore, we'll pick arbitrary  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ , then prove that  $n_1 = n_2$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:** Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ . We will prove that  $n_1 = n_2$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:** Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ . We  
will prove that  $n_1 = n_2$ .

Since  $f(n_1) = f(n_2)$ , we see that

$$2n_1 + 7 = 2n_2 + 7.$$



# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:** Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ . We  
will prove that  $n_1 = n_2$ .

Since  $f(n_1) = f(n_2)$ , we see that

$$2n_1 + 7 = 2n_2 + 7.$$

This in turn means that

$$2n_1 = 2n_2$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:** Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ . We  
will prove that  $n_1 = n_2$ .

Since  $f(n_1) = f(n_2)$ , we see that

$$2n_1 + 7 = 2n_2 + 7.$$

This in turn means that

$$2n_1 = 2n_2$$

so  $n_1 = n_2$ , as required.

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:** Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ . We  
will prove that  $n_1 = n_2$ .

Since  $f(n_1) = f(n_2)$ , we see that

$$2n_1 + 7 = 2n_2 + 7.$$

This in turn means that

$$2n_1 = 2n_2$$

so  $n_1 = n_2$ , as required. ■

# Injective Functions

**Theorem:** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be defined as  $f(n) = 2n + 7$ .  
Then  $f$  is injective.

**Proof:** Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ . We  
will prove that  $n_1 = n_2$ .

*How many of the following are correct ways of starting off this proof?*

Consider any  $n_1, n_2 \in \mathbb{N}$  where  $n_1 = n_2$ . We will prove that  $f(n_1) = f(n_2)$ .

Consider any  $n_1, n_2 \in \mathbb{N}$  where  $n_1 \neq n_2$ . We will prove that  $f(n_1) \neq f(n_2)$ .

Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) = f(n_2)$ . We will prove that  $n_1 = n_2$ .

Consider any  $n_1, n_2 \in \mathbb{N}$  where  $f(n_1) \neq f(n_2)$ . We will prove that  $n_1 \neq n_2$ .

So  $n_1 = n_2$ , as required. ■

Good exercise: Repeat this proof using  
the other definition of injectivity!

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

*How many of the following are correct ways of starting off this proof?*

Assume for the sake of contradiction that  $f$  is not injective.

Assume for the sake of contradiction that there are integers  $x_1$  and  $x_2$  where  $f(x_1) = f(x_2)$  but  $x_1 \neq x_2$ .

Consider arbitrary integers  $x_1$  and  $x_2$  where  $x_1 \neq x_2$ . We will prove that  $f(x_1) = f(x_2)$ .

Consider arbitrary integers  $x_1$  and  $x_2$  where  $f(x_1) = f(x_2)$ . We will prove that  $x_1 \neq x_2$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?



# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

$$\neg \forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

$$\neg \forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$
$$\exists x_1 \in \mathbb{Z}. \neg \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

$$\neg \forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

$$\exists x_1 \in \mathbb{Z}. \neg \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

$$\exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. \neg (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

$$\neg \forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

$$\exists x_1 \in \mathbb{Z}. \neg \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

$$\exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. \neg (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

$$\exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge \neg (f(x_1) \neq f(x_2)))$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

$$\neg \forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

$$\exists x_1 \in \mathbb{Z}. \neg \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

$$\exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. \neg (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

$$\exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge \neg (f(x_1) \neq f(x_2)))$$

$$\exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge f(x_1) = f(x_2))$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

$$\begin{aligned} & \neg \forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ & \exists x_1 \in \mathbb{Z}. \neg \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ & \exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. \neg (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ & \exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge \neg (f(x_1) \neq f(x_2))) \\ & \exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge f(x_1) = f(x_2)) \end{aligned}$$

Therefore, we need to find  $x_1, x_2 \in \mathbb{Z}$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ . Can we do that?



# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

$$\begin{aligned} & \neg \forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ & \exists x_1 \in \mathbb{Z}. \neg \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ & \exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. \neg (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ & \exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge \neg (f(x_1) \neq f(x_2))) \\ & \exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge f(x_1) = f(x_2)) \end{aligned}$$

Therefore, we need to find  $x_1, x_2 \in \mathbb{Z}$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ . Can we do that?

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

$$\begin{aligned} & \neg \forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ & \exists x_1 \in \mathbb{Z}. \neg \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ & \exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. \neg (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ & \exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge \neg (f(x_1) \neq f(x_2))) \\ & \exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge f(x_1) = f(x_2)) \end{aligned}$$

Therefore, we need to find  $x_1, x_2 \in \mathbb{Z}$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ . Can we do that?

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:**

What does it mean for  $f$  to be injective?

$$\forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2))$$

What is the negation of this statement?

$$\begin{aligned} &\neg \forall x_1 \in \mathbb{Z}. \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ &\exists x_1 \in \mathbb{Z}. \neg \forall x_2 \in \mathbb{Z}. (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ &\exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. \neg (x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)) \\ &\exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge \neg (f(x_1) \neq f(x_2))) \\ &\exists x_1 \in \mathbb{Z}. \exists x_2 \in \mathbb{Z}. (x_1 \neq x_2 \wedge f(x_1) = f(x_2)) \end{aligned}$$

Therefore, we need to find  $x_1, x_2 \in \mathbb{Z}$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ . Can we do that?

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:** We will prove that there exist integers  $x_1$  and  $x_2$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:** We will prove that there exist integers  $x_1$  and  $x_2$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ .

Let  $x_1 = -1$  and  $x_2 = +1$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:** We will prove that there exist integers  $x_1$  and  $x_2$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ .

Let  $x_1 = -1$  and  $x_2 = +1$ .

$$f(x_1) = f(-1) = (-1)^4 = 1$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:** We will prove that there exist integers  $x_1$  and  $x_2$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ .

Let  $x_1 = -1$  and  $x_2 = +1$ .

$$f(x_1) = f(-1) = (-1)^4 = 1$$

and

$$f(x_2) = f(1) = 1^4 = 1$$

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:** We will prove that there exist integers  $x_1$  and  $x_2$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ .

Let  $x_1 = -1$  and  $x_2 = +1$ .

$$f(x_1) = f(-1) = (-1)^4 = 1$$

and

$$f(x_2) = f(1) = 1^4 = 1,$$

so  $f(x_1) = f(x_2)$  even though  $x_1 \neq x_2$ , as required.



# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:** We will prove that there exist integers  $x_1$  and  $x_2$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ .

Let  $x_1 = -1$  and  $x_2 = +1$ .

$$f(x_1) = f(-1) = (-1)^4 = 1$$

and

$$f(x_2) = f(1) = 1^4 = 1,$$

so  $f(x_1) = f(x_2)$  even though  $x_1 \neq x_2$ , as required. ■

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:** We will prove that there exist integers  $x_1$  and  $x_2$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ .

*How many of the following are correct ways of starting off this proof?*

Assume for the sake of contradiction that  $f$  is not injective.

Assume for the sake of contradiction that there are integers  $x_1$  and  $x_2$  where  $f(x_1) = f(x_2)$  but  $x_1 \neq x_2$ .

Consider arbitrary integers  $x_1$  and  $x_2$  where  $x_1 \neq x_2$ . We will prove that  $f(x_1) = f(x_2)$ .

Consider arbitrary integers  $x_1$  and  $x_2$  where  $f(x_1) = f(x_2)$ . We will prove that  $x_1 \neq x_2$ .

# Injective Functions

**Theorem:** Let  $f : \mathbb{Z} \rightarrow \mathbb{N}$  be defined as  $f(x) = x^4$ . Then  $f$  is not injective.

**Proof:** We will prove that there exist integers  $x_1$  and  $x_2$  such that  $x_1 \neq x_2$ , but  $f(x_1) = f(x_2)$ .

Let  $x_1 = -1$  and  $x_2 = +1$ .

$$f(x_1) = f(-1) = (-1)^4 = 1$$

and

$$f(x_2) = f(1) = 1^4 = 1,$$

so  $f(x_1) = f(x_2)$  even though  $x_1 \neq x_2$ , as required. ■

# Injections and Composition

# Injections and Composition

- ***Theorem:*** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is an injection.
- Our goal will be to prove this result. To do so, we're going to have to call back to the formal definitions of injectivity and function composition.

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:**

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections.



**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective.

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective.

There are two definitions of injectivity that we can use here:

$$\forall a_1 \in A. \forall a_2 \in A. ((g \circ f)(a_1) = (g \circ f)(a_2) \rightarrow a_1 = a_2)$$

$$\forall a_1 \in A. \forall a_2 \in A. (a_1 \neq a_2 \rightarrow (g \circ f)(a_1) \neq (g \circ f)(a_2))$$

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective.

There are two definitions of injectivity that we can use here:

$$\forall a_1 \in A. \forall a_2 \in A. ((g \circ f)(a_1) = (g \circ f)(a_2) \rightarrow a_1 = a_2)$$

$$\forall a_1 \in A. \forall a_2 \in A. (a_1 \neq a_2 \rightarrow (g \circ f)(a_1) \neq (g \circ f)(a_2))$$

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective.

There are two definitions of injectivity that we can use here:

$$\forall a_1 \in A. \forall a_2 \in A. ((g \circ f)(a_1) = (g \circ f)(a_2) \rightarrow a_1 = a_2)$$

$$\forall a_1 \in A. \forall a_2 \in A. (a_1 \neq a_2 \rightarrow (g \circ f)(a_1) \neq (g \circ f)(a_2))$$

Therefore, we'll choose an arbitrary  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ , then prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ .

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ .

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ .

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ .

How is  $(g \circ f)(x)$  defined?

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ .

How is  $(g \circ f)(x)$  defined?

$$(g \circ f)(x) = g(f(x))$$



**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ .

How is  $(g \circ f)(x)$  defined?

$$(g \circ f)(x) = g(f(x))$$

So we need to prove that  $g(f(a_1)) \neq g(f(a_2))$ .

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ . Equivalently, we need to show that  $g(f(a_1)) \neq g(f(a_2))$ .


How is  $(g \circ f)(x)$  defined?

$$(g \circ f)(x) = g(f(x))$$

So we need to prove that  $g(f(a_1)) \neq g(f(a_2))$ .

**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.


**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ . Equivalently, we need to show that  $g(f(a_1)) \neq g(f(a_2))$ .



**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ . Equivalently, we need to show that  $g(f(a_1)) \neq g(f(a_2))$ .


Since  $f$  is injective and  $a_1 \neq a_2$ , we see that  $f(a_1) \neq f(a_2)$ .



**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ . Equivalently, we need to show that  $g(f(a_1)) \neq g(f(a_2))$ .


Since  $f$  is injective and  $a_1 \neq a_2$ , we see that  $f(a_1) \neq f(a_2)$ . Then, since  $g$  is injective and  $f(a_1) \neq f(a_2)$ , we see that  $g(f(a_1)) \neq g(f(a_2))$ , as required.



**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ . Equivalently, we need to show that  $g(f(a_1)) \neq g(f(a_2))$ .


Since  $f$  is injective and  $a_1 \neq a_2$ , we see that  $f(a_1) \neq f(a_2)$ . Then, since  $g$  is injective and  $f(a_1) \neq f(a_2)$ , we see that  $g(f(a_1)) \neq g(f(a_2))$ , as required. ■



**Theorem:** If  $f : A \rightarrow B$  is an injection and  $g : B \rightarrow C$  is an injection, then the function  $g \circ f : A \rightarrow C$  is also an injection.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary injections. We will prove that the function  $g \circ f : A \rightarrow C$  is also injective. To do so, consider any  $a_1, a_2 \in A$  where  $a_1 \neq a_2$ . We will prove that  $(g \circ f)(a_1) \neq (g \circ f)(a_2)$ . Equivalently, we need to show that  $g(f(a_1)) \neq g(f(a_2))$ .


Since  $f$  is injective and  $a_1 \neq a_2$ , we see that  $f(a_1) \neq f(a_2)$ . Then, since  $g$  is injective and  $f(a_1) \neq f(a_2)$ , we see that  $g(f(a_1)) \neq g(f(a_2))$ , as required. ■



Let's take a five minute break!



# Another Class of Functions



# Surjective Functions

A function  $f : A \rightarrow B$  is called **surjective** (or **onto**) if this first-order logic statement is true about  $f$ :

$$\forall b \in B. \exists a \in A. f(a) = b$$

*(“For every possible output, there's at least one possible input that produces it”)*

A function with this property is called a **surjection**.

How does this compare to our first rule of functions?

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:**

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:**

What does it mean for  $f$  to be surjective?

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:**

What does it mean for  $f$  to be surjective?

$$\forall y \in \mathbb{R}. \exists x \in \mathbb{R}. f(x) = y$$

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:**

What does it mean for  $f$  to be surjective?

$$\forall y \in \mathbb{R}. \exists x \in \mathbb{R}. f(x) = y$$

Therefore, we'll choose an arbitrary  $y \in \mathbb{R}$ , then prove that there is some  $x \in \mathbb{R}$  where  $f(x) = y$ .



# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:**

What does it mean for  $f$  to be surjective?

$$\forall y \in \mathbb{R}. \exists x \in \mathbb{R}. f(x) = y$$

Therefore, we'll choose an arbitrary  $y \in \mathbb{R}$ , then prove that there is some  $x \in \mathbb{R}$  where  $f(x) = y$ .

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:**

What does it mean for  $f$  to be surjective?

$$\forall y \in \mathbb{R}. \exists x \in \mathbb{R}. f(x) = y$$

Therefore, we'll choose an arbitrary  $y \in \mathbb{R}$ , then prove that there is some  $x \in \mathbb{R}$  where  $f(x) = y$ .

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:**

What does it mean for  $f$  to be surjective?

$$\forall y \in \mathbb{R}. \exists x \in \mathbb{R}. f(x) = y$$

Therefore, we'll choose an arbitrary  $y \in \mathbb{R}$ , then prove that there is some  $x \in \mathbb{R}$  where  $f(x) = y$ .

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:** Consider any  $y \in \mathbb{R}$ .

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:** Consider any  $y \in \mathbb{R}$ . We will prove that there is a choice of  $x \in \mathbb{R}$  such that  $f(x) = y$ .

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:** Consider any  $y \in \mathbb{R}$ . We will prove that there is a choice of  $x \in \mathbb{R}$  such that  $f(x) = y$ .

Let  $x = 2y$ .

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:** Consider any  $y \in \mathbb{R}$ . We will prove that there is a choice of  $x \in \mathbb{R}$  such that  $f(x) = y$ .

Let  $x = 2y$ . Then we see that

$$f(x) = f(2y)$$

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:** Consider any  $y \in \mathbb{R}$ . We will prove that there is a choice of  $x \in \mathbb{R}$  such that  $f(x) = y$ .

Let  $x = 2y$ . Then we see that

$$f(x) = f(2y) = 2y / 2$$



# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:** Consider any  $y \in \mathbb{R}$ . We will prove that there is a choice of  $x \in \mathbb{R}$  such that  $f(x) = y$ .

Let  $x = 2y$ . Then we see that

$$f(x) = f(2y) = 2y / 2 = y.$$

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:** Consider any  $y \in \mathbb{R}$ . We will prove that there is a choice of  $x \in \mathbb{R}$  such that  $f(x) = y$ .

Let  $x = 2y$ . Then we see that

$$f(x) = f(2y) = 2y / 2 = y.$$

So  $f(x) = y$ , as required.

# Surjective Functions

**Theorem:** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = x / 2$ . Then  $f(x)$  is surjective.

**Proof:** Consider any  $y \in \mathbb{R}$ . We will prove that there is a choice of  $x \in \mathbb{R}$  such that  $f(x) = y$ .

Let  $x = 2y$ . Then we see that

$$f(x) = f(2y) = 2y / 2 = y.$$

So  $f(x) = y$ , as required. ■

# Composing Surjections

**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:**

**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections.

**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective.



**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective.

What does it mean for  $g \circ f : A \rightarrow C$  to be surjective?

**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective.

What does it mean for  $g \circ f : A \rightarrow C$  to be surjective?

$$\forall c \in C. \exists a \in A. (g \circ f)(a) = c$$

**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective.

What does it mean for  $g \circ f : A \rightarrow C$  to be surjective?

$$\forall c \in C. \exists a \in A. (g \circ f)(a) = c$$


Therefore, we'll choose arbitrary  $c \in C$  and prove that there is some  $a \in A$  such that  $(g \circ f)(a) = c$ .

**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective. To do so, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $(g \circ f)(a) = c$ .

**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.


**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective. To do so, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $(g \circ f)(a) = c$ . Equivalently, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $g(f(a)) = c$ .



**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective. To do so, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $(g \circ f)(a) = c$ . Equivalently, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $g(f(a)) = c$ .


Consider any  $c \in C$ .



**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective. To do so, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $(g \circ f)(a) = c$ . Equivalently, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $g(f(a)) = c$ .


Consider any  $c \in C$ . Since  $g : B \rightarrow C$  is surjective, there is some  $b \in B$  such that  $g(b) = c$ .



**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective. To do so, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $(g \circ f)(a) = c$ . Equivalently, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $g(f(a)) = c$ .

Consider any  $c \in C$ . Since  $g : B \rightarrow C$  is surjective, there is some  $b \in B$  such that  $g(b) = c$ . Similarly, since  $f : A \rightarrow B$  is surjective, there is some  $a \in A$  such that  $f(a) = b$ .






**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective. To do so, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $(g \circ f)(a) = c$ . Equivalently, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $g(f(a)) = c$ .

Consider any  $c \in C$ . Since  $g : B \rightarrow C$  is surjective, there is some  $b \in B$  such that  $g(b) = c$ . Similarly, since  $f : A \rightarrow B$  is surjective, there is some  $a \in A$  such that  $f(a) = b$ . This means that there is some  $a \in A$  such that

$$g(f(a)) = g(b) = c,$$

which is what we needed to show.




**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective. To do so, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $(g \circ f)(a) = c$ . Equivalently, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $g(f(a)) = c$ .

Consider any  $c \in C$ . Since  $g : B \rightarrow C$  is surjective, there is some  $b \in B$  such that  $g(b) = c$ . Similarly, since  $f : A \rightarrow B$  is surjective, there is some  $a \in A$  such that  $f(a) = b$ . This means that there is some  $a \in A$  such that

$$g(f(a)) = g(b) = c,$$

which is what we needed to show. ■



**Theorem:** If  $f : A \rightarrow B$  is surjective and  $g : B \rightarrow C$  is surjective, then  $g \circ f : A \rightarrow C$  is also surjective.

**Proof:** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be arbitrary surjections. We will prove that the function  $g \circ f : A \rightarrow C$  is also surjective. To do so, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $(g \circ f)(a) = c$ . Equivalently, we will prove that for any  $c \in C$ , there is some  $a \in A$  such that  $g(f(a)) = c$ .

Consider any  $c \in C$ . Since  $g : B \rightarrow C$  is surjective, there is some  $b \in B$  such that  $g(b) = c$ . Similarly, since  $f : A \rightarrow B$  is surjective, there is some  $a \in A$  such that  $f(a) = b$ . This means that there is some  $a \in A$  such that

$$g(f(a)) = g(b) = c,$$

which is what we needed to show. ■

# Injections and Surjections

An injective function associates *at most* one element of the domain with each element of the codomain.

A surjective function associates *at least* one element of the domain with each element of the codomain.


What about functions that associate *exactly one* element of the domain with each element of the codomain?



**Katniss Everdeen**



**Elsa**



**Hermione Granger**

# Bijections

A function that associates each element of the codomain with a unique element of the domain is called ***bijjective***.

Such a function is a ***bijection***.

Formally, a bijection is a function that is both *injective* and *surjective*.

Bijections are sometimes called ***one-to-one correspondences***.

Not to be confused with “one-to-one functions.”

# Bijections and Composition

Suppose that  $f : A \rightarrow B$  and  $g : B \rightarrow C$  are bijections.

Is  $g \circ f$  necessarily a bijection?

**Yes!**

Since both  $f$  and  $g$  are injective, we know that  $g \circ f$  is injective.

Since both  $f$  and  $g$  are surjective, we know that  $g \circ f$  is surjective.

Therefore,  $g \circ f$  is a bijection.

# Inverse Functions







**Katniss Everdeen**





**Elsa**








**Hermione Granger**
















# Inverse Functions

In some cases, it's possible to “turn a function around.”

Let  $f : A \rightarrow B$  be a function. A function  $f^{-1} : B \rightarrow A$  is called an **inverse of  $f$**  if the following first-order logic statements are true about  $f$  and  $f^{-1}$

$$\forall a \in A. (f^{-1}(f(a)) = a) \quad \forall b \in B. (f(f^{-1}(b)) = b)$$

In other words, if  $f$  maps  $a$  to  $b$ , then  $f^{-1}$  maps  $b$  back to  $a$  and vice-versa.

Not all functions have inverses (we just saw a few examples of functions with no inverses).

If  $f$  is a function that has an inverse, then we say that  $f$  is **invertible**.



# Inverse Functions

***Theorem:*** Let  $f : A \rightarrow B$ . Then  $f$  is invertible if and only if  $f$  is a bijection.

These proofs are in the course reader. Feel free to check them out if you'd like!

***Really cool observation:*** Look at the formal definition of a function. Look at the rules for injectivity and surjectivity. Do you see why this result makes sense?