

Welcome to CS103!


Are there “laws of physics”  
in computer science?

# Key Questions in CS103

- What problems can you solve with a computer?
  - *Computability Theory*
- Why are some problems harder to solve than others?
  - *Complexity Theory*
- How can we be certain in our answers to these questions?
  - *Discrete Mathematics*



***Keith Schwarz***  
(Instructor)



***Akankshita Dash***  
(TA)



***Caroline Gao***  
(TA)



***Goli Emami***  
(TA)



***Grant McClearn***  
(TA)



***Jennie Chung***  
(TA)



***Julie Liu***  
(TA)




***Lori Liu***  
(TA)



***Ophir Horovitz***  
(TA)



***Wanyue Zhai***  
(TA)



***Warren Xia***  
(TA)



***Zheng Lian***  
(Head TA)

***Keith's Email: [htiek@cs.stanford.edu](mailto:htiek@cs.stanford.edu)***  
***Staff Email List: [cs103-aut2223-staff@lists.stanford.edu](mailto:cs103-aut2223-staff@lists.stanford.edu)***

# Asking Questions

- We've set up an online system you can use to ask us questions in lecture.
- First, visit our EdStem page. You can use this link, get there through Canvas, or follow the link from the lecture page on the course website.

<https://edstem.org/us/courses/27833/discussion/>

- Next, find the pinned thread at the top entitled  
***L00: Introduction, Set Theory.***
- Once you've found that thread, give it a      to let us know you've found it.
- Feel free to post questions here during lecture. The TAs will answer in real time. (You can also raise your hand if you have a question, too!)

# Course Website

**<https://cs103.stanford.edu>**

All course content  
will be hosted  
here, except for  
lecture videos.

# Prerequisite / Corequisite

# CS106B

Some problem sets will have small coding components. We'll also reference some concepts from CS106B, particularly recursion, throughout the quarter.

There aren't any math prerequisites for this course  
– high-school algebra should be enough!

# Another Option

# CS154

CS154 is more appropriate if you have a background in the topics from the first half of this quarter (set theory, proofwriting, discrete math, formal logic, graphs, etc.)

Come talk to me after class if you're curious about this!

# CS103A

- **CS103A** is an optional, one-unit companion course to CS103.
- CS103A meets Tuesdays, 4:30PM – 5:50PM, to provide additional practice with the course material in a small group setting.
- The first course meeting is next week.
- Interested? Apply online using [this link](#).





***Pranav Sriram***  
(ACE Instructor)

# Problem Set 0

- Your first assignment, Problem Set 0, goes out today. It's due Friday at 2:30PM Pacific.
- This assignment requires you to set up your development environment and to get set up on GradeScope.
- There's no coding involved, but it's good to start early in case you encounter any technical issues.

# *Recommended* Reading



# Grading




■ Problem Sets


## ***Ten Problem Sets***

Problem sets may be completed individually or in pairs.

# Grading



# Grading



We've got a big journey ahead of us.

***Let's get started!***

# Introduction to Set Theory


“CS103 students”

“Cool people”

“The chemical elements”


“Cute animals”

“US coins”



---






A **set** is an unordered collection of distinct objects, which may be anything, including other sets.




These are two different descriptions of exactly the same set.

---

Two sets are equal when they have the same contents, ignoring order.



Sets cannot contain duplicate elements.  
Any repeated elements are ignored.



These are two different descriptions of exactly the same set.

---

Sets cannot contain duplicate elements.  
Any repeated elements are ignored.



This symbol means "is  
an element of."

---

The objects that make up a set are called the  
***elements*** of that set.



This symbol means "is not an element of."


---

The objects that make up a set are called the ***elements*** of that set.

$$\{\} = \emptyset$$

The *empty set* is the set with no elements.

We denote the empty set using this symbol.



---


Sets can contain any number of elements.

$$1 \neq \{1\}$$

1

This is a  
number.

This is a set.  
It contains a  
number.




**Question:** Are these objects equal?


$$\emptyset \neq \{\emptyset\}$$



This is the  
empty set.




This is a set  
with the empty  
set in it.



**Question:** Are these objects equal?

$$x \neq \{x\}$$



No object  $x$  is equal to the set containing  $x$ .

# Infinite Sets

- Some sets contain *infinitely many* elements!
- The set  $\mathbb{N} = \{ 0, 1, 2, 3, \dots \}$  is the set of all the ***natural numbers***.
  - Some mathematicians don't include zero; in this class, assume that 0 is a natural number.
- The set  $\mathbb{Z} = \{ \dots, -2, -1, 0, 1, 2, \dots \}$  is the set of all the ***integers***.
  - Z is from German “Zahlen.”
- The set  $\mathbb{R}$  is the set of all ***real numbers***.
  - $e \in \mathbb{R}$ ,  $\pi \in \mathbb{R}$ ,  $4 \in \mathbb{R}$ ,  $-137$ , etc.

# Describing Complex Sets

- Here are some English descriptions of infinite sets:
  - “The set of all even natural numbers.”
  - “The set of all real numbers less than 137.”
  - “The set of all negative integers.”
- To describe complex sets like these mathematically, we'll use ***set-builder notation***.

# Even Natural Numbers

$$\{ n \mid n \in \mathbb{N} \text{ and } n \text{ is even} \}$$

The set of all  $n$

where

$n$  is a natural  
number

and  $n$  is even

# Even Natural Numbers

$$\{ n \mid n \in \mathbb{N} \text{ and } n \text{ is even} \}$$

The set of all  $n$

where

$n$  is a natural  
number

and  $n$  is even

$$\{ 0, 2, 4, 6, 8, 10, 12, 14, 16, \dots \}$$

# Set Builder Notation

- A set may be specified in ***set-builder notation***:

$$\{ x \mid \text{some property } x \text{ satisfies} \}$$

$$\{ x \in S \mid \text{some property } x \text{ satisfies} \}$$

- For example:

$$\{ n \mid n \in \mathbb{N} \text{ and } n \text{ is even} \}$$


$$\{ C \mid C \text{ is a set of US currency} \}$$

$$\{ r \in \mathbb{R} \mid r < 3 \}$$

$$\{ n \in \mathbb{N} \mid n < 3 \} \quad (\text{the set } \{0, 1, 2\})$$

# Combining Sets

# Venn Diagrams




*A*

$$A = \{ 1, 2, 3 \}$$

$$B = \{ 3, 4, 5 \}$$

# Venn Diagrams




*B*

$$A = \{ 1, 2, 3 \}$$

$$B = \{ 3, 4, 5 \}$$


# Venn Diagrams



Union  
 $A \cup B$   
 $\{ 1, 2, 3, 4, 5 \}$

$$A = \{ 1, 2, 3 \}$$
$$B = \{ 3, 4, 5 \}$$

# Venn Diagrams



Intersection


$$A \cap B$$

$$\{ 3 \}$$

$$A = \{ 1, 2, 3 \}$$

$$B = \{ 3, 4, 5 \}$$

# Venn Diagrams



Difference


$$A - B$$

$$\{ 1, 2 \}$$

$$A = \{ 1, 2, 3 \}$$

$$B = \{ 3, 4, 5 \}$$

# Venn Diagrams




Difference

$$A \setminus B$$
$$\{ 1, 2 \}$$

$$A = \{ 1, 2, 3 \}$$

$$B = \{ 3, 4, 5 \}$$


# Venn Diagrams




Symmetric  
Difference  
 $A \Delta B$   
 $\{ 1, 2, 4, 5 \}$

$$A = \{ 1, 2, 3 \}$$
$$B = \{ 3, 4, 5 \}$$


# Venn Diagrams




$$A \Delta B$$




# Venn Diagrams



# Venn Diagrams for Three Sets




# Venn Diagrams for Four Sets



Question to ponder:  
why don't we just  
draw four circles?

# Venn Diagrams for Five Sets



# Venn Diagrams for Seven Sets

**<http://moebio.com/research/sevensets/>**


# Subsets and Power Sets

# Subsets

- A set  $S$  is called a **subset** of a set  $T$  (denoted  **$S \subseteq T$** ) if all elements of  $S$  are also elements of  $T$ .
- Examples:
  - $\{ 1, 2, 3 \} \subseteq \{ 1, 2, 3, 4 \}$
  - $\{ b, c \} \subseteq \{ a, b, c, d \}$
  - $\{ \text{H}, \text{He}, \text{Li} \} \subseteq \{ \text{H}, \text{He}, \text{Li} \}$
  - $\mathbb{N} \subseteq \mathbb{Z}$  (*every natural number is an integer*)
  - $\mathbb{Z} \subseteq \mathbb{R}$  (*every integer is a real number*)

# Subsets and Elements


Set  $S$



$$S = \{ 2, \star, \{2\}, \text{smiley face}, \text{penny} \}$$

# Subsets and Elements


Set  $S$



$\star \in S$

# Subsets and Elements

Set  $S$




2

$2 \in S$

# Subsets and Elements

Set  $S$




General intuition:  
 $x \in S$  means you  
can ***point at  $x$***   
***inside of  $S$*** .

$$\{2\} \in S$$

# Subsets and Elements


Set  $S$



$$\left\{ \text{coin}, 2 \right\} \subseteq S$$

# Subsets and Elements


Set  $S$



$\{ \text{coin}, 2 \} \notin S$

# Subsets and Elements


Set  $S$



$\{2\} \in S$

# Subsets and Elements

Set  $S$




General intuition:  
 $A \subseteq B$  if you can  
form  $A$  by ***circling  
elements of  $B$ .***

$$\{2\} \subseteq S$$

# Subsets and Elements

Set  $S$




$$2 \notin S$$

(Since 2  
isn't a set.)

# Subsets and Elements


Set  $S$



$$\emptyset \subseteq S$$

# Subsets and Elements

Set  $S$



# Subsets and Elements

- We say that  $S \in T$  if, among the elements of  $T$ , one of them is *exactly* the object  $S$ .
- We say that  $S \subseteq T$  if  $S$  is a set and every element of  $S$  is also an element of  $T$ . ( $S$  has to be a set for the statement  $S \subseteq T$  to be true.)
- Although these concepts are similar, ***they are not the same!*** Not all elements of a set are subsets of that set and vice-versa.
- We have a resource on the course website, the Guide to Elements and Subsets, that explores this in more depth.

$$S = \left\{ \text{Lincoln Penny}, \text{Lincoln Dime} \right\}$$

$$\wp(S) = \left\{ \emptyset, \{ \text{Lincoln Dime} \}, \{ \text{Lincoln Penny} \}, \{ \text{Lincoln Penny}, \text{Lincoln Dime} \} \right\}$$

This is the **power set** of  $S$ , the set of all subsets of  $S$ . We write the power set of  $S$  as  $\wp(S)$ .

Formally,  $\wp(S) = \{ T \mid T \subseteq S \}$ .  
*(Do you see why?)*

What is  $\wp(\emptyset)$ ?

**Answer:**  $\{\emptyset\}$

*Remember that  $\emptyset \neq \{\emptyset\}$ !*

# Cardinality

# Cardinality

- The **cardinality** of a set is the number of elements it contains.
- If  $S$  is a set, we denote its cardinality as  $|S|$ .
- Examples:
  - $|\{\textit{whimsy}, \textit{mirth}\}| = 2$
  - $|\{\{a, b\}, \{c, d, e, f, g\}, \{h\}\}| = 3$
  - $|\{1, 2, 3, 3, 3, 3, 3\}| = 3$
  - $|\{n \in \mathbb{N} \mid n < 4\}| = |\{0, 1, 2, 3\}| = 4$
  - $|\emptyset| = 0$
  - $|\{\emptyset\}| = 1$

# The Cardinality of $\mathbb{N}$


- What is  $|\mathbb{N}|$ ?
  - There are infinitely many natural numbers.
  - $|\mathbb{N}|$  can't be a natural number, since it's infinitely large.
- We need to introduce a new term.
- Let's define  $\aleph_0 = |\mathbb{N}|$ .
  - $\aleph_0$  is pronounced “aleph-zero,” “aleph-nought,” or “aleph-null.”

Consider the set

$$S = \{ n \mid n \in \mathbb{N} \text{ and } n \text{ is even} \}.$$


What is  $|S|$ ?

# How Big Are These Sets?




# Comparing Cardinalities

- Two sets have the same cardinality if there is a way to pair their elements off without leaving any elements uncovered.
- The intuition:




# Comparing Cardinalities

- Two sets have the same cardinality if there is a way to pair their elements off without leaving any elements uncovered.
- The intuition:



# Infinite Cardinalities




$$n \leftrightarrow 2n$$

$$S = \{ n \mid n \in \mathbb{N} \text{ and } n \text{ is even} \}$$

$$|S| = |\mathbb{N}| = \aleph_0$$

# Infinite Cardinalities



$$|\mathbb{N}| = |\mathbb{Z}| = \aleph_0$$

Pair nonnegative integers with even natural numbers.  
Pair negative integers with odd natural numbers.

***Important Question:***

Do all infinite sets have  
the same cardinality?

$$S = \left\{ \text{Lincoln Penny}, \text{Kennedy Half Dollar} \right\}$$

$$\wp(S) = \left\{ \emptyset, \left\{ \text{Kennedy Half Dollar} \right\}, \left\{ \text{Lincoln Penny} \right\}, \left\{ \text{Lincoln Penny}, \text{Kennedy Half Dollar} \right\} \right\}$$

$$|S| < |\wp(S)|$$

$$S = \left\{ \text{Lincoln Penny}, \text{Jefferson Nickel}, \text{Button} \right\}$$

$$\wp(S) = \left\{ \begin{array}{l} \emptyset, \{ \text{Lincoln Penny} \}, \{ \text{Jefferson Nickel} \}, \{ \text{Button} \}, \\ \{ \text{Lincoln Penny}, \text{Jefferson Nickel} \}, \{ \text{Lincoln Penny}, \text{Button} \}, \{ \text{Jefferson Nickel}, \text{Button} \}, \\ \{ \text{Lincoln Penny}, \text{Jefferson Nickel}, \text{Button} \} \end{array} \right\}$$

$$|S| < |\wp(S)|$$

$$S = \{a, b, c, d\}$$

$$\wp(S) = \{ \\
\emptyset, \\
\{a\}, \{b\}, \{c\}, \{d\}, \\
\{a, b\}, \{a, c\}, \{a, d\}, \{b, c\}, \{b, d\}, \{c, d\} \\
\{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}, \\
\{a, b, c, d\} \\
\}$$

$$|S| < |\wp(S)|$$

If  $|S|$  is infinite, what is the relation between  $|S|$  and  $|\wp(S)|$ ?

Does  $|S| = |\wp(S)|$ ?

If  $|S| = |\wp(S)|$ , we can pair up the elements of  $S$  and the subsets of  $S$  without leaving anything out.

What would that look like?

$$\begin{array}{lcl}
x_0 & \longleftrightarrow & \{ x_0, x_2, x_4, \dots \} \\
x_1 & \longleftrightarrow & \{ x_3, x_5, \dots \} \\
x_2 & \longleftrightarrow & \{ x_0, x_1, x_2, x_5, \dots \} \\
x_3 & \longleftrightarrow & \{ x_1, x_4, \dots \} \\
x_4 & \longleftrightarrow & \{ x_2, \dots \} \\
x_5 & \longleftrightarrow & \{ x_0, x_4, x_5, \dots \} \\
\dots & \longleftrightarrow & \{ \dots \}
\end{array}$$



|         | $x_0$  | $x_1$  | $x_2$  | $x_3$  | $x_4$  | $x_5$  | ... |
|---------|--------|--------|--------|--------|--------|--------|-----|
| $x_0$ ↔ | $x_0,$ |        | $x_2,$ |        | $x_4,$ |        | ... |
| $x_1$ ↔ |        |        |        | $x_3,$ |        | $x_5,$ | ... |
| $x_2$ ↔ | $x_0,$ |        | $x_2,$ |        |        | $x_5,$ | ... |
| $x_3$ ↔ |        | $x_1,$ |        |        | $x_4,$ |        | ... |
| $x_4$ ↔ |        |        | $x_2,$ |        |        |        | ... |
| $x_5$ ↔ | $x_0,$ |        |        |        | $x_4,$ | $x_5,$ | ... |
| ... ↔   | ...    | ...    | ...    | ...    | ...    | ...    | ... |

{  $x_0,$     $x_2,$     $x_5,$    ... }

Which element is  
paired with this  
set?


|         | $x_0$  | $x_1$  | $x_2$  | $x_3$  | $x_4$  | $x_5$  | ... |
|---------|--------|--------|--------|--------|--------|--------|-----|
| $x_0$ ↔ | $x_0,$ |        | $x_2,$ |        | $x_4,$ |        | ... |
| $x_1$ ↔ |        |        |        | $x_3,$ |        | $x_5,$ | ... |
| $x_2$ ↔ | $x_0,$ |        | $x_2,$ |        |        | $x_5,$ | ... |
| $x_3$ ↔ |        | $x_1,$ |        |        | $x_4,$ |        | ... |
| $x_4$ ↔ |        |        | $x_2,$ |        |        |        | ... |
| $x_5$ ↔ | $x_0,$ |        |        |        | $x_4,$ | $x_5,$ | ... |
| ...     | ...    | ...    | ...    | ...    | ...    | ...    | ... |

"Flip" this set.  
Swap what's  
included and  
what's excluded.

{  $x_1,$   $x_3,$   $x_4,$  ... }

|       | $x_0$  | $x_1$  | $x_2$  | $x_3$  | $x_4$  | $x_5$  | ... |
|-------|--------|--------|--------|--------|--------|--------|-----|
| $x_0$ | $x_0$  |        | $x_2,$ |        | $x_4,$ |        | ... |
| $x_1$ |        |        |        | $x_3,$ |        | $x_5,$ | ... |
| $x_2$ | $x_0,$ |        | $x_2,$ |        |        | $x_5,$ | ... |
| $x_3$ |        | $x_1,$ |        |        | $x_4,$ |        | ... |
| $x_4$ |        |        | $x_2,$ |        |        |        | ... |
| $x_5$ | $x_0,$ |        |        |        | $x_4,$ | $x_5,$ | ... |
| ...   | ...    | ...    | ...    | ...    | ...    | ...    | ... |

Which element is paired with this set?




|         | $x_0$  | $x_1$  | $x_2$  | $x_3$  | $x_4$  | $x_5$  | ... |
|---------|--------|--------|--------|--------|--------|--------|-----|
| $x_0$ ↔ | $x_0,$ |        | $x_2,$ |        | $x_4,$ |        | ... |
| $x_1$ ↔ |        |        |        | $x_3,$ |        | $x_5,$ | ... |
| $x_2$ ↔ | $x_0,$ |        | $x_2,$ |        |        | $x_5,$ | ... |
| $x_3$ ↔ |        | $x_1,$ |        |        | $x_4,$ |        | ... |
| $x_4$ ↔ |        |        | $x_2,$ |        |        |        | ... |
| $x_5$ ↔ | $x_0,$ |        |        |        | $x_4,$ | $x_5,$ | ... |
| ... ↔   | ...    | ...    | ...    | ...    | ...    | ...    | ... |

{  $x_1,$   $x_3,$   $x_4,$  ... }

Which element is paired with this set?

|         | $x_0$  | $x_1$  | $x_2$  | $x_3$  | $x_4$  | $x_5$  | ... |
|---------|--------|--------|--------|--------|--------|--------|-----|
| $x_0$ ↔ | $x_0,$ |        | $x_2,$ |        | $x_4,$ |        | ... |
| $x_1$ ↔ |        |        |        | $x_3,$ |        | $x_5,$ | ... |
| $x_2$ ↔ | $x_0,$ |        | $x_2$  |        |        | $x_5,$ | ... |
| $x_3$ ↔ |        | $x_1,$ |        |        | $x_4,$ |        | ... |
| $x_4$ ↔ |        |        | $x_2,$ |        |        |        | ... |
| $x_5$ ↔ | $x_0,$ |        |        |        | $x_4,$ | $x_5,$ | ... |
| ...     | ...    | ...    | ...    | ...    | ...    | ...    | ... |



Which element is paired with this set?



|         | $x_0$  | $x_1$  | $x_2$  | $x_3$  | $x_4$  | $x_5$  | ... |
|---------|--------|--------|--------|--------|--------|--------|-----|
| $x_0$ ↔ | $x_0,$ |        | $x_2,$ |        | $x_4,$ |        | ... |
| $x_1$ ↔ |        |        |        | $x_3,$ |        | $x_5,$ | ... |
| $x_2$ ↔ | $x_0,$ |        | $x_2,$ |        |        | $x_5,$ | ... |
| $x_3$ ↔ |        | $x_1,$ |        |        | $x_4,$ |        | ... |
| $x_4$ ↔ |        |        | $x_2,$ |        |        |        | ... |
| $x_5$ ↔ | $x_0,$ |        |        |        | $x_4,$ | $x_5,$ | ... |
| ...     | ...    | ...    | ...    | ...    | ...    | ...    | ... |


|   |  |        |  |        |        |  |     |   |
|---|--|--------|--|--------|--------|--|-----|---|
| { |  | $x_1,$ |  | $x_3,$ | $x_4,$ |  | ... | } |
|---|--|--------|--|--------|--------|--|-----|---|

Which element is paired with this set?

|         | $x_0$  | $x_1$  | $x_2$  | $x_3$  | $x_4$  | $x_5$  | ... |
|---------|--------|--------|--------|--------|--------|--------|-----|
| $x_0$ ↔ | $x_0,$ |        | $x_2,$ |        | $x_4,$ |        | ... |
| $x_1$ ↔ |        |        |        | $x_3,$ |        | $x_5,$ | ... |
| $x_2$ ↔ | $x_0,$ |        | $x_2,$ |        |        | $x_5,$ | ... |
| $x_3$ ↔ |        | $x_1,$ |        |        | $x_4,$ |        | ... |
| $x_4$ ↔ |        |        | $x_2,$ |        |        |        | ... |
| $x_5$ ↔ | $x_0,$ |        |        |        | $x_4,$ | $x_5,$ | ... |
| ...     | ...    | ...    | ...    | ...    | ...    | ...    | ... |


Which element is paired with this set?

|      |  |        |  |        |        |  |         |      |
|------|--|--------|--|--------|--------|--|---------|------|
| $\{$ |  | $x_1,$ |  | $x_3,$ | $x_4,$ |  | $\dots$ | $\}$ |
|------|--|--------|--|--------|--------|--|---------|------|



|         | $x_0$  | $x_1$  | $x_2$  | $x_3$  | $x_4$  | $x_5$  | ... |
|---------|--------|--------|--------|--------|--------|--------|-----|
| $x_0$ ↔ | $x_0,$ |        | $x_2,$ |        | $x_4,$ |        | ... |
| $x_1$ ↔ |        |        |        | $x_3,$ |        | $x_5,$ | ... |
| $x_2$ ↔ | $x_0,$ |        | $x_2,$ |        |        | $x_5,$ | ... |
| $x_3$ ↔ |        | $x_1,$ |        |        | $x_4,$ |        | ... |
| $x_4$ ↔ |        |        | $x_2,$ |        |        |        | ... |
| $x_5$ ↔ | $x_0,$ |        |        |        | $x_4,$ | $x_5,$ | ... |
| ... ↔   | ...    | ...    | ...    | ...    | ...    | ...    | ... |

Which element is paired with this set?



# The Diagonalization Proof

- No matter how we pair up elements of  $S$  and subsets of  $S$ , the complemented diagonal won't appear in the table.
  - In row  $n$ , the  $n$ th element must be wrong.
- No matter how we pair up elements of  $S$  and subsets of  $S$ , there is *always* at least one subset left over.
- This result is ***Cantor's theorem***: Every set is strictly smaller than its power set:

**If  $S$  is a set, then  $|S| < |\wp(S)|$ .**

# ...And Beyond!

- By Cantor's Theorem:

$$|\mathbb{N}| < |\wp(\mathbb{N})|$$

$$|\wp(\mathbb{N})| < |\wp(\wp(\mathbb{N}))|$$

$$|\wp(\wp(\mathbb{N}))| < |\wp(\wp(\wp(\mathbb{N})))|$$

$$|\wp(\wp(\wp(\mathbb{N})))| < |\wp(\wp(\wp(\wp(\mathbb{N}))))|$$

...

- ***Not all infinite sets have the same size!***
- ***There is no biggest infinity!***
- ***There are infinitely many infinities!***

What does this have to do  
with computation?

*“The set of all computer programs”*

*“The set of all problems to solve”*

# Things on Strings

- A *string* is a sequence of characters.
- Two fun facts about strings:
  - There are *at most* as many programs as there are strings. (All programs are strings)
  - There are *at least* as many problems as there are sets of strings.
- There's an appendix to this slide deck that provides an overview of why these claims are true.
- These facts, plus Cantor's theorem, have terrifying implications.

Every computer program is a string.

So, the number of programs is at most the number of strings.

From Cantor's Theorem, we know that there are more sets of strings than strings.

There are at least as many problems as there are sets of strings.

$$|\mathbf{Programs}| \leq |\mathbf{Strings}| < |\wp(\mathbf{Strings})| \leq |\mathbf{Problems}|$$

*There are more problems to  
solve than there are programs  
to solve them.*

**|Programs| < |Problems|**

# It Gets Worse

- Using more advanced set theory, we can show that there are *infinitely more* problems than solutions.
- In fact, if you pick a totally random problem, the probability that you can solve it is *zero*.
- ***More troubling fact:*** We've just shown that *some* problems are impossible to solve with computers, but we don't know *which* problems those are!

We need to develop a more nuanced understanding of computation.

# Where We're Going

- ***What makes a problem impossible to solve with computers?***
  - Is there a deep reason why certain problems can't be solved with computers, or is it completely arbitrary?
  - How do you know when you're looking at an impossible problem?
  - Are these real-world problems, or are they highly contrived?
- ***How do we know that we're right?***
  - How can we back up our pictures with rigorous proofs?
  - How do we build a mathematical framework for studying computation?


# Next Time

- ***Mathematical Proof***
  - What is a mathematical proof?
  - How can we prove things with certainty?

# Appendix: String Things

# Strings and Programs

- The source code of a computer program is just a (long, structured, well-commented) string of text.
- All programs are strings, but not all strings are necessarily programs.



$$|\mathbf{Programs}| \leq |\mathbf{Strings}|$$

# Strings and Problems

- There is a connection between the number of sets of strings and the number of problems to solve.
- Let  $S$  be any set of strings. This set  $S$  gives rise to a problem to solve:

**Given a string  $w$ , determine whether  $w \in S$ .**

# Strings and Problems

**Given a string  $w$ , determine whether  $w \in S$ .**

- Suppose that  $S$  is the set

$$S = \{ "a", "b", "c", \dots, "z" \}$$

- From this set  $S$ , we get this problem:

**Given a string  $w$ , determine whether  $w$  is a single lower-case English letter.**

# Strings and Problems

**Given a string  $w$ , determine whether  $w \in S$ .**

- Suppose that  $S$  is the set
$$S = \{ "0", "1", "2", \dots, "9", "10", "11", \dots \}$$
- From this set  $S$ , we get this problem:

**Given a string  $w$ , determine whether  $w$  represents a natural number.**

# Strings and Problems

**Given a string  $w$ , determine whether  $w \in S$ .**

- Suppose that  $S$  is the set


$$S = \{ p \mid p \text{ is a legal C++ program} \}$$

- From this set  $S$ , we get this problem:

**Given a string  $w$ , determine whether  $w$  is a legal C++ program.**

# Strings and Problems

- Every set of strings gives rise to a unique problem to solve.
- Other problems exist as well.



$$|\mathbf{Sets\ of\ Strings}| \leq |\mathbf{Problems}|$$