Lecture Notes #14 May 6, 2020

Conditional Expectation

Based on a chapter by Chris Piech

Conditional Distributions

Before we looked at conditional probabilities for events. Here we formally go over conditional probabilities for random variables. The equations for the discrete case is an intuitive extension of our understanding of conditional probability:

Discrete

The conditional probability mass function (PMF) for the discrete case:

$$
p_{X|Y}(x|y) = P(X = x|Y = y) = \frac{P(X = x, Y = y)}{P(Y = y)} = \frac{p_{X,Y}(x, y)}{p_Y(y)}
$$

The conditional cumulative density function (CDF) for the discrete case:

$$
F_{X|Y}(a|y) = P(X \le a|Y = y) = \frac{\sum_{x \le a} p_{X,Y}(x, y)}{p_Y(y)} = \sum_{x \le a} p_{X|Y}(x|y)
$$

Conditional Expectation

We have gotten to know a kind and gentle soul, conditional probability. And we know another funky fool, expectation. Let's get those two crazy kids to play together.

Let *X* and *Y* be jointly discrete random variables. We define the conditional expectation of *X* given $Y = y$ to be:

$$
E[X|Y = y] = \sum_{x} x p_{X|Y}(x|y)
$$

Properties of Conditional Expectation

Here are some helpful, intuitive properties of conditional expectation:

$$
E[g(X)|Y = y] = \sum_{x} g(x)p_{X|Y}(x|y)
$$

$$
E[\sum_{i=1}^{n} X_i|Y = y] = \sum_{i=1}^{n} E[X_i|Y = y]
$$

g(*x*)*pX*|*^Y* (*x*|y) if X and Y are discrete

Law of Total Expectation

The law of total expectation states that: $E[E[X|Y]] = E[X]$.

What?! How is that a thing? Check out this proof:

$$
E[E[X|Y]] = \sum_{y} E[X|Y=y]P(Y=y)
$$

=
$$
\sum_{y} \sum_{x} xP(X=x|Y=y)P(Y=y)
$$

=
$$
\sum_{y} \sum_{x} xP(X=x, Y=y)
$$

=
$$
\sum_{x} \sum_{y} xP(X=x, Y=y)
$$

=
$$
\sum_{x} x \sum_{y} P(X=x, Y=y)
$$

=
$$
\sum_{x} xP(X=x)
$$

=
$$
E[X]
$$

Example 1

You roll two 6-sided dice D_1 and D_2 . Let $X = D_1 + D_2$ and let $Y =$ the value of D_2 .

• What is $E[X|Y = 6]$?

$$
E[X|Y = 6] = \sum_{x} xP(X = x|Y = 6)
$$

= $\left(\frac{1}{6}\right)(7 + 8 + 9 + 10 + 11 + 12) = \frac{57}{6} = 9.5,$

which makes intuitive sense since $6 + E$ [value of *D*₁] = $6 + 3.5$.

• What is $E[X|Y = y]$, where $y = 1, ..., 6$? Let $W =$ the value of D_1 . Then $X = Y + W$, and Y and W are independent.

$$
E[X|Y = y] = E[W + Y|Y = y] = E[W + y|Y = y]
$$

= $y + E[W|Y = y]$ (y is a constant with respect to W)
= $y + \sum_{w} wP(W = w|Y = y)$
= $y + \sum_{w} wP(W = w)$ (W, Y are independent)
= $y + 3.5$

Note that $E[X|Y = y]$ depends on the value y. In other words, $E[X|Y]$ is a function of the random variable *Y*.

Example 2

Consider the following code with random numbers:

```
int Recurse() {
   int x = randomInt(1, 3); // Equally likely values
   if (x == 1) return 3;
   else if (x == 2) return (5 + \text{Recursive}());
   else return (7 + Recurse());
}
```
Let Y = value returned by "Recurse". What is $E[Y]$. In other words, what is the expected return value. Note that this is the exact same approach as calculating the expected run time.

$$
E[Y] = E[Y|X=1]P(X=1) + E[Y|X=2]P(X=2) + E[Y|X=3]P(X=3)
$$

First lets calculate each of the conditional expectations:

$$
E[Y|X = 1] = 3
$$

\n
$$
E[Y|X = 2] = E[5 + Y] = 5 + E[Y]
$$

\n
$$
E[Y|X = 3] = E[7 + Y] = 7 + E[Y]
$$

Now we can plug those values into the equation. Note that the probability of X taking on 1, 2, or 3 is 1/3:

$$
E[Y] = E[Y|X = 1]P(X = 1) + E[Y|X = 2]P(X = 2) + E[Y|X = 3]P(X = 3)
$$

= 3(1/3) + (5 + E[Y])(1/3) + (7 + E[Y])(1/3)
= 15

Hiring Software Engineers

You are interviewing *n* software engineer candidates and will hire only 1 candidate. All orderings of candidates are equally likely. Right after each interview you must decide to hire or not hire. You can not go back on a decision. At any point in time you can know the relative ranking of the candidates you have already interviewed.

The strategy that we propose is that we interview the first *k* candidates and reject them all. Then you hire the next candidate that is better than all of the first *k* candidates. What is the probability that the best of all the *n* candidates is hired for a particular choice of *k*? Let's denote that result $P_k(Best)$. Let *X* be the position in the ordering of the best candidate:

$$
P_k(Best) = \sum_{i=1}^n P_k(Best|X=i)P(X=i)
$$

$$
= \frac{1}{n} \sum_{i=1}^n P_k(Best|X=i)
$$

P^k (*Best*|*X* = *i*) since each position is equally likely

What is $P_k(Best|X = i)$? if $i \leq k$ then the probability is 0 because the best candidate will be rejected without consideration. Sad times. Otherwise we will chose the best candidate, who is in position *i*, only if the best of the first *i* − 1 candidates is among the first *k* interviewed. If the best among the first *i* − 1 is not among the first *k*, that candidate will be chosen over the true best. Since all orderings are equally likely the probability that the best among the *i* − 1 candidates is in the first *k* is:

$$
\frac{k}{i-1} \qquad \qquad \text{if } i > k
$$

Now we can plug this back into our original equation:

$$
P_k(Best) = \frac{1}{n} \sum_{i=1}^n P_k(Best|X = i)
$$

=
$$
\frac{1}{n} \sum_{i=k+1}^n \frac{k}{i-1}
$$
 since we know $P_k(Best|X = i)$

$$
\approx \frac{1}{n} \int_{i=k+1}^n \frac{k}{i-1} di
$$
By Riemann Sum approximation
=
$$
\frac{k}{n} \ln(i = 1) \Big|_{k+1}^n = \frac{k}{n} \ln \frac{n-1}{k} \approx \frac{k}{n} \ln \frac{n}{k}
$$

If we think of $P_k(Best) = \frac{k}{n}$ $\frac{k}{n}$ ln $\frac{n}{k}$ as a function of *k* we can take find the value of *k* that optimizes it by taking its derivative and setting it equal to 0. The optimal value of *k* is *n*/*e*. Where *e* is Euler's number.