

o8: Poisson and More

Lisa Yan and Jerry Cain
September 30, 2020

Quick slide reference

3	Poisson	08a_poisson
11	Poisson, continued	08b_poisson_ii
17	Other Discrete RVs	08c_other_discrete
25	Exercises	LIVE
34	Poisson approximation	LIVE
52	Extra: Modeling exercise: Hurricanes	08e_extra_modeling

Poisson RV

Before we start

The natural exponent e :


$$\lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^n = \underline{e^{-\lambda}}$$

[https://en.wikipedia.org/wiki/E_\(mathematical_constant\)](https://en.wikipedia.org/wiki/E_(mathematical_constant))

Jacob Bernoulli
while studying
compound interest
in 1683



Algorithmic ride sharing



Probability of k requests from this area in the next 1 minute?

Suppose we know:


On average, $\lambda = 5$ requests per minute

Algorithmic ride sharing, approximately

Probability of k requests from this area in the next 1 minute?

On average, $\lambda = 5$ requests per minute

Break a minute down into 60 seconds:



At each second:

- Independent trial
- You get a request (1) or you don't (0).

Let $X = \#$ of requests in minute.

$$E[X] = \lambda = 5 = np$$

$$X \sim \text{Bin}(n = 60, p = 5/60)$$

$$P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$$




But what if there are *two* requests in the same second?

Algorithmic ride sharing, approximately

Probability of k requests from this area in the next 1 minute?

On average, $\lambda = 5$ requests per minute

Break a minute down into 60,000 **milliseconds**:



At each **millisecond**:

- Independent trial
- You get a request (1) or you don't (0).

Let $X = \#$ of requests in minute.

$$E[X] = \lambda = 5 = np$$

$$X \sim \text{Bin}(n = 60000, p = \lambda/n)$$

$$P(X = k) = \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k}$$




But what if there are *two* requests in the same **millisecond**?

Algorithmic ride sharing, approximately

Probability of k requests from this area in the next 1 minute?

On average, $\lambda = 5$ requests per minute

Break a minute down into **infinitely small** buckets:



For each time bucket:

- Independent trial
- You get a request (1) or you don't (0).

Let $X = \#$ of requests in minute.

$$E[X] = \lambda = 5 = np$$

$$X \sim \text{Bin}(n, p = \lambda/n)$$

$$P(X = k) = \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k}$$

Who wants to see some cool math?

Binomial in the limit

$$\lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^n = e^{-\lambda}$$

$$P(X = k) = \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \quad \text{Expand} = \lim_{n \rightarrow \infty} \frac{n!}{k!(n-k)!} \frac{\lambda^k}{n^k} \frac{\left(1 - \frac{\lambda}{n}\right)^n}{\left(1 - \frac{\lambda}{n}\right)^k}$$

$$\text{Rearrange} = \lim_{n \rightarrow \infty} \frac{n!}{n^k (n-k)!} \frac{\lambda^k}{k!} \frac{\left(1 - \frac{\lambda}{n}\right)^n}{\left(1 - \frac{\lambda}{n}\right)^k} \quad \text{Def natural exponent} = \lim_{n \rightarrow \infty} \frac{n!}{n^k (n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

$$\text{Expand} = \lim_{n \rightarrow \infty} \frac{\overbrace{n(n-1)\cdots(n-k+1)}^{k \text{ terms}}}{n^k} \frac{(n-k)!}{(n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

$$\text{Limit analysis + cancel} = \lim_{n \rightarrow \infty} \frac{n^k}{n^k} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{1} \quad \text{Simplify} = \frac{\lambda^k}{k!} e^{-\lambda}$$

$\underbrace{\left(1 - \frac{\lambda}{n}\right)^k}_{(1-p)^k} \quad \lambda = np$
 $\underbrace{1^k}_{1^k}$

Algorithmic ride sharing



Probability of k requests from this area in the next 1 minute?

On average, $\lambda = 5$ requests per minute

$$P(X = k) = \frac{\lambda^k}{k!} e^{-\lambda}$$


**Poisson
distribution**

Poisson, continued

Poisson Random Variable

Consider an experiment that lasts a fixed interval of time.

def A **Poisson** random variable X is the number of successes over the experiment duration, assuming **the time that each success occurs is independent** and the average # of requests over time is constant. $E[X]$



Examples:

- # earthquakes per year
- # server hits per second
- # of emails per day

Poisson Random Variable

Consider an experiment that lasts a fixed interval of time.

def A **Poisson** random variable X is the number of successes over the experiment duration, assuming the time that each success occurs is independent and the average # of requests over time is constant.

$$X \sim \text{Poi}(\lambda)$$

Support: $\{0, 1, 2, \dots\}$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation $E[X] = \lambda$

Variance $\text{Var}(X) = \lambda$

Examples:

- # earthquakes per year
- # server hits per second
- # of emails per day

Yes, expectation == variance for Poisson RV! More later.

Simeon-Denis Poisson



French mathematician (1781 – 1840)

- Published his first paper at age 18
- Professor at age 21
- Published over 300 papers

“Life is only good for two things: doing mathematics and teaching it.”

Earthquakes

$$X \sim \text{Poi}(\lambda) \quad p(k) = e^{-\lambda} \frac{\lambda^k}{k!}$$
$$E[X] = \lambda$$

There are an average of 2.79 major earthquakes in the world each year, and major earthquakes occur independently.


What is the probability of 3 major earthquakes happening next year?

1. Define RVs

X : # major quakes in the next year
 $E[X] = 2.79$ $X \sim \text{Poi}(\lambda = 2.79)$

2. Solve

$$P(X=3) = e^{-2.79} \frac{(2.79)^3}{3!}$$
$$\hat{=} 0.23$$



Are earthquakes really Poissonian?

Bulletin of the Seismological Society of America

Vol. 64

October 1974

No. 5

IS THE SEQUENCE OF EARTHQUAKES IN SOUTHERN CALIFORNIA,
WITH AFTERSHOCKS REMOVED, POISSONIAN?


BY J. K. GARDNER and L. KNOPOFF

ABSTRACT

Yes.

Other Discrete RVs

Grid of random variables



Geometric RV

Consider an experiment: independent trials of $\text{Ber}(p)$ random variables.

def A **Geometric** random variable X is the # of trials until the ^{fixed} first success.


$X \sim \text{Geo}(p)$

Support: $\{1, 2, \dots\}$

PMF	$P(X = k) = (1 - p)^{k-1} p$
Expectation	$E[X] = \frac{1}{p}$ <i>average # of trials until first success</i>
Variance	$\text{Var}(X) = \frac{1-p}{p^2}$

Examples:

- Flipping a coin ($P(\text{heads}) = p$) until first heads appears
- Generate bits with $P(\text{bit} = 1) = p$ until first 1 generated



$\text{Ber}(1/2) \rightarrow \text{Geo}(p=1/2)$ $(1-p)^{k-1} p$
 $E[X] = 2$

Negative Binomial RV

Consider an experiment: independent trials of $\text{Ber}(p)$ random variables.

def A **Negative Binomial** random variable X is the # of trials until r successes.

Fixed

$$X \sim \text{NegBin}(r, p)$$

Support: $\{r, r + 1, \dots\}$

PMF

$$P(X = k) = \binom{k-1}{r-1} (1-p)^{k-r} p^r$$

Expectation

$$E[X] = \frac{r}{p}$$



Variance

$$\text{Var}(X) = \frac{r(1-p)}{p^2}$$

derived later
~~next slide~~

Examples:


- Flipping a coin until r^{th} heads appears
- # of strings to hash into table until bucket 1 has r entries



$$\binom{k-1}{r-1} \cdot \dots \cdot p^r (1-p)^{k-r}$$

$$\text{Geo}(p) = \text{NegBin}(1, p)$$

Grid of random variables




Catching Pokemon

Wild Pokemon are captured by throwing Pokeballs at them.

- Each ball has probability $p = 0.1$ of capturing the Pokemon.
- Each ball is an independent trial.

What is the probability that you catch the Pokemon on the 5th try?



1. Define events/
RVs & state goal

$X \sim$ some distribution

Want: $P(X = 5)$

2. Solve

- A. $X \sim \text{Bin}(5, 0.1)$
- B. $X \sim \text{Poi}(0.5)$
- C. $X \sim \text{NegBin}(5, 0.1)$
- D. $X \sim \text{NegBin}(1, 0.1)$
- E. $X \sim \text{Geo}(0.1)$
- F. None/other



Catching Pokemon


Wild Pokemon are captured by throwing Pokeballs at them.

- Each ball has probability $p = 0.1$ of capturing the Pokemon.
- Each ball is an independent trial.

What is the probability that you catch the Pokemon on the 5th try?

FFFFT

#trials



1. Define events/
RVs & state goal

2. Solve

$X \sim$ some distribution

Want: $P(X = 5)$

What does Sample Space look like?
• have to catch Pokemon
• # trials is flexible

- ~~A.~~ $X \sim \text{Bin}(5, 0.1)$
- ~~B.~~ $X \sim \text{Poi}(0.5)$ ✓
- ~~C.~~ $X \sim \text{NegBin}(5, 0.1)$
- D. $X \sim \text{NegBin}(1, 0.1)$
- E. $X \sim \text{Geo}(0.1)$
- F. None/other

must be r : #successes


Catching Pokemon

$$X \sim \text{Geo}(p) \quad p(k) = (1 - p)^{k-1} p$$

Wild Pokemon are captured by throwing Pokeballs at them.

- Each ball has probability $p = 0.1$ of capturing the Pokemon.
- Each ball is an independent trial.

What is the probability that you catch the Pokemon on the 5th try?



1. Define events/
RVs & state goal

2. Solve

$$X \sim \text{Geo}(0.1)$$

$$\begin{aligned} \text{Want: } P(X = \underline{5}) &= (1-p)^{k-1} p \\ &= (0.9)^4 \cdot 0.1 \\ &\approx 0.066 \end{aligned}$$

o8: Poisson and More (live)

Lisa Yan and Jerry Cain
September 30, 2020



Discrete RVs




The hardest part of problem-solving is determining what is a random variable .

CS109 Learning Goal: Use new RVs

Let's say you are learning about servers/networks.

You read about the M/D/1 queue:



“The service time busy period is distributed as a Borel with parameter $\mu = 0.2$.”

Goal: You can recognize terminology and understand experiment setup.

Big Q: Fixed parameter or random variable?

Parameter

What is **common** among all outcomes of our experiment?

Ber(p) Geo(p)
Bin(n, p) NegBin(r, p)
Poi(λ)

Examples so far:

- Prob. success
- # total trials n
- # target successes
- Average rate of success

Random variable


What **differentiates** our event from the rest of the sample space?

$X \sim \text{Geo}(p)$ $X=1$ vs $X=100$


Examples so far:

- # of successes
- Time until success (for some definition of time)

Grid of random variables



Grid of random variables



Breakout Rooms

Check out the question on the next slide (Slide 32). Post any clarifications here!

<https://us.edstem.org/courses/2678/discussion/134631>

Breakout rooms: 5 min. Introduce yourself!



Kickboxing with RVs

How would you model the following?

1. # of snapchats you receive in a day
2. # of children until the first one with brown eyes (same parents)
3. If stock went up (1) or down (0) in a day
4. # of probability problems you try until you get 5 correct (if you are randomly correct)
5. # of years in some decade with more than 6 Atlantic hurricanes

Choose from:

A. Ber(p)	C. Poi(λ)
B. Bin(n, p)	D. Geo(p)
	E. NegBin(r, p)



Kickboxing with RVs

How would you model the following?

1. # of snapchats you receive in a day
2. # of children until the first one with brown eyes (same parents)
3. If stock went up (1) or down (0) in a day
4. # of probability problems you try until you get 5 correct (if you are randomly correct)
5. # of years in some decade with more than 6 Atlantic hurricanes

Note: These exercises are designed to build intuition; in a problem statement, you will generally have more clues.

Choose from: C. Poi(λ)
A. Ber(p) D. Geo(p)
B. Bin(n, p) E. NegBin(r, p)

C. Poi(λ)

D. Geo(p) or E. NegBin($1, p$)

A. Ber(p) or B. Bin($1, p$)

E. NegBin($r = 5, p$)

B. Bin($n = 10, p$), where
 $p = P(\geq 6 \text{ hurricanes in a year})$
calculated from C. Poi(λ)

Poisson Approximation

$$X \sim \text{Poi}(\lambda)$$

Support: $\{0, 1, 2, \dots\}$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation $E[X] = \lambda$

Variance $\text{Var}(X) = \lambda$

In CS109, a Poisson RV $X \sim \text{Poi}(\lambda)$ most often models

1. # of successes in a fixed interval of time, where successes are independent
 $\lambda = E[X]$, average success/interval

1. Web server load

$$X \sim \text{Poi}(\lambda) \quad p(k) = e^{-\lambda} \frac{\lambda^k}{k!}$$
$$E[X] = \lambda$$

Consider requests to a web server in 1 second.

- In the past, server load averages 2 hits/second, where hits arrive independently.
- Let $X = \#$ hits the server receives in a second.

$$E[X] = 2 = \lambda$$

What is $P(X < 5)$?

Define RVs

$$X \sim \text{Poi}(\lambda = 2)$$

Solve

$$P(X < 5) = \sum_{k=0}^4 p(k) = \sum_{k=0}^4 e^{-2} \cdot \frac{2^k}{k!}$$

$$\approx 0.95$$

alternatively: $1 - P(X \geq 5)$

Poisson Random Variable

$$X \sim \text{Poi}(\lambda)$$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation $E[X] = \lambda$

Support: $\{0, 1, 2, \dots\}$


Variance $\text{Var}(X) = \lambda$

In CS109, a Poisson RV $X \sim \text{Poi}(\lambda)$ most often models

1. # of successes in a fixed interval of time, where successes are independent
 $\lambda = E[X]$, average success/interval
2. Approximation of $Y \sim \text{Bin}(n, p)$ where n is large and p is small.
 $\lambda = E[Y] = np$

Approximation works even when trials not entirely independent.

2. DNA



All the movies, images, emails and other digital data from more than 600 smartphones (10,000 GB) can be stored in the faint pink smear of DNA at the end of this test tube.

What is the probability that DNA storage stays uncorrupted?

2. DNA

What is the probability that DNA storage stays uncorrupted?


- In DNA (and real networks), we store large strings.
- Let string length be long, e.g., $n \approx 10^4$
- Probability of corruption of each base pair is very small, e.g., $p = 10^{-6}$
- Let $X = \#$ of corruptions.

What is $P(\text{DNA storage is uncorrupted}) = P(X = 0)$?

1. Approach 1:

$$X \sim \text{Bin}(n = 10^4, p = 10^{-6})$$

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}$$

unwieldy!  $= \binom{10^4}{0} 10^{-6 \cdot 0} (1 - 10^{-6})^{10^4 - 0}$
 ≈ 0.99049829

Handwritten notes: "what if k ≠ 0?" with an arrow pointing to the binomial coefficient, and "0" under the exponent of the second term.

2. Approach 2:

$$X \sim \text{Poi}(\lambda = 10^4 \cdot 10^{-6} = 0.01)$$

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!} = e^{-0.01} \frac{0.01^0}{0!}$$

$$= e^{-0.01}$$

$$\approx 0.99049834$$

a good approximation! 

Think

Slide 41 has a question to go over by yourself.

Post any clarifications here or in chat!

<https://us.edstem.org/courses/2678/discussion/134631>

Think by yourself: 1 min



(by yourself)

When is a Poisson approximation appropriate?

$$P(X = k) = \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} = \dots$$

Under which conditions will $X \sim \text{Bin}(n, p)$ behave like $\text{Poi}(\lambda)$, where $\lambda = np$?

Def natural exponent

$$= \lim_{n \rightarrow \infty} \frac{n!}{n^k (n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

Expand

$$= \lim_{n \rightarrow \infty} \frac{n(n-1)\dots(n-k+1)}{n \cdot n \dots n^k \dots n} \frac{(n-k)!}{(n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

Limit analysis


$$= \lim_{n \rightarrow \infty} \frac{n^k}{n^k} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{1} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

large n
small p
 $(1-p)^k$
 $\frac{\lambda}{n} = p$

Simplify

$$= \frac{\lambda^k}{k!} e^{-\lambda}$$

- A. Large n , large p
- B. Small n , small p
- C. Large n , small p**
- D. Small n , large p
- E. Other



Poisson approximation

$$X \sim \text{Poi}(\lambda)$$
$$E[X] = \lambda$$

$$Y \sim \text{Bin}(n, p)$$
$$E[Y] = np$$


Poisson approximates Binomial when n is large, p is small, and $\lambda = np$ is “moderate.”

Different interpretations of “moderate”:

- $n > 20$ and $p < 0.05$
- $n > 100$ and $p < 0.1$

Poisson is Binomial in the limit:

- $\lambda = np$, where $n \rightarrow \infty, p \rightarrow 0$



Poisson Random Variable

Consider an experiment that lasts a fixed interval of time.

def A **Poisson** random variable X is the number of occurrences over the experiment duration.

$$X \sim \text{Poi}(\lambda)$$

Support: $\{0, 1, 2, \dots\}$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation $E[X] = \lambda$

Variance $\text{Var}(X) = \lambda$

Examples:

- # earthquakes per year
- # server hits per second
- # of emails per day

Time to show intuition for why expectation == variance!

Properties of $\text{Poi}(\lambda)$ with the Poisson paradigm

Recall the Binomial:

$$Y \sim \text{Bin}(n, p) \quad \begin{array}{ll} \text{Expectation} & E[Y] = np \\ \text{Variance} & \text{Var}(Y) = np(1 - p) \end{array}$$

Consider $X \sim \text{Poi}(\lambda)$, where $\lambda = np$ ($n \rightarrow \infty, p \rightarrow 0$):

$$X \sim \text{Poi}(\lambda) \quad \begin{array}{ll} \text{Expectation} & E[X] = \lambda \\ \text{Variance} & \text{Var}(X) = \lambda \end{array}$$

Proof:

$$E[X] = np = \lambda$$
$$\text{Var}(X) = np(1 - p) \rightarrow \lambda(1 - 0) = \lambda$$

λ



Poisson Approximation, approximately

Poisson can still provide a **good approximation of the Binomial**, even when assumptions are “mildly” violated.

$\text{Bin}(n, p)$

You can apply the Poisson approximation when:



- “Successes” in trials are not entirely independent
e.g.: # entries in each bucket in large hash table.
- Probability of “Success” in each trial varies (slightly),
like a **small relative change** in a very small p
e.g.: Average # requests to web server/sec may fluctuate
slightly due to load on network



We won't explore this too much,
but I want you to know it exists.

Think

Slide 47 has a question to go over by yourself.

Post any clarifications here or in chat!

<https://us.edstem.org/courses/2678/discussion/134631>

Think by yourself: 2 min



(by yourself)

Can these Binomial RVs be approximated?

Poisson approximates Binomial when n is large, p is small, and $\lambda = np$ is “moderate.”

Different interpretations of “moderate”:

- $n > 20$ and $p < 0.05$
- $n > 100$ and $p < 0.1$




Poisson is Binomial in the limit:

- $\lambda = np$, where $n \rightarrow \infty, p \rightarrow 0$



type in
chat:

yes, yes, yes



Can these Binomial RVs be approximated?




Poisson approximates Binomial when n is large, p is small, and $\lambda = np$ is “moderate.”

Different interpretations of “moderate”:

- $n > 20$ and $p < 0.05$
- $n > 100$ and $p < 0.1$

Poisson is Binomial in the limit:

- $\lambda = np$, where $n \rightarrow \infty, p \rightarrow 0$



A Real License Plate Seen at Stanford



No, it's not mine...
but I kind of wish it was.



still
no idea

Interlude for jokes/announcements

Announcements

PS2 on-time Monday 1 PM
late Wed 1 PM ← solns

Quiz #1

Time frame: Wednesday 10/7⁷ 2:00pm – Friday 10/8⁹ 12:59^{1:00}pm PT

Covers: Up to end of Week 2 (including Lecture 6)

Anand and Sandra's Review session: Sunday 10/4 6 – 8pm PT

recorded [Zoom link](#)

Info and practice: <https://web.stanford.edu/class/cs109/exams/quizzes.html>

Python tutorial #2

When: today 3:30-4:30PT

Recorded? yes

Notes: posted online


Office Hour update

Lisa's Tea Hour Thursdays 9:30-11am PT

- Casual, any CS109 or non-CS109 questions here
- Collaborate on jigsaw puzzle

Modeling exercise: Hurricanes

Hurricanes





What is the probability of an extreme weather event?

How do we model the number of hurricanes in a season (year)?

1. Graph your distribution.

1. Graph: Hurricanes per year since 1851


Which graph is a histogram (i.e., distribution) of frequency (# of hurricanes per year)?




1. Graph: Hurricanes per year since 1851

Which graph is a histogram (i.e., distribution) of frequency (# of hurricanes per year)?

A.




B.



Looks kinda Poissonian!

Hurricanes



How do we model the number of hurricanes in a season (year)?

2. Find a reasonable distribution and compute parameters.

2. Find a distribution: Python SciPy RV methods

```
from scipy import stats           # great package
X = stats.poisson(8.5)           #  $X \sim \text{Poi}(\lambda = 8.5)$ 
X.pmf(2)                         #  $P(X = 2)$ 
```

Function	Description
<code>X.pmf(k)</code>	$P(X = k)$
<code>X.cdf(k)</code>	$P(X \leq k)$
<code>X.mean()</code>	$E[X]$
<code>X.var()</code>	$\text{Var}(X)$
<code>X.std()</code>	$\text{SD}(X)$


SciPy reference:

<https://docs.scipy.org/doc/scipy/reference/generated/scipy.stats.poisson.html>

2. Find a distribution

Until 1966, things look pretty Poisson.

What is the probability of over **15 hurricanes** in a season (year) given that the distribution doesn't change?



$$P(X > 15) = 1 - P(X \leq 15)$$


$$= 1 - \sum_{k=0}^{15} P(X = k)$$

$$= 1 - 0.986 = 0.014$$

$$X \sim \text{Poi}(\lambda = 8.5)$$

You can calculate this PMF using your favorite programming language. Or Python3.

Hurricanes




How do we model the number of hurricanes in a season (year)?

3. Identify and explain outliers.

3. Improbability

Since 1966, there have been two years with over 30 hurricanes.

What is the probability of over 30 hurricanes in a season (year) given that the distribution doesn't change?




$$P(X > 30) = 1 - P(X \leq 30)$$

$$= 1 - \sum_{k=0}^{30} P(X = k) \quad X \sim \text{Poi}(\lambda = 8.5)$$


$$= 2.2\text{E} - 09$$

3. The distribution has changed.



1851-
1966




Since
1966



3. What changed?



3. What changed?



It's not just climate change. We also have tools for better data collection.