

Class 16: Agenda/Questions

1 Announcements

- HW7 was due today, HW8 (last one!!) is due next Wednesday.
- Our final exam will go out next Wednesday night, due the following Tuesday (March 15th) at 11:30am.

2 Questions?

Any questions from the minilectures and/or the quiz? (Martingales, the Doob martingale, Azuma-Hoeffding)

3 Chromatic numbers

In this exercise we'll practice using Azuma-Hoeffding

Group Work

Let $G \sim G_{n,p}$ be a Erdos-Renyi random graph (so there are n vertices, and each edge is present independently with probability p). Let $A = \chi(G)$ be the chromatic number of G . That is, A is the minimum number of colors necessary to properly color G (ie color the nodes of the graph such that no pair of neighboring nodes are assigned the same color).

1. Consider the Doob *vertex exposure* martingale. That is:
 - For $i \in \{1, \dots, n\}$, let X_i denote the the status of the edges between vertex i and vertices $\{i + 1, \dots, n\}$.
 - $Z_i = \mathbb{E}[A|X_1, \dots, X_i]$

Use the Azuma-Hoeffding inequality to show that

$$\Pr[|A - \mathbb{E}[A]| > c\sqrt{n}] \leq 2 \exp(-c^2/2).$$

(Notice that you may not know what $\mathbb{E}[A]$ is—that's okay!)

Hint: To use Azuma-Hoeffding, you need to bound $|Z_i - Z_{i-1}|$. How much can your expectation of the chromatic color change if I tell you additional information about a single vertex?

2. Repeat the same exercise with the *edge exposure* martingale:

- Let X_i denote the status of the i 'th edge, for $i \in \{1, \dots, \binom{n}{2}\}$.
- $Z_i = \mathbb{E}[A|X_1, \dots, X_i]$

Do you get the same thing? Do you get something better? Worse?

3. (**CHALLENGING**, but something to think about if you finish early: What can you say about $\mathbb{E}[A]$?

Note: If you're interested, check out <https://arxiv.org/abs/0706.1725> for a surprisingly strong statement about the chromatic number of random graphs!!

4 Gambling

In this exercise, we'll get yet more practice applying Azuma-Hoeffding.

Group Work

Consider the following gambling game:

- At time t , you can choose to bet *any* amount you like in $[0, B]$, where B is a house limit.
- A fair coin is flipped. If it's heads, you win the amount that you bet; if tails, you lose the amount that you bet.

You're allowed to be in debt; you don't stop when you run out of money.

1. Suppose that the amount you bet is a deterministic function of everything that's happened so far. Set up a martingale $\{Z_t\}$ (with respect some sequence $\{X_t\}$ that you have to define) so that Z_t is the amount of money you have at time t .
2. Use the Azuma-Hoeffding inequality to bound

$$\Pr[|Z_n| \geq cB\sqrt{n}].$$

3. Now suppose that you can use *any* betting strategy you like, even a randomized one. Is your martingale from part 1 still a martingale? If not, repeat parts 1 and 2 when your betting strategy can be randomized.