

Class 4: Agenda, Questions, and Links

1 Warm-Up

Suppose you flip a p -biased coin n times. What does Markov's inequality tell you about the probability that you see more than $2pn$ heads? What does Chebyshev's inequality tell you about that probability? (When prompted, enter your answers into the ohay "text responses" box.)

2 Announcements

- HW1 was due before today's class! HW2 was posted Monday, and will be due before class in a week.

3 Questions?

Any questions from the minilectures or warmup? (Markov and Chebyshev's inequalities).

- Go into small groups and ask each other your questions.
- Ask any questions that the group can't resolve in the chat, or DM one of the course staff and we'll come to your room.

4 Sampling-Based Median

Sampling-Based Median Algorithm

Median:(A list S of n distinct numbers, where n is odd):

1. Let $t = n^{3/4}$. Sample $R = \{r_1, \dots, r_t\} \subseteq S$ by drawing r_i uniformly at random, independently.
2. Sort R in time $O(t \log t)$. Henceforth, assume that $r_1 \leq r_2 \leq \dots \leq r_t$.
3. Let $a = r_{t/2-\sqrt{n}}$, $b = r_{t/2+\sqrt{n}}$.
4. Let $N_{<a}$ and $N_{>b}$ denote the number of elements in S less than a and greater than b respectively.
5. Let $T = \{x \in S : a \leq x \leq b\}$. Construct T , and compute $N_{<a}$ and $N_{>b}$, in time $O(n)$.
6. If $|T| < 4t$, sort T in time $O(t \log t)$; otherwise output FAIL.

7. If $N_{<a}, N_{>b} \leq n/2$ (aka, $\text{median}(S) \in T$):
 - Return the i 'th smallest element of T , where $i = (n + 1)/2 - N_{<a}$.
8. Otherwise, output FAIL.

[We'll see an example on a slide.]

[**Note:** Above there should be some floors or ceilings or something. Don't worry about it, and ignore off-by-one errors throughout this class.]

5 Analyzing the sampling-based median algorithm

You will analyze this algorithm in group work.

Group Work

Note: Throughout this group work, don't worry about \leq vs $<$, or whether or not something is true up to ± 1 , or anything small like that.

1. Make sure that you all understand the algorithm. Pseudo-code is above, and the one-slide example is available on the course website (cs265.stanford.edu), in the class-by-class resources for Class 4. Ask/answer any questions that you have amongst yourselves, and ask in chat or flag down a member of the course staff if you still have questions.
2. Suppose that you could show that:
 - with probability ≥ 0.9 , the median of S is in the list T ; and
 - with probability ≥ 0.9 , $|T| < 4t$.

Explain (to each other) why these two things would imply that the algorithm returns the correct answer with probability ≥ 0.8 . And if it does not return the median then it returns FAIL.

3. Convince yourself that this algorithm uses at most $O(n)$ operations. What is the leading constant in this big-Oh notation? (Assuming that "sample a random element of S ", and comparing two numbers are each single operations).

At this point, please tick #3 on the Numbers poll

4. In the following parts, you will show that the median of S is in T , with probability at least 0.9. Let m be the median of S . Consider two events:
 - $|\{r_i \in R : r_i < m\}| < \frac{t}{2} + \sqrt{n}$
 - $|\{r_i \in R : r_i > m\}| < \frac{t}{2} + \sqrt{n}$(a) Explain why, if both of these events hold, then $\text{median}(S) \in T$.

- (b) Use Chebyshev's inequality to bound the probability that the first event does not hold. (Hint: let X_i be the indicator random variable that is 1 iff $r_i \leq m$, and consider $\sum_i X_i$).
- (c) Convince yourself that the same argument will work for the second event, and write a statement of the form:

$$\Pr[\text{median}(S) \in T] \geq 1 - \text{----}.$$

At this point, please tick #4 on the Numbers poll

5. Now, we turn our attention to the probability that $|T| < 4t$.
- (a) Explain why it is sufficient to show that a is *not* one of the smallest $n/2 - 2t$ elements of S , and b is *not* one of the largest $n/2 + 2t$ elements of S .
 - (b) Use Chebyshev's inequality to bound the probability that a is not one of the smallest $n/2 - 2t$ elements of S . (Hint: Consider the indicator random variable Y_i that is 1 if r_i is in the smallest $n/2 - 2t$ elements of S . Argue that a is one of the smallest $n/2 - 2t$ elements of S iff $\sum_i Y_i \geq t/2 - \sqrt{n}$ (why?) and apply Chebyshev's inequality.)
 - (c) Convince yourself that the analogous statement for b , and write a statement of the form:

$$\Pr[|T| < 4t] \geq 1 - \text{-----}.$$

At this point, please tick #5 on the Numbers poll