

## Class 18: Agenda and Questions

## 1 Warm-Up

**Group Work**

Suppose that  $X$  is a  $k$ -source on  $\{0, 1\}^n$ . Let  $N = 2^n$ . Let  $\sigma \in \mathbb{R}^N$  be the “vectorized” version of the pmf of  $X$ . That is,  $\sigma_i = \Pr[X = i] \quad \forall i \in \{0, \dots, N - 1\}$ , where we associate a number  $i < N$  with its binary expansion in  $\{0, 1\}^n$ .

1. Why is  $\|\sigma\|_\infty \leq 2^{-k}$ ?
2. Argue that  $\|\sigma\|_2 \leq 2^{-k/2}$ .

**Hint:** Use the fact that for any vector  $x$ ,  $\|x\|_2^2 \leq \|x\|_\infty \|x\|_1$  (why is this true?).

## 2 Announcements

- HW7 due Friday! Last one!
- **Exam:** Friday March 21, 8:30am-11:30am, Hewlett 201. Practice exam out now!

## 3 Questions?

Questions from the minilectures, quiz, warm-up? (Expanders, extractors?)

## 4 Recap

Recall the definition of a  $(k, \varepsilon)$ -extractor:

**Definition 1.** A function  $\text{Ext} : \{0, 1\}^n \times \{0, 1\}^d \rightarrow \{0, 1\}^m$  is a  $(k, \varepsilon)$  extractor if, for all  $k$ -sources  $X$  on  $\{0, 1\}^n$ ,  $\|\text{Ext}(X, U_d) - U_m\| \leq \varepsilon$ .

Above,  $\|\cdot\|$  is the total variation distance, and  $U_d$  refers to the uniform distribution on  $d$  bits. Suppose that  $G = (V, E)$  is an (undirected, unweighted) degree- $D$  expander graph with  $|V| = N$ , and with expansion parameters  $\lambda(G) \leq 1/2$ . Recall that  $\lambda(G) = \max\{\lambda_2, |\lambda_N|\}$ , where  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$  are the eigenvalues of  $A$ , where  $A$  is the *normalized adjacency matrix* of  $G$ . (aka,  $A_{ij}$  is  $1/D$  if  $\{i, j\} \in E$  and is zero otherwise).

## 5 Extractors *from* expanders

Some slides illustrating a construction of an extractor. A description is below for reference.

Let  $\varepsilon > 0$ . Let  $N = 2^n$  and fix a bijection between  $\{0, 1\}^n$  and  $V$ , where  $V$  is the vertex set of  $G$  above. Fix any  $k \leq n$ . Let  $d = \log(D) \cdot \ell$ , where  $\ell = (n - k)/2 + \log(1/\varepsilon)$ . Consider the following function  $\text{Ext} : \{0, 1\}^n \times \{0, 1\}^d \rightarrow \{0, 1\}^n$ . On input  $x, s \in \{0, 1\}^n \times \{0, 1\}^d$ :

- Treat  $x \in \{0, 1\}^n$  as an element of  $V$ .
- Treat  $s \in \{0, 1\}^d$  as a string of  $\ell$  numbers in  $\{0, 1, \dots, D\}$ . (That is, break up  $s$  into  $\ell$  chunks, each  $\log(D)$  bits long). Say these numbers are  $a_1, a_2, \dots, a_\ell$ .
- Consider the following walk on  $G$ : let  $x^{(0)} = x$ . For  $i = 1, 2, \dots, \ell$ , get from  $x^{(i-1)}$  to  $x^{(i)}$  by choosing the  $a_i$ 'th neighbor of  $x^{(i-1)}$ .
- Output  $x^{(\ell)} \in V$ , which we treat as an element of  $\{0, 1\}^n$ .

## 6 Group work: this is a good extractor!

In this section, you'll show that  $\text{Ext}$  is a  $(k, \varepsilon)$  extractor.

### Group Work

1. Let  $\sigma$  be the “vectorized” pmf of  $X$  (as in the warm-up). Explain why the distribution of  $\text{Ext}(X, U_d)$  is given by  $A^\ell \cdot \sigma$ . (Recall that  $A$  is the normalized adjacency matrix of  $G$ ).
2. Let  $\pi = \frac{1}{N}\mathbf{1}$  be the vector that corresponds to the uniform distribution. Explain why

$$\|U_n - \text{Ext}(X, U_d)\| = \|\pi - A^\ell \cdot \sigma\| \leq \frac{\sqrt{N}}{2} \lambda(G)^\ell \|\pi - \sigma\|_2.$$

**Hint:** Mimic a computation that we did in the Expanders minilecture to show that random walks mix quickly when  $\lambda(G)$  is small.

3. Argue that  $\|\pi - \sigma\|_2 \leq 2 \cdot 2^{-k/2}$ .

**Hint:** To get started, note that  $\|\pi - \sigma\|_2 \leq \|\pi\|_2 + \|\sigma\|_2$  by the triangle inequality.

4. Assume that  $G$  is a good enough expander that  $\lambda(G) \leq 1/2$ . (It turns out that these exist for large enough degrees  $D$ ). Conclude that  $\|U_n - \text{Ext}(X, U_d)\| \leq \varepsilon$  and thus  $\text{Ext}$  is a  $(k, \varepsilon)$  extractor.