

# CS156: The Calculus of Computation

Zohar Manna

Autumn 2008

## Chapter 7: Quantified Linear Arithmetic

## Quantifier Elimination (QE)

Algorithm for elimination of all quantifiers of formula  $F$  until quantifier-free formula (qff)  $G$  that is equivalent to  $F$  remains

Note: Could be enough if  $F$  is equisatisfiable to  $G$ , that is  $F$  is satisfiable iff  $G$  is satisfiable

A theory  $T$  admits quantifier elimination iff

there is an algorithm that given  $\Sigma$ -formula  $F$  returns a quantifier-free  $\Sigma$ -formula  $G$  that is  $T$ -equivalent to  $F$ .

## Example: $\exists x. 2x = y$

For  $\Sigma_{\mathbb{Q}}$ -formula

$$F : \exists x. 2x = y,$$

quantifier-free  $T_{\mathbb{Q}}$ -equivalent  $\Sigma_{\mathbb{Q}}$ -formula is

$$G : \top$$

For  $\Sigma_{\mathbb{Z}}$ -formula

$$F : \exists x. 2x = y,$$

there is no quantifier-free  $T_{\mathbb{Z}}$ -equivalent  $\Sigma_{\mathbb{Z}}$ -formula.

Let  $\widehat{T}_{\mathbb{Z}}$  be  $T_{\mathbb{Z}}$  with divisibility predicates  $|$ .

For  $\widehat{\Sigma}_{\mathbb{Z}}$ -formula

$$F : \exists x. 2x = y,$$

a quantifier-free  $\widehat{T}_{\mathbb{Z}}$ -equivalent  $\widehat{\Sigma}_{\mathbb{Z}}$ -formula is

$$G : 2 \mid y.$$

## About QE Algorithm

In developing a QE algorithm for theory  $T$ , we need only consider formulae of the form

$$\exists x. F$$

for quantifier-free  $F$ .

Example: For  $\Sigma$ -formula

$$G_1 : \exists x. \forall y. \underbrace{\exists z. F_1[x, y, z]}_{F_2[x, y]}$$

$$G_2 : \exists x. \forall y. F_2[x, y]$$

$$G_3 : \exists x. \underbrace{\neg \exists y. \neg F_2[x, y]}_{F_3[x]}$$

$$G_4 : \underbrace{\exists x. \neg F_3[x]}_{F_4}$$

$$G_5 : F_4$$

$G_5$  is quantifier-free and  $T$ -equivalent to  $G_1$  

## Quantifier Elimination for $T_{\mathbb{Z}}$

$\Sigma_{\mathbb{Z}} : \{\dots, -2, -1, 0, 1, 2, \dots, -3\cdot, -2\cdot, 2\cdot, 3\cdot, \dots, +, -, =, <\}$

Lemma:

Given quantifier-free  $\Sigma_{\mathbb{Z}}$ -formula  $F[y]$  s.t.  $\text{free}(F[y]) = \{y\}$ .

$F[t]$  represents the set of integers

$$S : \{n \in \mathbb{Z} : F[n] \text{ is } T_{\mathbb{Z}}\text{-valid}\} .$$

Either  $S \cap \mathbb{Z}^+$  or  $\mathbb{Z}^+ \setminus S$  is finite.

Note:  $\mathbb{Z}^+$  is the set of positive integers.

Example:  $\Sigma_{\mathbb{Z}}$ -formula  $F[y] : \exists x. 2x = y$

$S$ : even integers

$S \cap \mathbb{Z}^+$ : positive even integers — infinite

$\mathbb{Z}^+ \setminus S$ : positive odd integers — infinite

Therefore, by the lemma, there is no quantifier-free  $T_{\mathbb{Z}}$ -formula that is  $T_{\mathbb{Z}}$ -equivalent to  $F[y]$ .

Thus,  $T_{\mathbb{Z}}$  does not admit QE.

## Augmented theory $\widehat{T}_{\mathbb{Z}}$

$\widehat{\Sigma}_{\mathbb{Z}}$ :  $\Sigma_{\mathbb{Z}}$  with countable number of unary divisibility predicates  
 $k \mid \cdot$  for  $k \in \mathbb{Z}^+$

Intended interpretations:

$k \mid x$  holds iff  $k$  divides  $x$  without any remainder

Example:

$$x > 1 \wedge y > 1 \wedge 2 \mid x + y$$

is satisfiable (choose  $x = 2, y = 2$ ).

$$\neg(2 \mid x) \wedge 4 \mid x$$

is not satisfiable.

Axioms of  $\widehat{T}_{\mathbb{Z}}$ : axioms of  $T_{\mathbb{Z}}$  with additional countable set of axioms

$$\forall x. k \mid x \leftrightarrow \exists y. x = ky \quad \text{for } k \in \mathbb{Z}^+$$

## $\widehat{T}_{\mathbb{Z}}$ admits QE (Cooper's method)

Algorithm: Given  $\widehat{\Sigma}_{\mathbb{Z}}$ -formula

$$\exists x. F[x],$$

where  $F$  is quantifier-free, construct quantifier-free  $\widehat{\Sigma}_{\mathbb{Z}}$ -formula that is equivalent to  $\exists x. F[x]$ .

1. Put  $F[x]$  into Negation Normal Form (NNF).
2. Normalize literals:  $s < t$ ,  $k|t$ , or  $\neg(k|t)$ .
3. Put  $x$  in  $s < t$  on one side:  $hx < t$  or  $s < hx$ .
4. Replace  $hx$  with  $x'$  without a factor.
5. Replace  $F[x']$  by  $\bigvee F[j]$  for finitely many  $j$ .

## Cooper's Method: Step 1

Put  $F[x]$  in Negation Normal Form (NNF)  $F_1[x]$ , so that  $\exists x. F_1[x]$

- ▶ has negations only in literals (only  $\wedge$ ,  $\vee$ )
- ▶ is  $\widehat{T}_{\mathbb{Z}}$ -equivalent to  $\exists x. F[x]$

Example:

$$\exists x. \neg(x - 6 < z - x \wedge 4 \mid 5x + 1 \rightarrow 3x < y)$$

is equivalent to

$$\exists x. x - 6 < z - x \wedge 4 \mid 5x + 1 \wedge \neg(3x < y)$$

## Cooper's Method: Step 2

Replace (left to right)

$$s = t \Leftrightarrow s < t + 1 \wedge t < s + 1$$

$$\neg(s = t) \Leftrightarrow s < t \vee t < s$$

$$\neg(s < t) \Leftrightarrow t < s + 1$$

The output  $\exists x. F_2[x]$  contains only literals of form

$$s < t, \quad k \mid t, \quad \text{or} \quad \neg(k \mid t),$$

where  $s, t$  are  $\widehat{T}_{\mathbb{Z}}$ -terms and  $k \in \mathbb{Z}^+$ .

Example:

$$\neg(x < y) \wedge \neg(x = y + 3)$$

$\Downarrow$

$$y < x + 1 \wedge (x < y + 3 \vee y + 3 < x)$$

## Cooper's Method: Step 3

Collect terms containing  $x$  so that literals have the form

$$hx < t, \quad t < hx, \quad k \mid hx + t, \quad \text{or} \quad \neg(k \mid hx + t),$$

where  $t$  is a term (does not contain  $x$ ) and  $h, k \in \mathbb{Z}^+$ . The output is the formula  $\exists x. F_3[x]$ , which is  $\widehat{T}_{\mathbb{Z}}$ -equivalent to  $\exists x. F[x]$ .

Example:

$$x + x + y < z + 3z + 2y - 4x$$

$$\Downarrow$$

$$6x < 4z + y$$

$$5 \mid -7x + t$$

$$\Downarrow$$

$$5 \mid 7x - t$$

## Cooper's Method: Step 4 I

Let

$$\delta' = \text{lcm}\{h : h \text{ is a coefficient of } x \text{ in } F_3[x]\},$$

where lcm is the least common multiple. Multiply atoms in  $F_3[x]$  by constants so that  $\delta'$  is the coefficient of  $x$  everywhere:

$$hx < t \Leftrightarrow \delta'x < h't \quad \text{where } h'h = \delta'$$

$$t < hx \Leftrightarrow h't < \delta'x \quad \text{where } h'h = \delta'$$

$$k \mid hx + t \Leftrightarrow h'k \mid \delta'x + h't \quad \text{where } h'h = \delta'$$

$$\neg(k \mid hx + t) \Leftrightarrow \neg(h'k \mid \delta'x + h't) \quad \text{where } h'h = \delta'$$

The result  $\exists x. F'_3[x]$ , in which all occurrences of  $x$  in  $F'_3[x]$  are in terms  $\delta'x$ .

Replace  $\delta'x$  terms in  $F'_3$  with a fresh variable  $x'$  to form

$$F''_3 : F_3\{\delta'x \mapsto x'\}$$

## Cooper's Method: Step 4 II

Finally, construct

$$\exists x'. \underbrace{F_3''[x'] \wedge \delta' \mid x'}_{F_4[x']}$$

$\exists x'. F_4[x']$  is equivalent to  $\exists x. F[x]$  and each literal of  $F_4[x']$  has one of the forms:

- (A)  $x' < t$
- (B)  $t < x'$
- (C)  $k \mid x' + t$
- (D)  $\neg(k \mid x' + t)$

where  $t$  is a term that does not contain  $x$ , and  $k \in \mathbb{Z}^+$ .

## Cooper's Method: Step 4 III

Example:  $\widehat{T}_{\mathbb{Z}}$ -formula

$$\exists x. \underbrace{3x + 1 > y \wedge 2x - 6 < z \wedge 4 \mid 5x + 1}_{F[x]}$$

After step 3:

$$\exists x. \underbrace{2x < z + 6 \wedge y - 1 < 3x \wedge 4 \mid 5x + 1}_{F_3[x]}$$

Collecting coefficients of  $x$  (step 4):

$$\delta' = \text{lcm}(2, 3, 5) = 30$$

Multiply when necessary:

$$\exists x. 30x < 15z + 90 \wedge 10y - 10 < 30x \wedge 24 \mid 30x + 6$$

## Cooper's Method: Step 4 IV

Multiply when necessary:

$$\exists x. 30x < 15z + 90 \wedge 10y - 10 < 30x \wedge 24 \mid 30x + 6$$

Replacing  $30x$  with fresh  $x'$  and adding divisibility conjunct:

$$\exists x'. \underbrace{x' < 15z + 90 \wedge 10y - 10 < x' \wedge 24 \mid x' + 6 \wedge 30 \mid x'}_{F_4[x']}$$

$\exists x'. F_4[x']$  is equivalent to  $\exists x. F[x]$ .

## Cooper's Method: Step 5

Construct left infinite projection  $F_{-\infty}[x']$  of  $F_4[x']$  by

(A) replacing literals  $x' < t$  by  $\top$

(B) replacing literals  $t < x'$  by  $\perp$

Idea: very small numbers satisfy (A) literals but not (B) literals

Let

$$\delta = \text{lcm} \left\{ \begin{array}{l} k \text{ of (C) literals } k \mid x' + t \\ k \text{ of (D) literals } \neg(k \mid x' + t) \end{array} \right\}$$

and  $B$  be the set of terms  $t$  appearing in (B) literals of  $F_4[x']$ .

Construct

$$F_5 : \bigvee_{j=1}^{\delta} F_{-\infty}[j] \vee \bigvee_{j=1}^{\delta} \bigvee_{t \in B} F_4[t + j].$$

$F_5$  is quantifier-free and  $\widehat{T}_{\mathbb{Z}}$ -equivalent to  $\exists x. F[x]$ .

## Intuition of Step 5 I

### Property (Periodicity)

if  $m \mid \delta$

then  $m \mid n$  iff  $m \mid n + \lambda\delta$  for all  $\lambda \in \mathbb{Z}$

That is,  $m \mid \cdot$  cannot distinguish between  $m \mid n$  and  $m \mid n + \lambda\delta$ .

By the choice of  $\delta$  (lcm of the  $k$ 's) — no  $\mid$  literal in  $F_5$  can distinguish between  $n$  and  $n + \lambda\delta$ , for any  $\lambda \in \mathbb{Z}$ .

$$F_5 : \bigvee_{j=1}^{\delta} F_{-\infty}[j] \vee \bigvee_{j=1}^{\delta} \bigvee_{t \in B} F_4[t + j]$$

## Intuition of Step 5 II

- ▶ left disjunct  $\bigvee_{j=1}^{\delta} F_{-\infty}[j]$  :

Contains only | literals

Asserts: no least  $n \in \mathbb{Z}$  s.t.  $F_4[n]$ .

For if there exists  $n$  satisfying  $F_{-\infty}$ ,

then every  $n - \lambda\delta$ , for  $\lambda \in \mathbb{Z}^+$ , also satisfies  $F_{-\infty}$

- ▶ right disjunct  $\bigvee_{j=1}^{\delta} \bigvee_{t \in B} F_4[t + j]$  :

Asserts: There is least  $n \in \mathbb{Z}$  s.t.  $F_4[n]$ .

For let  $t^*$  be the largest  $t$  in  $(B) = \{\text{largest } t \mid t < x' \text{ in } (B)\}$ .

If  $n \in \mathbb{Z}$  is s.t.  $F_4[n]$ , then

$$\exists j(1 \leq j \leq \delta). t^* + j \leq n \wedge F_4[t^* + j]$$

In other words,

if there is a solution,

then one must appear in  $\delta$  interval to the right of  $t^*$

## Example of Step 5 I

$$\exists x. \underbrace{3x + 1 > y \wedge 2x - 6 < z \wedge 4 \mid 5x + 1}_{F[x]}$$

$\Downarrow$

$$\exists x'. \underbrace{x' < 15z + 90 \wedge 10y - 10 < x' \wedge 24 \mid x' + 6 \wedge 30 \mid x'}_{F_4[x']}$$

By step 5,

$$F_{-\infty}[x] : \top \wedge \perp \wedge 24 \mid x' + 6 \wedge 30 \mid x' ,$$

which simplifies to  $\perp$ .

## Example of Step 5 II

Compute

$$\delta = \text{lcm}\{24, 30\} = 120 \quad \text{and} \quad B = \{10y - 10\}.$$

Then replacing  $x'$  by  $10y - 10 + j$  in  $F_4[x']$  produces

$$F_5 : \bigvee_{j=1}^{120} \left[ \begin{array}{l} 10y - 10 + j < 15z + 90 \wedge 10y - 10 < 10y - 10 + j \\ \wedge 24 \mid 10y - 10 + j + 6 \wedge 30 \mid 10y - 10 + j \end{array} \right]$$

which simplifies to

$$F_5 : \bigvee_{j=1}^{120} \left[ \begin{array}{l} 10y + j < 15z + 100 \wedge 0 < j \\ \wedge 24 \mid 10y + j - 4 \wedge 30 \mid 10y - 10 + j \end{array} \right].$$

$F_5$  is quantifier-free and  $\widehat{T}_{\mathbb{Z}}$ -equivalent to  $\exists x. F[x]$ .

## Cooper's Method: Example I

$$\underbrace{\exists x. (3x + 1 < 10 \vee 7x - 6 > 7) \wedge 2 \mid x}_{F[x]}$$

Isolate  $x$  terms

$$\exists x. (3x < 9 \vee 13 < 7x) \wedge 2 \mid x ,$$

so

$$\delta' = \text{lcm}\{3, 7\} = 21 .$$

After multiplying coefficients by proper constants,

$$\exists x. (21x < 63 \vee 39 < 21x) \wedge 42 \mid 21x ,$$

we replace  $21x$  by  $x'$ :

$$\exists x'. \underbrace{(x' < 63 \vee 39 < x')}_{F_4[x']} \wedge 42 \mid x' \wedge 21 \mid x' .$$

## Cooper's Method: Example II

Then

$$F_{-\infty}[x'] : (\top \vee \perp) \wedge 42 \mid x' \wedge 21 \mid x' ,$$

or, simplifying,

$$F_{-\infty}[x'] : 42 \mid x' \wedge 21 \mid x' .$$

Finally,

$$\delta = \text{lcm}\{21, 42\} = 42 \quad \text{and} \quad B = \{39\} ,$$

so  $F_5$  :

$$\bigvee_{j=1}^{42} (42 \mid j \wedge 21 \mid j) \vee \bigvee_{j=1}^{42} ((39 + j < 63 \vee 39 < 39 + j) \wedge 42 \mid 39 + j \wedge 21 \mid 39 + j) .$$

Since  $42 \mid 42$  and  $21 \mid 42$ , the left main disjunct simplifies to  $\top$ , so that  $F_5$  is  $\widehat{T}_{\mathbb{Z}}$ -equivalent to  $\top$ . Thus,  $\exists x. F[x]$  is  $\widehat{T}_{\mathbb{Z}}$ -valid.

## Cooper's Method: Example I

$$\exists x. \underbrace{2x = y}_{F[x]}$$

Rewriting

$$\exists x. \underbrace{y - 1 < 2x \wedge 2x < y + 1}_{F_3[x]}$$

Then

$$\delta' = \text{lcm}\{2, 2\} = 2,$$

so by Step 4

$$\exists x'. \underbrace{y - 1 < x' \wedge x' < y + 1 \wedge 2 \mid x'}_{F_4[x']}$$

$F_{-\infty}$  produces  $\perp$ .

## Cooper's Method: Example II

However,

$$\delta = \text{lcm}\{2\} = 2 \quad \text{and} \quad B = \{y - 1\},$$

so

$$F_5 : \bigvee_{j=1}^2 (y - 1 < y - 1 + j \wedge y - 1 + j < y + 1 \wedge 2 \mid y - 1 + j)$$

Simplifying,

$$F_5 : \bigvee_{j=1}^2 (0 < j \wedge j < 2 \wedge 2 \mid y - 1 + j)$$

and then

$$F_5 : 2 \mid y,$$

which is quantifier-free and  $\widehat{T}_{\mathbb{Z}}$ -equivalent to  $\exists x. F[x]$ .

## Improvement: Symmetric Elimination

In step 5, if there are fewer

(A) literals  $x' < t$

than

(B) literals  $t < x'$ ,

construct the right infinite projection  $F_{+\infty}[x']$  from  $F_4[x']$  by replacing

(A) literal  $x' < t$  by  $\perp$

than

(B) literal  $t < x'$  by  $\top$

Then right elimination.

$$F_5 : \bigvee_{j=1}^{\delta} F_{+\infty}[-j] \vee \bigvee_{j=1}^{\delta} \bigvee_{t \in A} F_4[t - j].$$

## Improvement: Eliminating Blocks of Quantifiers I

Given

$$\exists x_1. \dots \exists x_n. F[x_1, \dots, x_n]$$

where  $F$  quantifier-free.

Eliminating  $x_n$  (left elimination) produces

$$G_1 : \exists x_1. \dots \exists x_{n-1}. \bigvee_{j=1}^{\delta} F_{-\infty}[x_1, \dots, x_{n-1}, j] \vee \bigvee_{j=1}^{\delta} \bigvee_{t \in B} F_4[x_1, \dots, x_{n-1}, t + j]$$

which is equivalent to

$$G_2 : \bigvee_{j=1}^{\delta} \exists x_1. \dots \exists x_{n-1}. F_{-\infty}[x_1, \dots, x_{n-1}, j] \vee \bigvee_{j=1}^{\delta} \bigvee_{t \in B} \exists x_1. \dots \exists x_{n-1}. F_4[x_1, \dots, x_{n-1}, t + j]$$

## Improvement: Eliminating Blocks of Quantifiers II

Treat  $j$  as a free variable and examine only  $1 + |B|$  formulae

- ▶  $\exists x_1. \cdots \exists x_{n-1}. F_{-\infty}[x_1, \dots, x_{n-1}, j]$
- ▶  $\exists x_1. \cdots \exists x_{n-1}. F_4[x_1, \dots, x_{n-1}, t + j]$  for each  $t \in B$

## Example 1

$$F : \exists y. \exists x. x < -2 \wedge 1 - 5y < x \wedge 1 + y < 13x$$

Since  $\delta' = \text{lcm}\{1, 13\} = 13$

$$\exists y. \exists x. 13x < -26 \wedge 13 - 65y < 13x \wedge 1 + y < 13x$$

Then

$$\exists y. \exists x'. x' < -26 \wedge 13 - 65y < x' \wedge 1 + y < x' \wedge 13 \mid x'$$

There is one (A) literal  $x' < \dots$  and two (B) literals  $\dots < x'$ , we use right elimination.

$$F_{+\infty} = \perp \quad \delta = \{13\} = 13 \quad A = \{-26\}$$

$$\exists y. \bigvee_{j=1}^{13} \left[ \begin{array}{l} -26 - j < -26 \wedge 13 - 65y < -26 - j \\ \wedge 1 + y < -26 - j \wedge 13 \mid -26 - j \end{array} \right]$$

## Example II

Commute

$$G[j] : \bigvee_{j=1}^{13} \exists y. j > 0 \wedge 39 + j < 65y \wedge y < -27 - j \wedge 13 \mid -26 - j$$

Treating  $j$  as free variable (and removing  $j > 0$ ), apply QE to

$$H[j] : \exists y. 39 + j < 65y \wedge y < -27 - j \wedge 13 \mid -26 - j$$

Simplify...

$$H'[j] : \bigvee_{k=1}^{65} (k < -1794 - 66j \wedge 13 \mid -26 - j \wedge 65 \mid 39 + j + k)$$

Replace  $H[j]$  with  $H'[j]$  in  $G[j]$

$$\bigvee_{j=1}^{13} \bigvee_{k=1}^{65} (k < -1794 - 66j \wedge 13 \mid -26 - j \wedge 65 \mid 39 + j + k)$$

## Example III

This qff formula is  $\widehat{T}_{\mathbb{Z}}$ -equivalent to  $F$ .

# Quantifier Elimination over Rationals

$$\Sigma_{\mathbb{Q}} : \{0, 1, +, -, =, \geq\}$$

Recall: we use  $>$  instead of  $\geq$ , as

$$x \geq y \Leftrightarrow x > y \vee x = y \quad x > y \Leftrightarrow x \geq y \wedge \neg(x = y).$$

## Ferrante & Rackoff's Method

Given a  $\Sigma_{\mathbb{Q}}$ -formula  $\exists x. F[x]$ , where  $F[x]$  is quantifier-free, generate quantifier-free formula  $F_4$  (four steps) s.t.

$$F_4 \text{ is } \Sigma_{\mathbb{Q}}\text{-equivalent to } \exists x. F[x]$$

by

1. putting  $F[x]$  in NNF,
2. replacing negated literals,
3. solving literals s.t.  $x$  appears isolated on one side, and
4. taking finite disjunction  $\bigvee_t F[t]$ .

## Ferrante & Rackoff's Method: Steps 1 and 2

Step 1: Put  $F[x]$  in NNF. The result is  $\exists x. F_1[x]$ .

Step 2: Replace literals (left to right)

$$\neg(s < t) \Leftrightarrow t < s \vee t = s$$

$$\neg(s = t) \Leftrightarrow t < s \vee t > s$$

The result  $\exists x. F_2[x]$  does not contain negations.

## Ferrante & Rackoff's Method: Step 3

Solve for  $x$  in each atom of  $F_2[x]$ , e.g.,

$$t_1 < cx + t_2 \quad \Rightarrow \quad \frac{t_1 - t_2}{c} < x$$

where  $c \in \mathbb{Z} - \{0\}$ .

All atoms in the result  $\exists x. F_3[x]$  have form

(A)  $x < t$

(B)  $t < x$

(C)  $x = t$

where  $t$  is a term that does not contain  $x$ .

## Ferrante & Rackoff's Method: Step 4 I

Construct from  $F_3[x]$

▶ left infinite projection  $F_{-\infty}$  by replacing

(A) atoms  $x < t$  by  $\top$

(B) atoms  $t < x$  by  $\perp$

(C) atoms  $x = t$  by  $\perp$

▶ right infinite projection  $F_{+\infty}$  by replacing

(A) atoms  $x < t$  by  $\perp$

(B) atoms  $t < x$  by  $\top$

(C) atoms  $x = t$  by  $\perp$

Let  $S$  be the set of  $t$  terms from (A), (B), (C) atoms.

Construct the final

$$F_4 : F_{-\infty} \vee F_{+\infty} \vee \bigvee_{s,t \in S} F_3 \left[ \frac{s+t}{2} \right],$$

which is  $T_{\mathbb{Q}}$ -equivalent to  $\exists x. F[x]$ .

## Ferrante & Rackoff's Method: Step 4 II

- ▶  $F_{-\infty}$  captures the case when small  $x \in \mathbb{Q}$  satisfy  $F_3[x]$
- ▶  $F_{+\infty}$  captures the case when large  $x \in \mathbb{Q}$  satisfy  $F_3[x]$
- ▶ last disjunct: for  $s, t \in S$ 
  - if  $s \equiv t$ , check whether  $s \in S$  satisfies  $F_3[s]$
  - if  $s \not\equiv t$ , in any  $T_{\mathbb{Q}}$ -interpretation,
    - ▶  $|S| - 1$  pairs  $s, t \in S$  are adjacent. For each such pair,  $(s, t)$  is an interval in which no other  $s' \in S$  lies.
    - ▶ Since  $\frac{s+t}{2}$  represents the whole interval  $(s, t)$ , simply check  $F_3[\frac{s+t}{2}]$ .

## Ferrante & Rackoff's Method: Intuition

Step 4 says that four cases are possible:

1. There is a left open interval s.t. all elements satisfy  $F(x)$ .

$$\overline{\leftarrow)}$$

2. There is a right open interval s.t. all elements satisfy  $F(x)$ .

$$\overline{\leftarrow)} \quad (\rightarrow)$$

3. Some term  $t$  satisfies  $F(x)$ .

$$\overline{\dots \quad t \quad \dots}$$

4. There is an open interval between two  $s, t$  terms s.t. every element satisfies  $F(x)$ .

$$\overline{(\leftarrow \rightarrow)}$$
$$\dots \quad s \quad \uparrow \quad t \quad \dots$$
$$\frac{s+t}{2}$$

## Correctness of Step 4 I

### Theorem

Let  $F_4$  be the formula constructed from  $\exists x. F_3[x]$  as in Step 4.  
Then  $\exists x. F_3[x] \Leftrightarrow F_4$ .

### Proof:

- $\Leftarrow$  If  $F_4$  is true, then  $F_{-\infty}$ ,  $F_{\infty}$  or  $F_3[\frac{s+t}{2}]$  is true.
  - If  $F_3[\frac{s+t}{2}]$  is true, then obviously  $\exists x. F_3[x]$  is true.
  - If  $F_{-\infty}$  is true, choose some small  $x, x < t$  for all  $t \in S$ .  
Then  $F_3[x]$  is true.
  - If  $F_{+\infty}$  is true, choose some big  $x, x > t$  for all  $t \in S$ .  
Then  $F_3[x]$  is true.

## Correctness of Step 4 II

⇒ If  $I \models \exists x. F_3[x]$  then there is value  $v$  such that

$$I \triangleleft \{x \mapsto v\} \models F_3.$$

If  $v < \alpha_I[t]$  for all  $t \in S$ , then  $I \models F_{-\infty}$ .

If  $v > \alpha_I[t]$  for all  $t \in S$ , then  $I \models F_{+\infty}$ .

If  $v = \alpha_I[t]$  for some  $t \in S$ , then  $I \models F[\frac{t+t}{2}]$ .

Otherwise choose largest  $s \in S$  with  $\alpha_I[s] < v$  and smallest  $t \in S$  with  $\alpha_I[t] > v$ .

Since no atom of  $F_3$  can distinguish between values in interval  $(s, t)$ ,

$$I \models F_3[v] \text{ iff } I \models F_3\left[\frac{s+t}{2}\right].$$

Hence,  $I \models F[\frac{s+t}{2}]$ .

## Correctness of Step 4 III

In all cases  $I \models F_4$ .

# Ferrante & Rackoff's Method: Example I

$\Sigma_{\mathbb{Q}}$ -formula

$$\exists x. \underbrace{3x + 1 < 10 \wedge 7x - 6 > 7}_{F[x]}$$

Solving for  $x$

$$\exists x. \underbrace{x < 3 \wedge x > \frac{13}{7}}_{F_3[x]}$$

$$\begin{array}{ll} \text{Step 4: } x < 3 \text{ in (A)} & \Rightarrow F_{-\infty} = \perp \\ x > \frac{13}{7} \text{ in (B)} & \Rightarrow F_{+\infty} = \perp \end{array}$$

$$F_4 : \bigvee_{s,t \in S} \underbrace{\left( \frac{s+t}{2} < 3 \wedge \frac{s+t}{2} > \frac{13}{7} \right)}_{F_3[\frac{s+t}{2}]}$$

## Ferrante & Rackoff's Method: Example II

$$S = \left\{3, \frac{13}{7}\right\} \Rightarrow$$

$$F_3 \left[ \frac{3+3}{2} \right] = \perp \quad F_3 \left[ \frac{\frac{13}{7} + \frac{13}{7}}{2} \right] = \perp$$

$$F_3 \left[ \frac{\frac{13}{7} + 3}{2} \right] : \frac{\frac{13}{7} + 3}{2} < 3 \wedge \frac{\frac{13}{7} + 3}{2} > \frac{13}{7}$$

simplifies to  $\top$ .

Thus,  $F_4 : \top$  is  $T_{\mathbb{Q}}$ -equivalent to  $\exists x. F[x]$ ,  
so  $\exists x. F[x]$  is  $T_{\mathbb{Q}}$ -valid.

## Example

$$\exists x. \underbrace{2x > y \wedge 3x < z}_{F[x]}$$

Solving for  $x$

$$\exists x. \underbrace{x > \frac{y}{2} \wedge x < \frac{z}{3}}_{F_3[x]}$$

Step 4:  $F_{-\infty} = \perp$ ,  $F_{+\infty} = \perp$ ,  $F_3[\frac{y}{2}] = \perp$  and  $F_3[\frac{z}{3}] = \perp$ .

$$F_4 : \frac{\frac{y}{2} + \frac{z}{3}}{2} > \frac{y}{2} \wedge \frac{\frac{y}{2} + \frac{z}{3}}{2} < \frac{z}{3}$$

which simplifies to:

$$F_4 : 2z > 3y$$

$F_4$  is  $T_{\mathbb{Q}}$ -equivalent to  $\exists x. F[x]$ .