

Class 7: Agenda and Questions

1 Warm-Up

Group Work

Let $G = (V, E)$ be a weighted, undirected graph, on n vertices with edge weights w_{uv} on the edge $\{u, v\} \in E$. Let $d : V \times V \rightarrow \mathbb{R}$ be the associated graph metric.

Explain how to efficiently find and apply a map $f : V \rightarrow \mathbb{R}^k$, for $k = O(\log^2 n)$, so that

$$\frac{\sum_{\{u,v\} \in E} \|f(u) - f(v)\|_1}{\sum_{\{u,v\} \in \binom{V}{2}} \|f(u) - f(v)\|_1} \leq O(\log n) \frac{\sum_{\{u,v\} \in E} d(u, v)}{\sum_{\{u,v\} \in \binom{V}{2}} d(u, v)}$$

holds with high probability. Above, $\binom{V}{2}$ refers to the set of all unordered pairs $\{u, v\}$ for $u, v \in V$ and $u \neq v$.

2 Lecture Recap and Questions?

Any questions from the mini-lectures or pre-class-quiz? (Metric Embeddings; Bourgain's Embedding)

3 Sparsest Cuts

[Some slides; summary is below]

For a graph $G = (V, E)$, define

$$\phi(G, S) = \frac{|E(S, \bar{S})|}{|S||\bar{S}|},$$

and

$$\phi(G) = \min_{S \subset V, S \neq \emptyset, V} \phi(G, S),$$

where above $\bar{S} := V \setminus S$ denotes the complement of S , and $E(S, \bar{S})$ denotes the set of edges that have one endpoint in S and one endpoint in \bar{S} .

Intuitively, if $\phi(G, S)$ is small, then S is pretty “disconnected” from \bar{S} . Notice that the denominator, $|S||\bar{S}|$, is the number of edges that would be between S and \bar{S} in the complete graph, so $\phi(G, S)$ is the fraction of possible edges between S and \bar{S} that actually exist in G .

Finding S to minimize $\phi(G, S)$ is useful, for example, in clustering applications. However, it's also NP-hard. Today we'll see a randomized algorithm to find an S so that $\phi(G, S)$ is *approximately* minimized. More precisely, we'll find S so that $\phi(S, G) \leq O(\log n)\phi(G)$.

Question: How is this definition of $\phi(G)$ different than simply asking for the minimum cut? When might you prefer a sparsest cut to a min cut? (Recall we saw a randomized algorithm for the minimum cut back in Week 1...)

3.1 Connection to metrics

Group Work

In this group work, you will show that

$$\phi(G) = \min_f \frac{\sum_{\{u,v\} \in E} \|f(u) - f(v)\|_1}{\sum_{\{u,v\} \in \binom{V}{2}} \|f(u) - f(v)\|_1}, \quad (1)$$

where the minimum is over all functions $f : V \rightarrow \mathbb{R}^k$ for some k , so that f takes on at least two distinct values. (This last bit is needed so that the denominator doesn't vanish).

1. Show that

$$\phi(G) = \min_{f:V \rightarrow \{0,1\}} \frac{\sum_{\{u,v\} \in E} |f(u) - f(v)|}{\sum_{\{u,v\} \in \binom{V}{2}} |f(u) - f(v)|},$$

where the minimum is over all functions $f : V \rightarrow \{0, 1\}$ so that f takes on both values 0 and 1. (The difference between this and the expression above is that f maps to $\{0, 1\}$ instead of \mathbb{R}^k for some k).

Hint: What's a natural way to relate a set $S \subseteq V$ to a function $f : V \rightarrow \{0, 1\}$?

The next two parts are optional. If you have time, just try to get some intuition.

2. Think about why the above extends to show that

$$\phi(G) = \inf_{f:V \rightarrow \mathbb{R}} \frac{\sum_{\{u,v\} \in E} |f(u) - f(v)|}{\sum_{\{u,v\} \in \binom{V}{2}} |f(u) - f(v)|},$$

where now the minimum is over $f : V \rightarrow \mathbb{R}$ instead of $f : V \rightarrow \{0, 1\}$.

Hint: Suppose that f takes on three values, say $a < b < c$. What happens if you "slide" the value b closer to a or c in the ratio $R(f) = \frac{\sum_{\{u,v\} \in E} |f(u) - f(v)|}{\sum_{\{u,v\} \in \binom{V}{2}} |f(u) - f(v)|}$? Show

that $R(f)$ always either increases or decreases if you do this, and conclude that the f that minimizes $R(f)$ actually takes on only two values. Then argue that those two values might as well be 0 and 1.

3. Think about why the above extends to show that

$$\phi(G) = \min_{f:V \rightarrow \mathbb{R}^k} \frac{\sum_{\{u,v\} \in E} \|f(u) - f(v)\|_1}{\sum_{\{u,v\} \in \binom{V}{2}} \|f(u) - f(v)\|_1},$$

where the minimum is over all functions $f : V \rightarrow \mathbb{R}^k$ for any k .

Hint: You may want to use the inequality that $\frac{\sum_i a_i}{\sum_i b_i} \geq \min_i \frac{a_i}{b_i}$ for $a_i, b_i > 0$.

3.2 A randomized algorithm

Based on the result that we got in the first group work, we might think of the following approach:

Find $f : V \rightarrow \mathbb{R}^k$ to minimize

$$R(f) := \frac{\sum_{\{u,v\} \in E} \|f(u) - f(v)\|_1}{\sum_{\{u,v\} \in \binom{V}{2}} \|f(u) - f(v)\|_1}$$

Unfortunately, this doesn't turn out to be an easy optimization problem to solve. Instead, we'll consider the optimization problem:

Find values $d_{u,v} \in \mathbb{R}$ for all $u \neq v \in V$ to minimize

$$Q(d) := \sum_{\{u,v\} \in E} d_{u,v}$$

subject to:

- $d_{u,v} = d_{v,u} \geq 0$ for all u, v
- $d_{u,v} + d_{v,w} \geq d_{u,w}$ for all u, v, w
- $\sum_{\{u,v\} \in \binom{V}{2}} d_{u,v} = 1$

It turns out that this problem *can* be solved efficiently, using linear programming. (If you don't know what that is, it's okay, all that matters now is that we can find \vec{d} to minimize this efficiently).

Below, we'll show how to use this to find an efficient approximate sparsest-cut algorithm.

Group Work

1. Suppose that d^* is the minimizer of the problem above.

Explain why $Q(d^*) \leq \phi(G)$.

Hint: Find some d so that $Q(d) = \Phi(G)$, and observe that $Q(d^*) \leq Q(d)$.

2. Find a randomized algorithm to approximate $\phi(G)$. More precisely, give a random-

ized algorithm that finds $f : V \rightarrow \mathbb{R}^k$ so that, with high probability,

$$\frac{\sum_{\{u,v\} \in E} \|f(u) - f(v)\|_1}{\sum_{\{u,v\} \in \binom{V}{2}} \|f(u) - f(v)\|_1} \leq O(\log n)\phi(G).$$

Hint: Your warm-up exercise might be relevant.

Hint: If it comes up, you may assume that Bourgain's embedding works just fine on pseudo-metrics, which are functions $d(u, v)$ that obey all of the axioms of metrics except that maybe $d(u, v) = 0$ for $u \neq v$.

3. Given f as in the previous part, explain how to efficiently find a set $S \subset V$ so that

$$\phi(G, S) \leq O(\log n)\phi(G).$$

Hint: Our approach in the first group-work was somewhat algorithmic...