



Observations over the last 2 days

Models:

Many types and “classes” but are they really that different?

For climate work they must be Process driven – otherwise




Brian's Questions


- How sensitive are LULC to the model of decision making
 - That depends on how the land cover information is passed from one model to the next. Errors in representation/translation may be greater than the signal modeled
 - Difficult to link cause and effect if the LULC model is not process based

PCM Present Day Uncertainty


a) LSM original present day land cover: LSMIc




b) IMAGE 2.2 present day land cover: IMAIc



c) Hybrid present day land cover: HYBIc



d) IMAGE 2.2 potential vegetation land cover: POTIc



LSM Land Cover Types

	0 - Ocean		8 - Broadleaf decid		15 - Forest crop		22 - Semi-Desert
	1 - Ice		10 - Tropical broadleaf		17 - Cool grassland/steppe		26 - Crop
	2 - Desert		11 - Trop seasonal decid tree		18 - Warm grassland		27 - Forest wetland
	3 - Needleleaf evergreen		12 - Savanna		19 - Tundra		28 - Non-forest wetland
	4 - Needleleaf decid		13 - Evergreen forest tundra		20 - Evergreen shrub		
	6 - Temp mixed forest		14 - Decid forest tundra		21 - Decid Shrub		




PCM Uncertainty v.s. Historical Land Cover

PRESENT DAY UNCERTAINTY

- Arctic – albedo
- Amazon – latent heat flux
- Australia – albedo



Present Day (IMAGE) Land Surface minus control




HISTORICAL CHANGE

Climate difference from land cover classification is as large as the climate difference from land cover change

- Primarily shift due to agriculture



Natural Vegetation minus Present Day (IMAGE)





Brian's Questions

- Are IAMs missing important processes
 - Yes
 - Details about land use and land type
 - Inconsistency between base scenarios for IAMs vs other models
 - Simplified land cover representation lead to interpolation/interpretation errors



Brian's Questions

- How to connect regional land use scenarios and data to global scenarios?
 - Splicing of dataset
 - Boundary conditions
 - Integration of data




Brian's Questions

- **How sensitive are climate and biogeochem to LULC**
 - Models are potentially very sensitive to some variables but much less so to others. We need to test the variables that are most critical




Year 2000 Crop comparison


HYDE 3.0




% Grid Cell




Ramankutty 2000



Difference




% Difference






Year 2000 Grazing comparison


HYDE 3.0




% Grid Cell




Ramankutty 2000



Difference




% Difference






Urban land cover


HYDE 3.0




% Grid Cell




Jackson and Feddema 2004



Difference




% Difference






Comparison of Discover and MODIS Bare Ground Fraction


Discover - CLM 3.0



MODIS




Percent Bare




Source: Peter Lawrence, CIRES, University of Colorado

Comparison of Agriculture land classes from 3 satellite products 10 degree tile over East Africa


MODIS




IGBP



GLC2000



Agreement



- No Ag
- 1 product
- 2 products
- All products



Brian's Questions



- **Can validation be improved**
 - This process has hardly started!
 - IAM MIP of 20th century
 - Validation of source data between models




High Resolution Modeling – North East US

From:
Peter Lawrence


NCEP Summer Avg Temp F



NCEP Winter Avg Temp F



NCEP FV1 Sum Avg Temp F



NCEP FV1 Win Avg Temp F

