


Global Warming Levels as an organizing dimension of future climate projections

Stanford-EPRI Public Educational Webcast Series on Options for Estimating the Global Economic Impacts of Climate Change and the Social Cost of Carbon Module-by-Module


Modeling global climate change and earth system responses to greenhouse gas emissions

Claudia Tebaldi
Joint Global Change Research Institute (PNNL)
Claudia.Tebaldi@pnnl.gov


Twenty-year
running
averages:
climate, not
weather!




Twenty-year
running
averages:
climate, not
weather!




Twenty-year
running
averages:
Climate, not
weather!





Twenty-year
running
averages:
Climate, not
weather!



Twenty-year running averages:
Climate, not weather!




Response at different GWLs (Ensemble average)





Timing of GWLs
Depends on scenarios


Twenty-year running averages:
Climate, not weather!



Response at different GWLs (Ensemble average)



Timing of GWLs
Depends on scenarios



Timing 2°C	Best estimate	Very likely range
SSP1-1.9	NA	NA
SSP1-2.6	NA	(2031–50 to NA)
SSP2-4.5	2043-62	(2028–47 to 2075–94)
SSP3-7.0	2037–56	(2026–45 to 2053–72)
SSP5-8.5	2032–51	(2023–42 to 2044–63)




Most atmospheric variables relevant to impact modeling behave in sync with global temperature, i.e., are path/scenario independent when conditioned upon the global warming level. This is the case for both mean and extreme metrics.

Change in temperature of the hottest day of the year

(a) At 1.5°C global warming

(b) At 2.0°C global warming

(c) At 4.0°C global warming








Change in temperature of the coldest day of the year


(d) At 1.5°C global warming

(e) At 2.0°C global warming


(f) At 4.0°C global warming




Change in length of longest dry spell of the year




Change in soil moisture



Change in 10-year drought events (JJA)




Change in 10-year drought events (DJF)




Change in precipitation amount during the wettest day of the year


(a) At 1.5°C global warming




(b) At 2.0°C global warming



(c) At 4.0°C global warming



Colour High model agreement
Hatched Low model agreement





Quantities with long memory do not depend only on the global temperature level reached, but on how the system got there, for how long it has been there -> Sea Level Rise

Non-reversible impacts may look very different on one side or the other of the temperature trajectory peak -> Ice Sheets, Desertification

Impacts may depend on the **rate of change** besides the level of warming.


At which GWL **by 2100** will today's 100-yr event become an annual event?







Land areas where temperature and precipitation show irreversible changes

GWLs and uncertainty: where we ‘clasp’ does not change the sources and relevance, but may change the tools we use to address the various modeling steps.



GWLs and uncertainty: Clasping it does not change the sources and relevance, but may change the tools we use to characterize the various steps.





We still need **scenarios** of future emissions to tell a coherent story of future developments and action/choices, and to add a consistent representation of the human component to the evaluation of impacts.

Getting to 2C by 2050 will have very different impacts than getting to it **by 2090**; In fact getting to 2C by 2090 but along **different socio-economic pathways** will have very different impacts.

Exposure, vulnerability/adaptation are critical determinants and depend on the **scenario, down to its sub-global assumptions.**

Thank you