

A Study of Acoustic Ultrahigh-energy Neutrino Detection (SAUND)

<http://hep.stanford.edu/neutrino/SAUND/>



Justin Vandembroucke

Stanford / Berkeley

justinav@socrates.berkeley.edu

8th International Conference on Advanced Technology and Particle Physics

Como, Italy

October 8, 2003

Thanks to my SAUND collaborators:

Current:

G. Gratta (Stanford)

M. Buckingham (Scripps)

J. Vandenbroucke (Stanford, now at Berkeley)

Past:

N. Lehtinen (Stanford, now at Hawaii)

S. Adam (Stanford, now at Cornell)

Y. Zhao (Stanford)

T. Berger (Scripps)

And to AUTECH and the U. S. Navy:

D. Belasco


J. Cecil

D. Deveau


D. Kapolka

T. Kelly-Bissonnette

Cosmic ray spectrum



The Greisen-Zatsepin-Kuzmin (GZK) cutoff



1 event / km² / century → detector innovation key!


So what?

A crossroads of exciting physics

- active galactic nuclei
- gamma ray bursts
- grand unified theories
- topological defects (magnetic monopoles, cosmic strings, domain walls)
- supersymmetry
- dark matter
- Lorentz invariance violation
- extra dimensions
- gravity at a TeV
- relic neutrinos


All key to understanding the universe!

Cosmic ray showers




http://zebu.uoregon.edu/~js/glossary/cosmic_rays.html

The ocean as a particle detector




Simulated neutrino signal

$\theta=8^\circ$



Extreme pancakes

Radiation lobes from $5e+20$ eV to $3e+21$ eV



Sound from charged particles in liquids: some background

- first discussion of the idea

G. A. Askaryan Sov. J. Atom. Energy 3 (1957) 921

- extensive theoretical analysis

J. G. Learned Phys. Rev. D 19 (1979) 3293

- experimental confirmation

L. Sulak et al. NIM 161 (1979) 203

- sensitivity studies for neutrino detection

N. Lehtinen et al. Astroparticle Physics 17 (2002) 279—292



- growing interest in acoustic arrays

*Stanford workshop on acoustic cosmic ray and neutrino detection,
September 2003 <http://hep.stanford.edu/neutrino/SAUND/workshop>*

The Atlantic Undersea Test and Evaluation Center (AUTEC)




AUTEC hydrophones



SAUND
7 km²

The SAUND array



Site 3



DAQ



8th ICATPP, Como, Italy

Justin Vandembroucke

October 8, 2003

DAQ

Software

- 179 kHz sampling
- digital matched filter
- variable threshold
- 60 events/minute target
- 1-2 GB / 24 hrs



$$\text{signal} : S(t) \propto \frac{t}{\tau} e^{-t^2/2\tau^2}$$

$$\text{noise} : N(t) \propto f^{-2}$$


$$\rightarrow \text{response function} : H(t) \propto \left(\left(\frac{t}{\tau} \right)^3 - 3 \frac{t}{\tau} \right) e^{-t^2/2\tau^2}$$

Hardware


- 1.7 GHz Pentium 4
- ADC card
- 60 GB external hard drive



Adaptive threshold




Calibration sources




One bulb at all 7 hydrophones

Bulb 1, 30-Jul-2001 09:08:01




Two reflections!

Bulb 1, 30-Jul-2001 09:08:01




Direct signal


Bulb 1, 30-Jul-2001 09:08:01



Lightbulb positions reconstructed





Lightbulb positions reconstructed



Lightbulb energies reconstructed

Bulb	Depth (m)	$P = \rho gh$ (kPa)	$E_0 = PV$ (J)
1	160	1563	234
2	107	1047	157
3	139	1360	204
4	166	1626	244
5	126	1237	186
6	101	990	148
7	86	838	126
8	135	1324	199
9	188	1842	276
10	290	2846	427

Data: a large livetime has been integrated




211 days livetime	25 million events	350 GB	70% duty cycle
-------------------	-------------------	--------	----------------

Five-phone coincidence




Require

- 1) Events obey causality: $t_{ij} \leq d_{ij} / v_{sound}$
- 2) Geometry consistent with pancake (2D circle) shape:


accepted:




rejected:




AUTEC sound velocity profile (SVP)




Refraction




Refraction shadow




Refracted pancake




Position reconstruction achieved (10 m resolution)




Localization → energy reconstruction





Background event type 1




Background event type 2



Examples of dolphin signals recorded by AUTECH personnel




Background categorization



Monte Carlo neutrinos

Simulated 9508 fake events (thr=0.01)




Background rejection

<u>Cut</u>	<u>Events passing cut</u> (Run II, 163 days integrated livetime)	
1) Filter trigger	40 million	single-phone events
2) Electronic noise	25 million	singles
3) 5-phone coincidence	5 million	combinations
4) Waveform analysis	3 thousand	combinations
a) Periods < 4		
b) $20 \text{ kHz} < \text{freq} < 40 \text{ kHz}$		
c) Diamond metric < 0.7		
5) Threshold ≤ 0.024		
6) 5-phone localization	300	combinations
7) Threshold crossings < 2	0	combinations

([online](#), [offline](#))

Diffuse neutrino flux



...SAUND measurements coming soon

Conclusions

Lessons and Accomplishments

- Large data set collected (211 days integrated)
- Backgrounds determined
- Refraction significant (esp. beyond 1 km)
 - position + direction reconstruction nontrivial; sea-floor phones bad
- $c_{\text{sound}} = c_{\text{light}} / 200,000$!! (coincidence not as powerful)
- first-generation threshold: 10^{21} eV (10^{19} eV possible?)

Upcoming

- Rigorous efficiency checks
- Flux limits

Onward and downward (SAUND-II)

- On to more phones (52), effective area (250 km^2), online rejection (7 CPU $\tilde{\text{Q}}$)
- Down to the Gaussian floor

**A first effort has been made toward acoustic neutrino detection.
The method will be fleshed out by SAUND-II and others
in the next few years.**

<http://hep.stanford.edu/neutrino/SAUND>