

Richard L. Hallett, MD

Chief, Cardiovascular Imaging
Northwest Radiology Network
Indianapolis, IN


Clinical Assistant Professor – Radiology
Stanford University
Stanford, CA

PERIPHERAL CTA

RC 712B

5 December 2013

1630 - 1800



Outline

- CTA Acquisition Techniques
 - Scan Acquisition
 - Contrast Medium injection
 - Reconstruction
- Clinical Efficacy in PAD
- Cost Effectiveness

CTA Indications in PAD

- ✦ Intermittent Claudication
- ✦ Chronic Limb Ischemia
- ✦ Acute Ischemia (urgent)
- ✦ Monitoring of Tx (complications)

CTA Benefits / Limitations

- Non-invasive (DSA)
 - Spatial Resolution (MRA)
 - Quick Acquisition
 - (mostly) Operator Independent
- Ionizing Radiation
 - Nephrotoxic Contrast
 - Spatial Resolution (DSA)
 - No same session Tx
 - No flow or pressure measurements

CTA Acquisition

- ✦ Scan Acquisition
- ✦ Contrast Medium Injection

CTA Scan Acquisition

Peripheral CTA Scan Acquisition / Recon

Scanning Range 1

celiac artery (~T12) → toes
(105 – 130 cm)

Optional Scanning Range 2


above the knees → toes

Always pre-programmed, but only initiated
by RT if no contrast in pedal vessels

Recons:

Thin, overlapped

FOV = greater trochanters



Detector Configuration (mm)

TI / 360° (mm)

Table Speed (mm/s)

Scanning Time (s)

Anatomic coverage:
105 – 130cm

16-Channel MDCT

| | | | |
|---------|----|----|-------|
| 16×.75 | 18 | 36 | 30-40 |
| 16×.63 | 18 | 35 | 30-40 |
| 16×1.5 | 33 | 66 | 15-20 |
| 16×1.25 | 35 | 70 | 15-20 |

slow

~35 mm/s

slow

fast

~65 mm/s

fast

64-Channel MDCT

| | | | |
|--------|----|----|-------|
| 64×.63 | 55 | 92 | 11-14 |
| 64×.60 | 29 | 78 | 13-17 |

very

~85 mm/s

fast

Speed considerations for ≥ 64 slice CTA

- Outrunning Bolus
- Delayed filling of distal arteries


Free-Flap Planning CTA




Arteriomegaly



preprogrammed,
optional 2nd acquisition




Peripheral arterial bolus propagation



Fleischmann D and Rubin GD.
Radiology 2005, 1076-1082

Contrast Administration for peripheral CTA



INPUT





intravenous injection rate (mL/s)

Biphasic Injection

Phase I
(surge phase)



Biphasic Injection for Peripheral CTA

Fleischmann et al. *JVIR* 2006, 17(1) 3-26.





OUTPUT

arterial



Phase II
(continuing phase)

Patient Factors



- Arterial enhancement is inversely related to:
 - Cardiac output (CO)
 - Central blood volume (CBV) } usually unknown
 - CO (and CBV) correlate with body weight
 - at least in pts. with ~ normal cardiac function
 - Weight-based dosing helps consistency

1) Hittmair & Fleischmann, JCAT 2001

Integrated Contrast/Scan Protocol

- ✦ Simple, weight based injection volumes and flow rates, combined with a fixed scan time or scan time/diagnostic delay sum.
- ✦ automated bolus triggering
- ✦ Use physiology not scanner speed
- ✦ **BENEFITS:**
 - ✦ Decrease patient to patient variability in scan quality
 - ✦ Optimize imaging timing
 - ✦ Image all of the contrast given!
 - ✦ (Potentially) save contrast

STANFORD Integrated Scanning-Injection Protocol: (Siemens)

- Scan time: 40s for ALL patients (pitch variable)
- Inj.duration: 35s for ALL patients
- Delay: bolus triggering

weight

Biphasic Injection

| | |
|-------|------------------------------------|
| <55kg | 20 mL (4.0mL/s) + 96 mL (3.2mL/s) |
| <65kg | 23 mL (4.5mL/s) + 108 mL (3.6mL/s) |
| 75kg | 25 mL (5.0mL/s) + 120 mL (4.0mL/s) |
| >85kg | 28 mL (5.5mL/s) + 132 mL (4.4mL/s) |
| >95kg | 30 mL (6.0mL/s) + 144 mL (4.8mL/s) |


ST. VINCENT Integrated Scanning-Injection Protocol: (GE HD-750, VCT)

- Scan time: Variable (can't specify time)
- Add "diagnostic delay" to make 40 sec
- Inj.duration: 35s for ALL patients
- Delay: bolus triggering




| <u>weight</u> | <u>Biphasic Injection</u> |
|---------------|------------------------------------|
| <55 kg | 20 mL (4.0mL/s) + 96 mL (3.2mL/s) |
| 55-95 kg | 25 mL (5.0mL/s) + 120 mL (4.0mL/s) |
| >95 kg | 30 mL (6.0mL/s) + 144 mL (4.8mL/s) |

CTA Reconstruction and Interpretation


- Iterative Reconstruction
- **Recon** thin, overlapping images and **review in 3D**
 - VR / MIP overview then MPR, CPR
 - Axials in A/P
- Recon larger matrix



1024 Matrix Examples



1024 Matrix Examples



Efficacy of LE CTA in PAD

I.C.

CLI

PAD Classification

| Fontaine Stage | | Rutherford Classification |
|----------------|--|--|
| I | Asymptomatic | 0 |
| IIa | Mild Claudication (>200m walk) | 1 |
| IIb | Moderate to Severe Claudication (<200m walk) | 2 (moderate) 3 (severe) |
| III | Ischemic Rest Pain | 4 |
| IV | Ulceration or Gangrene | 5 (minor tissue loss) 6 (major tissue loss) |

CTA: Diagnostic Performance vs. DSA

Performance

| CT Channels | Sens (95% CI) | Spec (95% CI) |
|-------------|---------------|---------------|
| 2-4 | 92 (88-96) | 98 (95-99) |
| 16-64 | 97 (95-98) | 98 (96-99) |

Detection of $\geq 50\%$
Stenosis or Occlusion
By Anatomical Region


| Vessels | Sens (95% CI) | Spec (95% CI) |
|-----------------|---------------|---------------|
| Aortoiliac | 96 (91-99) | 98 (95-99) |
| Femoropopliteal | 97 (95-99) | 94 (85-99) |
| Trifurcation | 95 (85-99) | 91 (79-97) |

Met R et al. JAMA 2009;301:415-424


Diagnostic Performance: 64-slice CTA

- Symptomatic PAD: 242 pts, 7420 segments
- CTA and DSA performed
- For >70% stenosis:
 - SENS/SPEC 96% PPV 98% NPV 99%
 - No sig difference vs DSA findings
 - Results similar in Ca++ vs. Non-Ca++ lesions

The Achilles' Heel of Extremity CTA.....



Predictors of Vascular Calcification



- ✦ Above knee:¹ Severe PAD (Fontaine III-IV), Diabetes
- ✦ Below Knee:¹ Renal Failure (esp. dialysis), Diabetes
- ✦ Also:² Age, cardiac disease
- ✦ If heavy, significant decrease in SENS/SPEC in calf ¹

¹ Meyer BC Eur Radiol (2010) 20:497-505

² Ouwendijk R. Radiology (2006) 241, 603-608

CTA for stent assessment


- Most stents assessable (76%) by CTA
 - Gold / platinum markers
 - Motion
 - Strecker stent (Tantalum): Increased luminal density²
- If evaluable, **sens/spec ~ 95%** for significant in-stent restenosis (vs. DSA)

¹ Li X, et al. Eur J Radiol 2010; 98-103

² Strotzer, Invest. Radiol. 2001:36(11)



CTA for assessment of complications




Clinical Utility of LE CTA in PAD

- Intermittent Claudication (IC)
- Critical Limb Ischemia (CLI)

Intermittant Claudication (IC)

- Only 1/4 progress clinically
- Amputation uncommon (unless diabetic)
- More likely ilio-femoral than trifurcation dz



Buttock Claudication



Management of Intermittent Claudication by CTA

- Fontaine IIb patients, Tx decisions by TASC II criteria
- 57/58 correct Tx decision-making by CTA
 - One CFA stenosis missed
 - 29 endovasc/surg Tx
 - 29 conservative mgmt

Schernthaner R, et al. AJR 2007; 189:1215-1222

Critical Limb Ischemia (CLI)

- Duration > 2 weeks
- rest pain, tissue loss, ulcers, gangrene (TASC II)
 - Fontaine III / IV
- Higher incidence DM, trifurcation disease, comorbidities than IC



CTA assessment in CLI

- 41 pts, 1435 segments
- 64-CTA
- Fontaine IIb, III, IV
- 2.2% segments non-diagnostic
 - not included in calculation
 - 91% infrapop segments evaluable
- **For $\geq 50\%$ stenosis:**
 - **Sens 99%** **Spec 98%** **Acc: 98%**

Management Decisions in CLI


- 28 pts, Fontaine IV
- 64-detector CTA
- 14/28 → endovascular and/or surg. Tx
- correct decision-making for interventions, amputation, and medical Tx based on DSA and Tx response

Schernthaner R, et al. AJR 2009; 192: 1416-1424

Management by CTA in both IC and CLI

- Tx using TASC II guidelines
 - 49 conservative TX
 - 87 Endovascular
 - 38 surgery
 - 17 hybrid
- Tx recommendations from CTA same as DSA in all but ONE

Napoli A. Radiology. 2011 Dec 1;261(3):976-86.



Cost-Effectiveness of CTA

CTA as cost-effective care

- ✦ 2005: Randomized, controlled trial: 4-DCT vs DSA¹
 - ✦ Dx confidence slightly lower with CTA (calcifications)
 - ✦ CT cost-effective and provides sufficient information for Tx planning
 - ✦ DSA costs (564 Euro) greater than CTA (363)

¹ Kock, MC, et al. Radiology 2005. 237 (2) pp. 727-37

CTA as cost-effective care

- 2005: RCT - 156 pts CTA vs MRA
- CTA/MRA utility similar
- CTA lower diagnostic costs /patient
 - Average costs: \$199 vs \$627
 - Difference from imaging test itself, not from additional procedures

CTA as cost-effective care

- 2008: DIPAD Trial (Multicenter RCT)
- 514 PAD pts, randomized to Doppler/MRA/CTA
- CTA and MRA:
 - significantly higher diagnostic confidence
 - less additional imaging needed
- Total costs higher for MRA and US

Conclusions

- LE CTA is an accurate and cost-effective tool for assessment of various forms of peripheral arterial disease
- Implementation of integrated CM/scan protocol will improve consistency
- Clinical integration and uses will continue to expand



Thanks for your Attention!

- Special thanks to.....
 - Dominik Fleischmann, MD