

# Rotation Curve Modeler

## A Modeling and Simulating Tool for Arbitrary Galaxies

Robert J. Moss\* and Dr. James G. O'Brien†

*Department of Computer Science and Networking\**, *Department of Applied Math and Sciences†*  
*Wentworth Institute of Technology, Boston, Massachusetts 02115*  
*mossr@wit.edu\**, *obrienj10@wit.edu†*

### Abstract

In order to best understand the current dark matter paradigm, one must have a firm understanding of the rotation curve problem, as well as the underlying theories which explain the rotation curves themselves. Also in modern literature, many alternative gravitational theories are appearing with more frequency and greater relevance. Although specific galaxies can be modeled in a fairly straight forward manner (depending on the assumed theory), it may prove more difficult to compare various models on a particular galactic rotation data set. Moreover, it can prove difficult to sift through peer-reviewed articles to collect data and parameter sets that astronomers have published on various galaxies. Here, we present potential solutions to these problems. First, a Rotation Curve Modeler (RoCM) to model arbitrary galaxies. Second, a Scholarly Observed Celestial Measurements (SOCM) database to house all the galactic data in a central location. Finally, a Rotation Curve Simulation (RoCS) to simulate the spin of star clusters around the center of a galaxy.

### Keywords

*galaxy, rotation curve, modeling, simulating, general relativity, lambda cold dark matter, conformal gravity.*

## 1. Introduction

To model the rotation of single galaxies, astrophysicists would use programs like MATLAB or Mathematica, but there doesn't exist a singular tool to expedite this process in a universal format. More importantly, there doesn't exist a tool which can simultaneously model galaxies across competing gravitational theories. In this paper, we introduce the Rotation Curve Modeler (RoCM) to serve as this tool to model the rotation of star clusters around the center of a galaxy. The tool will also aid theorists working to study alternate theories to dark matter, by its ability to test all existing galactic models against the observational data for the specified galaxy simultaneously. With observable data from astronomers as the input, any arbitrary galaxy can be imported into the tool. The RoCM tool is also designed to be dynamic, and enable users to import their own galactic models to test against existing theories. Perhaps the most useful and unique features of RoCM, are its parameter value sliders, which allow users to control free fitting parameters within the models with real-time visual feedback in the generated graph. This way, each model can be finely tuned to the data, in a real time method while still yielding all of the same statistical measurements of single run galaxy modelers. The modeler makes no bias against the physicality of the specific model, but can thus be used as a full range exploration of the parameter space in question. RoCM is designed to be a stand alone program which can import data sets from astronomical papers, but has also been designed to work with the second phase of the project, the Scholarly Observed Celestial Measurements database (SOCM). The functionality of both RoCM and SOCM will be discussed in the body of the paper. First however, we briefly introduce the history of rotation curves in order to better explain the impact of this work.

## 2. The Rotation Curve Problem

The physicist Fritz Zwicky working in 1933 for the California Institute of Technology observed the red shift of star clusters within galaxies and stated that the expected velocity was entirely off pace with prediction. The proposed theory was that a matter invisible to the eye – dark matter – was in need to account for missing mass that would otherwise hold the clusters together [2]. He proposed that there were “faint galaxies and diffused gas” that filled in this void [2]. As telescopes moved from optical to radio, the fidelity was higher when measuring the amount of gas thanks to the success of spectroscopy. Although we can now account for this “missing mass”, many years of evidence has shown that the amount from gas was no where near the volume that was missing in the overall observation.

Later in the 1970's, Vera Rubin and her colleague, Kent Ford, worked with a new spectrograph that could observe the rotation velocity of spiral-galaxies with incredible accuracy in both the optical and radio bands [4]. Upon further research, they concluded that the stars were moving with a uniform velocity around the center of the galaxies, rather than gradually declining in velocity further out (as predicted by general relativity). Moreover, as their observations advanced, the uniform velocity seemed to trend across all sizes and shapes of galaxies, which would rule out effects of star formation and or galactic evolution. Their

findings brought along much speculation. Since at its heart, the predicted velocity  $v$  for the star clusters is a function of only two observational parameters, the distance from the galactic center and the mass of the galaxy. It was realized that if there was invisible mass in the outer regions of galaxies, the prediction and data could be reconciled. This missing factor that increased the predicted velocity is highly thought to be dark matter, yet as can be seen in current literature, with the appearance of many competing alternative gravitational theories, the true answer still remains a mystery. The rotation curve problem has been around for almost a century. Since it's origin, theories have surfaced that claim to solve it. RoCM was built for the purpose of comparing the theories that attempt to solve the “missing mass” problem in galaxy rotation curves.

### 3. Parameter Fitting for Models in RoCM

Each galaxy has a common set of parameters that may be used as model inputs:

Parameter	Units
Distance	Mpc
Luminosity	$10^{10} L_{\odot}$
Scale Length	kpc
Luminous Disk Mass	$10^{10} M_{\odot}$
Hydrogen Mass	$10^{10} M_{\odot}$

A quick summary is as follows: *Distance* is the assumed distance from the observed galaxy to us here on earth making the measurements. It should be noted that distance is the most important input parameter for galactic rotation curves, as all of the other inputs in the above table are quantities derived from the assumed distance. *Luminosity* is the assumed luminosity of the observed galaxy. *Scale Length* is the observed luminosity profile exponential falloff, assuming that galactic density is given by

$$\Sigma(R) = \Sigma_0 e^{-\frac{R}{R_0}} \quad (1)$$

where  $R_0$  is the scale length, and  $\Sigma_0$  is the central density and is thus ultimately related to the later described parameter  $N^*$ . *Luminous Disk Mass* is the amount of luminous matter in the galaxy, and *Hydrogen Mass* is the assumed mass of the hydrogen gas present in the galaxy. Due to the nature of the measurements, astronomers always provide a range for observed parameters. Each galaxy has an estimated range for the number of stars,  $N^*$ , which is calculated as a ratio of the luminous disk mass over solar mass,

$$N^* = \frac{M_{disk}}{M_{\odot}}. \quad (2)$$

Hence, in the workstation, RoCM provides a measurement for the mass to light ratio for the current galaxy, which can then be compared to the physical bounds as set by the morphology of the galaxy.

#### 3.1. Statistical $\chi^2$ Variance

To check the goodness of fit for the models used in RoCM, we've implemented a  $\chi^2$  variance. The  $\chi^2$  variance, unlike the traditional method, takes the measured error into account when calculating the fit. Using the observational data error,  $\mu$ , the model can be tested with

$$\chi^2 = \sum \frac{(O - E)^2}{\mu^2} \quad (3)$$

where  $O$  is the observed data and  $E$  is the experimental/model data. The objective is to minimize the  $\chi^2$  by adjusting the free parameters in the tested model. More details will be discussed in Section 4 on how RoCM can be used to quickly minimize the variance in the models.

#### 3.2. General Relativity Prediction

Referenced from [3], the normalization free parameter,  $N^*$ , scale length,  $R_0$ , Schwarzschild radius,  $\beta^*$ , and the speed of light,  $c$ , are used in general relativity (GR) to model a galaxy's rotation velocity as a function of galactocentric distance,  $R$ , in the form

$$v_{GR}(R) = \sqrt{\frac{N^* \beta^* c^2 R^2}{2R_0^3}} F_b \quad (4)$$

where the bessel functions  $I_0$ ,  $I_1$ ,  $K_0$ , and  $K_1$  are used to formulate the curve

$$F_b = \left[ I_0 \left( \frac{R}{2R_0} \right) K_0 \left( \frac{R}{2R_0} \right) - I_1 \left( \frac{R}{2R_0} \right) K_1 \left( \frac{R}{2R_0} \right) \right]. \quad (5)$$

It is assumed that each parameter is fixed (other than the input  $R$ ). Since these fixed parameters have an accepted range as dictated by the astronomer, it's difficult to alter them and see the behavior of the model in a single run program. Later we shall show how RoCM gives the user the flexibility to alter these fixed parameters within their acceptable ranges to best see their impact across multiple models of galactic rotation.

### 3.3. Lambda Cold Dark Matter Prediction

An issue when modeling galaxies is how to best adjust a free parameter like  $N^*$ . Some models have more free parameters than others, all dictated by the theory. The Lambda Cold Dark Matter ( $\Lambda$ CDM) theory states that each galaxy has two unobservable parameters, namely the spherical dark matter density,  $\sigma_0$ , and the dark matter halo radius,  $r_0$ . As described by [3], the  $\Lambda$ CDM rotation velocity contribution

$$v_{dark}(R) = \sqrt{4\pi\beta^*c^2\sigma_0 \left[ 1 - \frac{r_0}{R} \arctan \left( \frac{R}{r_0} \right) \right]} \quad (6)$$

is summed with the general relativity contribution to produce a total dark matter rotation velocity of

$$v_{total}(R) = \sqrt{v_{GR}^2 + v_{dark}^2}. \quad (7)$$

In order for  $\Lambda$ CDM to work,  $\sigma_0$  and  $r_0$  have to be fit to the data using a  $\chi^2$  test against  $v_{total}$ . The two free parameters effectively shape the modeled rotation curve to make it precisely fit the data. The problem with parameter fitting is that it's time consuming and can be tough to initialize minimum and maximum ranges without any empirical evidence to suggest an initial value range. To expedite this process, RoCM provides parameter fitting sliders to quickly and visually test each parameter value in the models, thus allowing the user to have a true interaction with the model on a real time basis.

### 3.4. Alternative Gravitational Models

Since one of the goals of RoCM is to compare models of rotation curves on the fly, we shall without loss of generality use the alternative gravitational theory known as conformal gravity (CG) to illustrate the scope of RoCM. Starting with the principle equations of GR, conformal gravity intends to replace GR, but allows it to stay true under certain scales [3]. Not intended to solve the rotation curve problem, conformal gravity seeks to formulate an equally good theory of gravity that is more inclusive than GR. Conformal gravity retains a metric theory of gravity but also includes the feature of conformal invariance. In short, conformal gravity requires three additional terms to the GR equation to be appended,  $\gamma^*$ ,  $\gamma_0$ , and  $\kappa$  [3]. These constants include missing feasible physical parameters that Einstein didn't need in order for him to model the solar system. Now that physicists need to model objects at a greater scale, the parameters that were initially scrapped from the theory of general relativity have now been added back in. As described by [3], the final rotation velocity function derived from conformal gravity is given by

$$v_{CG}(R) = \sqrt{v_{GR}^2 + \frac{N^*\gamma^*c^2R^2}{2R_0} I_1 \left( \frac{R}{2R_0} \right) K_1 \left( \frac{R}{2R_0} \right) + \frac{\gamma_0c^2R}{2} - \kappa c^2R} \quad (8)$$

where the constants are

$$\begin{aligned} \gamma^* &= 5.42 \times 10^{-41} \text{ cm}^{-1}, \\ \gamma_0 &= 3.06 \times 10^{-30} \text{ cm}^{-1}, \\ \text{and } \kappa &= 9.54 \times 10^{-54} \text{ cm}^{-2}. \end{aligned}$$

These small terms will contribute only at certain scales. This allows conformal gravity to scale down to the solar system, making the three additional terms negligible. GR has the ability to scale down to obey Newtonian physics. Thus, by transitivity, conformal gravity encompasses not only GR, but Newtonian laws of motion as well.

### 3.5. Bulge contribution

In order to complete the RoCM tool to include all other effects that most rotation curve modeling encompasses, it is important to include a bulge contribution. This formula from [3] uses the number of stars in the bulge,  $N_b^*$ , and bulge scale length,  $t$ , to derive the bulge contribution of

$$v_{bulge}(R) = \sqrt{\frac{2N_b^* \beta^* c^2}{\pi R} \int_0^{R/t} dz z^2 K_0(z)}. \tag{9}$$

Not every galaxy includes a bulge, thus RoCM provides a means to turn on and off the bulge contribution. This again allows the user to explore all possible options in modeling a galaxy while keeping true to physical observations (for example, some galaxies are noted to have a small but possibly insignificant bulge).

## 4. RoCM Functionality

RoCM seeks to bring together the equations described above, and create a workstation for astrophysicists to quickly and easily model galaxies. The tool consists of 7 major components that all contribute to the powerful functionality we've provided in RoCM:

- 1) SOCM drop-down table for easy access to the galaxy database (to select galaxies to plot or download).
- 2) Rotation velocity over distance graph (Vel vs. R) to plot models against collected data.
- 3) Parameter sliders to manipulate ALL values within galactic models (for parameter fitting purposes).
- 4) Workstation for users to import custom models and parameters outside the ones already described.
- 5) Simulation of the rotation curve in an animated visualization (Rotation Curve Simulation – RoCS).
- 6) LaTeX equation viewer for users to import their models in LaTeX format (to better understand the behavior of parameters within each model).
- 7) Exportable SVG of the plotted graph for use in articles or presentations.

### 4.1. Scholarly Observed Celestial Measurements

Researchers must first sift through peer-reviewed articles and gather galactic data one-by-one before modeling a galaxy. To minimize the time astrophysicists gather data, SOCM was created. SOCM is a public database that serves as a central repository for galactic parameters and observed velocity data. The database can also be used as an application programming interface (API) for researchers. Initially, SOCM is comprised of 112 galaxies, including the peer-reviewed galactic parameters and measurements of star velocities contained within. Although we are housing all of this information in one central location, all proper citations and references have still been maintained for proper use by theorists of the SOCM tool.

Scholarly Observed Celestial Measurements												
Display 5 galaxies											Search: <input type="text"/>	
(galaxy_name) Galaxy	(galaxy_type) Type	(distance) Distance (Mpc)	(luminosity) L <sub>B</sub> (10 <sup>10</sup> L <sub>⊙</sub> W)	(scale_length) R <sub>0</sub> (kpc)	(mass_hydrogen) M <sub>H</sub> (10 <sup>10</sup> M <sub>⊙</sub> kg)	(mass_disk) M <sub>disk</sub> (10 <sup>10</sup> M <sub>⊙</sub> kg)	(r_last) R <sub>last</sub> (kpc)	(mass_light_ratio) (M/L) <sub>stars</sub> (M <sub>⊙</sub> /L <sub>⊙</sub> kg/W)	(universal_constant) (v <sup>2</sup> /c <sup>2</sup> R) <sub>last</sub> (10 <sup>-30</sup> cm <sup>-1</sup> )	(velocities_count) Number of Observed Points	(citation_ids_array) Citations	Functions
MILKY-WAY	HSB	0.0081	1.6	2.56	0.43	5.527	100.72	3.45	0.422	635	Citations	Plot CSV
NGC-2403	HSB	4.3	1.647	2.7	0.46	2.37168	23.904	1.44	2.89	288	Citations	Plot CSV
NGC-6946	HSB	6.9	3.732	2.9	0.57	6.26976	22.38	1.68	6.387	207	Citations	Plot CSV
NGC-5055	HSB	9.2	3.622	2.9	0.76	6.77314	44.38	1.87	2.363	199	Citations	Plot CSV
NGC-2841	HSB	14.1	4.742	3.5	0.86	19.53704	51.611	4.12	5.831	141	Citations	Plot CSV

Showing 1-5 of 112 galaxies Previous 1 2 3 4 5 ... 23 Next

Figure 1: The user can sort by value, or search within the SOCM table to select a galaxy they would like to plot. The user can also download the parameter and velocity data of the galaxy in CSV format. Each galaxy has a set of associated citations to reference the origin of the data. Excluding the Milky Way, all of the data included in SOCM came from [3].

The idea is for astronomers to submit new measurements to the SOCM administrator (the authors) for approval. Thus creating a single means of distributing validated measurements of hundreds of galaxies. The input data can then be sent to RoCM. This way, if new galaxies get submitted and accepted, RoCM will always have the most up-to-date data that is provided. As an added feature, SOCM was created to also serve as a simple central database. Thus, researchers may pull from the SOCM API to use the data for applications other than rotation curve modeling. When data and parameters are pulled from SOCM outside of RoCM, all of the proper units as well as citations are preserved.

## 4.2. Curve Plot for Rotation Velocity

The main focus of RoCM is the curve plotting tool. Each galactic model is provided in the legend and can be displayed if selected. The Milky Way’s rotation curve can be seen in Figure 2. To see a full screen-shot of the main RoCM page, see Figure 9. It should be noted that although the displayed variance below for the Milky Way may appear high, in the Milky Way data set [5] there are over 600 points, and some of which have almost zero error, which makes our choice of error analysis seem exaggerated for this galaxy. This highlights the impact of RoCM to allow visual feedback from parameter adjustments. Included below the curve plotter is a table of the current parameter values. The values in the table can be directly edited from the input text-box or from the parameter sliders drop-down (shown in Figure 4). The graph can be exported as a loss-less SVG image, to easily include in an article or presentation.




Figure 2: The plotted GR, CG, total  $\Lambda$ CDM models over the observational data for the Milky Way using the collected data from [1] and [5].

GR $\chi^2$	CONFORMAL $\chi^2$	TOTAL $\chi^2$	Distance (Mpc) (distance)	$L_B$ ( $10^{10} L_\odot W$ ) (luminosity)	$R_0$ (kpc) (scale_length)	$M_{HI}$ ( $10^{10} M_\odot$ kg) (mass_hydrogen)	$M_{disk}$ ( $10^{10} M_\odot$ kg) (mass_disk)	$(M/L)_{stars}$ ( $M_\odot/L_\odot$ kg/W) (mass_light_ratio)
$4.28 \times 10^5$	$3.63 \times 10^5$	$1.14 \times 10^6$	$8.10 \times 10^{-8}$	1.6	2.56	0.43	5.527	3.45

Figure 3: The  $\chi^2$  table and mutable parameters table for the Milky Way galaxy.

### 4.3. Parameter Fitting Sliders

The power of RoCM lies within these parameter sliders. The user can adjust the minimum and maximum of the parameters and slide the bar to see how the changes effect the curves on the plot. This responsive visualization helps aid in understanding how each parameter behaves in the theories that use it. Since the sliders are built into the workstation, one can see the effect they have on the enabled curves immediately. When writing a user defined model, the user can add their own parameters to be manipulated via the sliders.




Figure 4: Parameter fitting sliders with min and max values, units, and optional bulge contribution toggle.

### 4.4. User Defined Models and Parameters

The discussed models, namely GR,  $\Lambda$ CDM, and CG aren't the only theories that model galaxies. Thus, users have the ability to write a model in JavaScript to import directly into RoCM. The User Defined Model workbench opens RoCM up for any and all models that want to be compared against each other. The function should be in the form:

```
function MODELNAME (R) {
  var rotation_velocity;
  // The implemented JavaScript model

  return rotation_velocity;
}
```

On the actual RoCM tool, user tutorials are provided to allow for quick input of the suggested models. As RoCM grows in scope, we plan on having all of the major competing theories already hard coded into RoCM.

**4.4.1. Constants and Conversions:** Since it is the goal of RoCM to allow for any model, we must then also allow the user to manipulate some of the hard coding of RoCM as well. Thus, constants can be changed if needed (see for example some alternative gravitational theories based on  $G$  not being constant). The constants can be retrieved and manipulated based on the following CONST commands. We also have provided some useful and common conversions hard coded into RoCM that can be accessed via the CONVERT commands.

```
// Speed of light
CONST.get ("c")
CONST.get ("speed_of_light")

// Solar mass
CONST.get ("Msun")
CONST.get ("solar_mass")

// Solar luminosity
CONST.get ("Lsun")
CONST.get ("solar_luminosity")

// Gravitational constant
CONST.get ("G")
CONST.get ("gravitational_constant")

// Schwarzschild radius
CONST.get ("Rs")
CONST.get ("schwarzschild_radius")

CONVERT.pc_to_kpc(pc)
CONVERT.Mpc_to_kpc(Mpc)
CONVERT.kpc_to_Mpc(kpc)
CONVERT.kpc_to_pc(kpc)
CONVERT.kpc_to_km(kpc)
CONVERT.km_to_kpc(km)
CONVERT.km_to_m(km)
CONVERT.km_to_cm(km)
CONVERT.cm_to_km(cm)
CONVERT.cm_to_kpc(cm)
// GeV/cm^3 to kg/km*s^2
CONVERT.GeVcm3_to_kgkms2(GeVcm3)
CONVERT.arcsec_to_degree(arcsec)
CONVERT.degree_to_arcsec(degree)
CONVERT.radian_to_degree(radian)
CONVERT.degree_to_radian(degree)
```

## 4.5. Rotation Curve Simulation

Although theorists in the field of gravity are quite fluent in rotation curve discussions, to an audience that has never been exposed to the rotation curve problem, their use and influence on the current paradigm may be elusive. RoCM has a built in function in the workstation to simulate and animate the current selected model. The Rotation Curve Simulation (RoCS) compares a particular model to the observed data by animating their spins side by side. The color scale represents the relative **minimum** and **maximum** rotation velocity for the stars (as dictated by the y axis of the data). The scale helps recognize when the rotation curve simulation of a model doesn't match up with the observational data. The user can then compare the differences in rotational speed by both the difference in color scale, and by looking at the spinning animation in RoCS.




Figure 5: When viewing a model in RoCS side by side with the simulated observational data, it's clear to see where the velocity inconsistencies arise within the modeled galaxies. The biggest difference is when the data is compared to the GR prediction for NGC-2403 using the parameters provided by the astronomers. The inner parts of the galaxy seems to match the data. But about half way out, the velocity starts to slow down as described by the GR prediction, while the data stays uniform.




Figure 6: As an alternative, the CG prediction for the provided NGC-2403 parameters clearly matches the observational data.




Figure 7: The NGC-2403 galaxy predicted with GR and CG plotted within RoCM with the parameters given by the astronomers. This illustrates that the GR curve starts to gradually decline as the CG curve stays consistent with the observational data, without fitting the parameters.




Figure 8: The parameters given by the astronomers have been fitted within an acceptable range to minimize the  $\chi^2$  variance for CG. A bulge contribution was also added to each model. All this was achieved through the parameter sliders functionality. This can be compared to the legacy method of model fitting as demonstrated by [3].



## 5. Conclusion

In this project, we have successfully created a robust tool that can model the rotation curves of galaxies in a quite general and free form manner. Since the RoCM tool can be manipulated by the user, it serves currently as both a complete tool for individual modeling as well as model comparison. With the creation of SOCM, the project now becomes diverse in catering to researchers seeking data alone, or model fitting. After its implementation and some outreach by the authors, the hope is that astronomers will use SOCM to upload observational data into one central location. Theorists then can use RoCM to test that data against several galactic models to finally understand the dynamics of galaxies.

We are pleased to announce that SOCM is open to the public at [socm.herokuapp.com](http://socm.herokuapp.com) (shortly transitioning to [wit.edu/socm](http://wit.edu/socm)), where users can now view our database of collected measurements and developers may utilize our API endpoints to use in their own endeavors. RoCM is hosted online and can be found at [rotationcurve.herokuapp.com](http://rotationcurve.herokuapp.com) (shortly transitioning to [wit.edu/rotationcurve](http://wit.edu/rotationcurve)). Each project individually is a contribution to the open source community and can be found here: [github.com/RoCMSOCM](https://github.com/RoCMSOCM).

## Acknowledgments

The authors would like to thank the departments of Sciences and Computer Science at WIT for allowing the work on RoCM and SOCM as a senior capstone project. The authors would like to thank Crystal Bailey, APS and SPS for the generous travel grant for R. J. Moss to present this work at the 2014 April APS meeting. The authors would also like to thank Chuck Hotchkiss and Paloma Valverde for their continued support for undergraduate research projects at WIT, and for a matching travel grant for dissemination of this work. The authors would also like to thank Dr. Sophia Cizneros for her input into the creation of a useful database for theorists.

We would also like to thank Professor Franz Rueckert, Patrick McGee and David Miller for their help building the SOCM database. Alex Clement, Professor Lisa MacLean, and Professor Mohammed Anwaruddin have also helped shape the outcome of the RoCM tool, so we thank them for their support.

## References

- [1] P. Bhattacharjee, S. Chaudhury, and S. Kundu. Rotation Curve of the Milky Way out to  $\sim 200$  kpc. *Astrophysical Journal*, 785:63, Apr. 2014.
- [2] O. Knill. Bibliographical entries of Jost Buergi, Max Wolf, Arthur Schuster and Fritz Zwicky. *Springer*, 2007.
- [3] P. D. Mannheim and J. G. O'Brien. Fitting galactic rotation curves with conformal gravity and a global quadratic potential. *Physical Review D*, 85(12):124020, June 2012.
- [4] V. C. Rubin, W. K. J. Ford, and N. Thonnard. Rotational properties of 21 SC galaxies with a large range of luminosities and radii, from NGC 4605  $R = 4$  kpc to UGC 2885  $R = 122$  kpc. *Astrophysical Journal*, 238:471–487, June 1980.
- [5] Y. Sofue, M. Honma, and T. Omodaka. Unified Rotation Curve of the Galaxy – Decomposition into de Vaucouleurs Bulge, Disk, Dark Halo, and the 9-kpc Rotation Dip. *Publications of the ASJ*, 61:227–, Feb. 2009.




Figure 9: RoCM’s main page where users can plot the specified galaxy (the Milky Way galaxy is shown). The User Defined Model workbench provides a way to import a JavaScript function to locally run additional models.